Product Datasheet

TNF RI/TNFRSF1A Antibody - BSA Free NBP2-78785

Unit Size: 0.1 mg

Store at 4C for up to 3 months. For longer storage, aliquot and store at -20C.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-78785

Updated 9/9/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-78785



NBP2-78785


TNF RI/TNFRSF1A Antibody - BSA Free

TNF RI/TNFRSF1A Antibody - BSA Free	
Product Information	
Unit Size	0.1 mg
Concentration	1.0 mg/ml
Storage	Store at 4C for up to 3 months. For longer storage, aliquot and store at -20C.
Clonality	Polyclonal
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	PBS
Product Description	
Description	Novus Biologicals Rabbit TNF RI/TNFRSF1A Antibody - BSA Free (NBP2-78785) is a polyclonal antibody validated for use in IHC and WB. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	7132
Gene Symbol	TNFRSF1A
Species	Human, Mouse
Immunogen	Partial recombinant protein made to an N-terminal portion of the human TNFR1 protein (amino acids 2-211) [UniProt P19438]
Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry
Recommended Dilutions	Western Blot 1 - 5 ug/ml, Immunohistochemistry 1:100 - 1:500, Immunohistochemistry-Paraffin 1:100 - 1:500




Images

Western Blot: TNF RI/TNFRSF1A Antibody [NBP2-78785] - Total protein from human and mouse liver and placenta was separated on a 12% gel by SDS-PAGE, transferred to PVDF membrane and blocked in 5% non-fat milk in TBST. The membrane was probed with 2.0 ug/ml anti-TNFR1 in blocking buffer and detected with an anti-rabbit HRP secondary antibody using West Pico PLUS chemiluminescence detection reagent.



Immunohistochemistry-Paraffin: TNF RI/TNFRSF1A Antibody [NBP2-78785] - Analysis of a FFPE tissue section of human small intestine using 1:200 dilution of TNFRI antibody (NBP2-78785). The signal was developed using HRP-DAB method which followed counterstaining of the cells with hematoxylin. The antibody generated mainly moderately strong reactivity in glandular cell cytoplasm.



Procedures

Western Blot protocol for TNF RI/TNFRSF1A Antibody (NBP2-78785)

TNF RI/TNFRSF1A Antibody:

Western Blot Protocol

- 1. Perform SDS-PAGE on samples to be analyzed, loading 10-25 ug of total protein per lane.
- 2. Transfer proteins to PVDF membrane according to the instructions provided by the manufacturer of the membrane and transfer apparatus.
- 3. Stain the membrane with Ponceau S (or similar product) to assess transfer success, and mark molecular weight standards where appropriate.
- 4. Rinse the blot TBS -0.05% Tween 20 (TBST).
- 5. Block the membrane in 5% Non-fat milk in TBST (blocking buffer) for at least 1 hour.
- 6. Wash the membrane in TBST three times for 10 minutes each.
- 7. Dilute anti-TNFRI primary antibody in blocking buffer and incubate overnight at 4C with gentle rocking.
- 8. Wash the membrane in TBST three times for 10 minutes each.
- 9. Incubate the membrane in diluted HRP conjugated secondary antibody in blocking buffer (as per manufacturer's instructions) for 1 hour at room temperature.
- 10. Wash the blot in TBST three times for 10 minutes each (this step can be repeated as required to reduce background).
- 11. Apply the detection reagent of choice in accordance with the manufacturers instructions.

Immunohistochemistry-Paraffin protocol for TNF RI/TNFRSF1A Antibody (NBP2-78785)

TNF RI/TNFRSF1A Antibody:

Immunohistochemistry-Paraffin Embedded Sections

Antigen Unmasking:

Bring slides to a boil in 10 mM sodium citrate buffer (pH 6.0) then maintain at a sub-boiling temperature for 10 minutes. Cool slides on bench-top for 30 minutes.

Staining:

- 1. Wash sections in deionized water three times for 5 minutes each.
- 2. Wash sections in wash buffer for 5 minutes.
- 3. Block each section with 100-400 ul blocking solution for 1 hour at room temperature.
- 4. Remove blocking solution and add 100-400 ul diluted primary antibody. Incubate overnight at 4 C.
- 5. Remove antibody solution and wash sections in wash buffer three times for 5 minutes each.
- 6. Add 100-400 ul biotinylated diluted secondary antibody. Incubate 30 minutes at room temperature.
- 7. Remove secondary antibody solution and wash sections three times with wash buffer for 5 minutes each.
- 8. Add 100-400 ul Streptavidin-HRP reagent to each section and incubate for 30 minutes at room temperature.
- 9. Wash sections three times in wash buffer for 5 minutes each.
- 10. Add 100-400 ul DAB substrate to each section and monitor staining closely.
- 11. As soon as the sections develop, immerse slides in deionized water.
- 12. Counterstain sections in hematoxylin.
- 13. Wash sections in deionized water two times for 5 minutes each.
- 14. Dehydrate sections.
- 15. Mount coverslips.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP2-78785

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

NBP2-35274-5ug Recombinant Human TNF RI/TNFRSF1A Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-78785

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

