Product Datasheet

DPPIV/CD26 Antibody (JM11-42) NBP2-78791

Unit Size: 100 ul

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-78791

Updated 5/28/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-78791



NBP2-78791

DPPIV/CD26 Antibody (JM11-42)


- · · · · · · · · · · · · · · · ·	
Product Information	
Unit Size	100 ul
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	JM11-42
Preservative	0.05% Sodium Azide
Isotype	IgG
Purity	Protein A purified
Buffer	TBS (pH7.4), 0.05% BSA, 40% Glycerol
Product Description	
Host	Rabbit
Gono ID	1803

Product Description	
Host	Rabbit
Gene ID	1803
Gene Symbol	DPP4
Species	Human, Mouse, Rat
Immunogen	Synthetic peptide within Human DPPIV/CD26 aa 701-747 / 766 (Extracellular). (SwissProt: P27487 Human; SwissProt: P28843 Mouse; SwissProt: P14740 Rat)


Product Application Details	
Applications	Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Knockout Validated
Recommended Dilutions	Western Blot 1:500-1:2000, Flow Cytometry 1:50-1:100, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1:50-1:200, Immunohistochemistry-Paraffin 1:50-1:100, Knockout Validated

Images


Western Blot: DPPIV/CD26 Antibody (JM11-42) [NBP2-78791] - Western blot analysis of DPPIV/CD26 on different lysates. Proteins were transferred to a PVDF membrane and blocked with 5% BSA in PBS for 1 hour at room temperature. The primary antibody (1/500) was used in 5% BSA at room temperature for 2 hours. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1:5,000 dilution was used for 1 hour at room temperature. Positive control: Lane 1: mouse lung tissue lysate Lane 2: mouse heart tissue lysate Lane 3: human kidney tissue lysate Lane 4: Raji cell lysate Lane 5: PC-3 cell lysate




Immunocytochemistry/Immunofluorescence: DPPIV/CD26 Antibody (JM11-42) [NBP2-78791] - Staining CD26 in Hela cells (red). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.




Immunohistochemistry-Paraffin: DPPIV/CD26 Antibody (JM11-42) [NBP2 -78791] - Analysis of paraffin-embedded human liver tissue using anti-CD26 antibody. Counter stained with hematoxylin.




Flow Cytometry: DPPIV/CD26 Antibody (JM11-42) [NBP2-78791] - Analysis of Hela cells with CD26 antibody at 1/50 dilution (red) compared with an unlabelled control (cells without incubation with primary antibody; black). Alexa Fluor 488-conjugated goat anti rabbit IgG was used as the secondary antibody.




Immunocytochemistry/Immunofluorescence: DPPIV/CD26 Antibody (JM11-42) [NBP2-78791] - Staining CD26 in PC-12 cells (red). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



Immunohistochemistry-Paraffin: DPPIV/CD26 Antibody (JM11-42) [NBP2 -78791] - Analysis of paraffin-embedded rat epididymis tissue using anti-CD26 antibody. Counter stained with hematoxylin.



Knockout Validated: DPPIV/CD26 Antibody (JM11-42) [NBP2-78791] - All lanes: Western blot analysis of CD26 with anti-CD26 antibody at 1:500 dilution. Lane 1: Wild-type Hela whole cell lysate (10 ug). Lane 2: CD26 knockout Hela whole cell lysate (10 ug). NBP2-78791 was shown to specifically react with CD26 in wild-type Hela cells. NO band was observed when CD26 knockout sample was tested. Wild-type and CD26 knockout samples were subjected to SDS-PAGE. Proteins were transferred to a PVDF membrane and blocked with 5% NFDM in TBST for 1 hour at room temperature. The primary antibody (1:500) was used in 5% BSA at room temperature for 2 hours. Goat Anti-Rabbit IgG-HRP Secondary Antibody (HA1001) at 1:300,000 dilution was used for 1 hour at room temperature.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP2-78791

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

NBP3-21319PEP DPPIV/CD26 Recombinant Protein Antigen

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-78791

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

