

Product Datasheet

Mesothelin Antibody (MSLN/2131) NBP2-79724-100ug

Unit Size: 100 ug

Store at 4C.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-79724

Updated 7/16/2024 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-79724



NBP2-79724-100ug


Mesothelin Antibody (MSLN/2131)

Product Information	
Unit Size	100 ug
Concentration	0.2 mg/ml
Storage	Store at 4C.
Clonality	Monoclonal
Clone	MSLN/2131
Preservative	0.05% Sodium Azide
Isotype	IgG2b Kappa
Purity	Protein A or G purified
Buffer	10 mM PBS with 0.05% BSA
Product Description	
Description	200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP2-79858) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.
Host	Mouse
Gene ID	10232
Gene Symbol	MSLN
Species	Human, Mouse, Rat
Marker	Mesothelial Marker
Immunogen	Recombinant fragment (around aa 273-407) of human Mesothelin (MSLN) protein (exact sequence is proprietary) (Uniprot: Q13421)
Product Application Details	
Applications	Immunohistochemistry, Immunohistochemistry-Paraffin, Protein Array
Recommended Dilutions	Immunohistochemistry 1-2 ug/ml, Immunohistochemistry-Paraffin 1-2 ug/ml, Protein Array
Application Notes	Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes. Optimal dilution for a specific application should be determined.




Images


Immunohistochemistry-Paraffin: Mesothelin Antibody (MSLN/2131) [NBP2-79724] - Formalin-fixed, paraffin-embedded human endometrial carcinoma stained with Biotin-conjugated Mesothelin Mouse Monoclonal Antibody (MSLN/2131).




Immunohistochemistry-Paraffin: Mesothelin Antibody (MSLN/2131) [NBP2-79724] - Formalin-fixed, paraffin-embedded human Lung Mesothelioma stained with Mesothelin Antibody (MSLN/2131).




Immunohistochemistry-Paraffin: Mesothelin Antibody (MSLN/2131) [NBP2-79724] - Formalin-fixed, paraffin-embedded human Endometrial Carcinoma stained with Mesothelin Antibody (MSLN/2131).




Immunohistochemistry-Paraffin: Mesothelin Antibody (MSLN/2131) [NBP2-79724] - Formalin-fixed, paraffin-embedded human Tonsil stained with Mesothelin Antibody (MSLN/2131).



Immunohistochemistry-Paraffin: Mesothelin Antibody (MSLN/2131) [NBP2-79724] - Formalin-fixed, paraffin-embedded Mouse Lung stained with Mesothelin Antibody (MSLN/2131).



Immunohistochemistry-Paraffin: Mesothelin Antibody (MSLN/2131) [NBP2-79724] - Formalin-fixed, paraffin-embedded Rat Stomach stained with Mesothelin Antibody (MSLN/2131).



Protein Array: Mesothelin Antibody (MSLN/2131) [NBP2-79724] - Analysis of Protein Array containing more than 19,000 full-length human proteins using Mesothelin Antibody (MSLN/2131). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt (TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-79724-100ug

HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
NBP1-43317-0.5mg	Mouse IgG2b Kappa Light Chain Isotype Control (MG2b)
NBP1-88224PEP	Mesothelin Recombinant Protein Antigen

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-79724

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

