Product Datasheet

Spectrin alpha 1 Antibody (SPTA1/1832) NBP2-79772-100ug

Unit Size: 100 ug Store at 4C.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-79772

Updated 7/16/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-79772



NBP2-79772-100ug


| Spectrin alpha 1 Antibody (SPTA1/1832) | |
|--|---|
| Product Information | |
| Unit Size | 100 ug |
| Concentration | 0.2 mg/ml |
| Storage | Store at 4C. |
| Clonality | Monoclonal |
| Clone | SPTA1/1832 |
| Preservative | 0.05% Sodium Azide |
| Isotype | IgG2b Kappa |
| Purity | Protein A or G purified |
| Buffer | 10 mM PBS with 0.05% BSA |
| Target Molecular Weight | 240 kDa |
| Product Description | |
| Description | 200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP2-79906) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. |
| Host | Mouse |
| Gene ID | 6708 |
| Gene Symbol | SPTA1 |
| Species | Human |
| Marker | Erythrocyte Marker |
| Immunogen | Recombinant human Spectrin alpha 1 fragment (around aa356-475) (exact sequence is proprietary) (Uniprot: P02549) |
| Product Application Details | |
| Applications | Western Blot, ELISA, Immunohistochemistry, Immunohistochemistry-Paraffin, Protein Array |
| Recommended Dilutions | Western Blot 1-2 ug/ml, ELISA 2-4 ug/ml, Immunohistochemistry 1-2 ug/ml, Immunohistochemistry-Paraffin 1-2 ug/ml, Protein Array |
| Application Notes | ELISA: Use Ab at 2-4ug/ml for coating. Order Ab without BSA. Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes. Optimal dilution for a specific application should be determined. |




Optimal dilution for a specific application should be determined.

Images


Western Blot: Spectrin alpha 1 Antibody (SPTA1/1832) [NBP2-79772] - Western Blot Analysis of Human HeLa cell lysate using Spectrin alpha 1 Antibody (SPTA1/1832).




Immunohistochemistry-Paraffin: Spectrin alpha 1 Antibody (SPTA1/1832) [NBP2-79772] - Formalin-fixed, paraffin-embedded human Tonsil stained with Spectrin alpha 1 Antibody (SPTA1/1832).




Western Blot: Spectrin alpha 1 Antibody (SPTA1/1832) [NBP2-79772] - Western Blot Analysis of Human K562 cell lysate using Spectrin alpha 1 Antibody (SPTA1/1832).



Immunohistochemistry-Paraffin: Spectrin alpha 1 Antibody (SPTA1/1832) [NBP2-79772] - Formalin-fixed, paraffin-embedded human Placenta stained with Spectrin alpha 1 Antibody (SPTA1/1832).



Protein Array: Spectrin alpha 1 Antibody (SPTA1/1832) [NBP2-79772] - Analysis of Protein Array containing more than 19,000 full-length human proteins using Spectrin alpha 1 Antibody (SPTA1/1832) Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5.







Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP2-79772-100ug

HAF007 Goat anti-Mouse IgG Secondary Antibody [HRP]

NB720-B Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
NBP1-43317-0.5mg Mouse IgG2b Kappa Light Chain Isotype Control (MG2b)

NBP1-87953PEP Spectrin alpha 1 Recombinant Protein Antigen

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-79772

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

