Product Datasheet

Endoglin/CD105 Antibody (MJ7/18) - Azide and BSA Free NBP2-80718

Unit Size: 0.5 mg

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-80718

Updated 9/9/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-80718



NBP2-80718

Endoglin/CD105 Antibody (MJ7/18) - Azide and BSA Free


3	
Product Information	
Unit Size	0.5 mg
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	MJ7/18
Preservative	No Preservative
Isotype	IgG2a Kappa
Purity	Protein G purified
Buffer	Tris-Glycine

Novus Biologicals Rat Endoglin/CD105 Antibody (MJ7/18) - Azide and BSA Free (NB100-77666) is a monoclonal antibody validated for use in IHC, WB, Flow, ICC/IF and IP. All Novus Biologicals antibodies are covered by our 100% guarantee.
Rat
2022
ENG
Mouse
Inflamed mouse skin

Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Flow Cytometry, Flow (Cell Surface), Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Frozen, Immunoprecipitation
Recommended Dilutions	Western Blot 1:100-1:2000, Flow Cytometry 1:10-1:1000, Immunohistochemistry 1:10-1:500, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation 1:10-1:500, Immunohistochemistry-Paraffin, Immunohistochemistry-Frozen 1:10-1:500, Flow (Cell Surface)
Application Notes	For immunofluorescent staining, the suggested use of this reagent is: 0.5 microgram per one million cells in 100 microliter volume. It is recommended that the reagent be titrated for optimal performance for each application.


Images

Western Blot: Endoglin/CD105 Antibody (MJ7/18) - Azide and BSA Free [NBP2-80718] - Total protein from mouse placenta was separated on a 12% gel by SDS-PAGE, transferred to PVDF membrane and blocking in 5% non-fat milk in TBST. The membrane was probed with 2 ug/ml anti-CD105 (NB100-77666) in 1% milk/TBST and detected with an anti-rat HRP conjugated secondary antibody using chemiluminescence. Image from the standard format of this antibody.







Immunocytochemistry/Immunofluorescence: Endoglin/CD105 Antibody (MJ7/18) - Azide and BSA Free [NBP2-80718] - Using the DyLight 550 direct conjugate Staining of CD105 (endoglin) in 4T1 tumors after radioimmunotherapy shows CD105 expression (red), blood vessels (green), and nuclei (blue).




Immunohistochemistry-Paraffin: Endoglin/CD105 Antibody (MJ7/18) - Azide and BSA Free [NBP2-80718] - Analysis of a FFPE mouse ductus deference using a 1:50 dilution of CD105 antibody (clone MJ7/18) on a Bond Rx autostainer (Leica Biosystems). The assay involved 20 minutes of heat induced antigen retrieval (HIER) using 10mM sodium citrate buffer (pH 6.0)



Flow (Cell Surface): Endoglin/CD105 Antibody (MJ7/18) - Azide and BSA Free [NBP2-80718] - A surface stain was performed on MS-1 Cells with Endoglin/CD105 (MJ7/18) antibody NB100-77666 (blue) and a matched isotype control (orange). Cells were incubated in an antibody dilution of 1 ug/mL for 20 minutes at room temperature, followed by rat F (ab)2 IgG (H+L) APC-conjugated secondary antibody (F0113, R&D Systems). Image from the standard format of this antibody.



Flow Cytometry: Endoglin/CD105 Antibody (MJ7/18) - Azide and BSA Free [NBP2-80718] - Analysis of Biotin conjugate of NB100-77666. Surface staining of bEND.3 cell line with 0.25 ug of Anti-Mouse CD105 (Endoglin) Biotin followed by Streptavidin PE. Appropriate isotype controls were used (open histogram). Total viable cells were used for analysis.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP2-80718

HAF005 Goat anti-Rat IgG Secondary Antibody [HRP]

F0105B Goat anti-Rat IgG Secondary Antibody [Phycoerythrin]
NBP1-43321-0.5mg Rat IgG2a Kappa Light Chain Isotype Control (R2a)
NBP1-91212PEP Endoglin/CD105 Recombinant Protein Antigen

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-80718

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

