Product Datasheet

c-Myc Antibody (SPM237) [Alexa Fluor® 350] NBP2-86682AF350

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-86682AF350

Updated 10/26/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-86682AF350



NBP2-86682AF350

c-Myc Antibody (SPM237) [Alexa Fluor® 350]

Product Information Unit Size 0.1 ml Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonality Monoclonal Clone SPM237 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate Alexa Fluor 350 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description 4609 Gene ID 4609 Gene Symbol MYC Species Human, Chicken (Negative), Mouse (Negative), Rat (Negative) Reactivity Notes Does not react with Mouse, Rat or Chicken. Species Human, Chicken (Negative), Mouse (Negative), Maithing as c-myc. Its epitope spans between aa 410-419 (EQKLISEEDL), which is a specific portion of an alpha helical region of human c-myc protein. This monoclonal antibody shows no differentiation and is amplified and/or overexpressed in a variety of tumors. Overexpression of c-myc protein incore of cores-reaction with -myc. c-myc is involved in the control of cell proliferation and differentiation and is amplified and/or overexpressed in a variety of tumors. Overexpression of c-myc protein increaction and differentiation and is amplified and/or overexpressed in a variety of tumors. Over		-
ConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneSPM237Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 350PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseGene ID4609Gene SymbolMYCSpeciesHuman, Chicken (Negative), Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse, Rat or Chicken.Specificity/SensitivityIt recognizes a transcription factor of 64-67kDa, identified as c-myc. Its epitope spans between aa 410-419 (EQKLISEEDL) which is a specific portion of an alpha helical region of human c-myc protein. This monoclonal antibody shows no cross-reaction with v-myc. c-myc is involved in the control of cell proliferation and differentiation and is amplified and/or overexpressed in a variety of tumors. Overexpressed of a variety of tumors. Overexpression of c-myc protein. This monoclonal antibody shows no cross-reaction with v-myc. c-myc is involved in the control of cell proliferation and differentiation and is amplified and/or overexpressed in a variety of tumors. Overexpression of c-myc protein. This monoclonal antibody shows no cross-reaction with v-myc. c-myc is involved in the control of cell proliferation and differentiation and is amplified and/or overexpressed in a variety of tumors. Overexpressed of a variety of tu	Product Information	
services. Storage Store at 4C in the dark. Clonality Monoclonal Clone SPM237 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate Alexa Fluor 350 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Mouse Gene ID 4609 Gene Symbol MYC Species Human, Chicken (Negative), Mouse (Negative), Rat (Negative) Reactivity Notes Does not react with Mouse, Rat or Chicken. Specificity/Sensitivity It recognizes a transcription factor of 64-67kDa, identified as c-myc. Its epitope spans between aa 410-419 (EQKLISEEDL) which is a specific portion of an alpha helical region of human c-myc protein. This monoclonal antibody shows no cross-reaction with v-myc. c-myc is involved in the control of cell proliferation and differentiation and is amplified and/or overexpressed in a variety of tumors. Overexpression of c-myc protein a variety of tumors. Overexpression of c-myc protein adverted of tumors. Overexpression of c-myc protein adverted of tumors. Overexpression of c-myc protein occurs frequently in luminal cells of prostate intraepithelial neoplasia as well as in most primary carcinomas and metastatic disease. Immunogen A synthetic peptide, corresponding to aa 408-439	Unit Size	0.1 ml
ClonalityMonoclonalCloneSPM237Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 350PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description4609HostMouseGene ID4609Gene SymbolMYCSpeciesHuman, Chicken (Negative), Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse, Rat or Chicken.Specificity/SensitivityIt recognizes a transcription factor of 64-67kDa, identified as c-myc. Its epitope trans a 410-419 (EQKLISEEDL) which is a specific portion of an alpha helical region of human c-myc protein. This monoclonal antibody shows no cross-reaction with v-myc. c-myc is involved in the control of cell proliferation and differentiation and is amplified and/or overexpressed in a variety of tumors. Over-expression of c-myc protein Occurs frequently in luminal cells of prostate intraepithelial neoplasia as well as in most primary carcinomas and metastatic disease.ImmunogenA synthetic peptide, corresponding to aa 408-439 (AEEQKLISEEDLLRKRREQLKHKLEQLRNSCA) from C-terminus of human c-	Concentration	
CloneSPM237Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 350PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID4609Gene SymbolMYCSpeciesHuman, Chicken (Negative), Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse, Rat or Chicken.Specificity/SensitivityIt recognizes a transcription factor of 64-67kDa, identified as c-myc. Its epitope spans between aa 410-419 (EQKLISEEDL) which is a specific portion of an alpha helical region of human c-myc protein. This monoclonal antibody shows no cross-reaction with v-myc. c-myc is involved in the control of cell proliferation and differentiation and is amplified and/or overexpressed in a variety of tumors. Over-expression of c-myc protein occurs frequently in luminal cells of prostate intraepithelial neoplasia as well as in most primary carcinomas and metastatic disease.ImmunogenA synthetic peptide, corresponding to aa 408-439 (AEEQKLISEEDLLRKRREQLKHKLEQLRNSCA) from C-terminus of human c-	Storage	Store at 4C in the dark.
Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 350PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID4609Gene SymbolMYCSpeciesHuman, Chicken (Negative), Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse, Rat or Chicken.Specificity/SensitivityIt recognizes a transcription factor of 64-67kDa, identified as c-myc. Its epitope spans between aa 410-419 (EQKLISEEDL) which is a specific portion of an alpha helical region of human c-myc protein. This monoclonal antibody shows no cross-reaction with v-myc. c-myc is involved in the control of cell proliferation and differentiation and is amplified and/or overexpressed in a variety of turors. Overexpression of c-myc protein occurs frequently in luminal cells of prostate intraepithelial neoplasia as well as in most primary carcinomas and metastatic disease.ImmunogenA synthetic peptide, corresponding to aa 408-439 (AEEQKLISEEDLLRKRREQLKHKLEQLRNSCA) from C-terminus of human c-	Clonality	Monoclonal
IsotypeIgG1 KappaConjugateAlexa Fluor 350PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID4609Gene SymbolMYCSpeciesHuman, Chicken (Negative), Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse, Rat or Chicken.Specificity/SensitivityIt recognizes a transcription factor of 64-67kDa, identified as c-myc. Its epitope spans between aa 410-419 (EQKLISEEDL) which is a specific portion of an alpha helical region of human c-myc protein. This monoclonal antibody shows no cross-reaction with v-myc. c-myc is involved in the control of cell proliferation and differentiation and is amplified and/or overexpressed in a variety of tumors. Overexpression of c-myc protein occurs frequently in luminal cells of prostate intraepithelial neoplasia as well as in most primary carcinomas and metastatic disease.ImmunogenA synthetic peptide, corresponding to aa 408-439 (AEEQKLISEEDLLRKRREQLKHKLEQLRNSCA) from C-terminus of human c-	Clone	SPM237
ConjugateAlexa Fluor 350PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID4609Gene SymbolMYCSpeciesHuman, Chicken (Negative), Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse, Rat or Chicken.Specificity/SensitivityIt recognizes a transcription factor of 64-67kDa, identified as c-myc. Its epitope spans between aa 410-419 (EQKLISEEDL) which is a specific portion of an alpha helical region of human c-myc protein. This monoclonal antibody shows no cross-reaction with v-myc. c-myc is involved in the control of cell proliferation and differentiation and is amplified and/or overexpressed in a variety of tumors. Over-expression of c-myc protein occurs frequently in luminal cells of prostate intraepithelial neoplasia as well as in most primary carcinomas and metastatic disease.ImmunogenA synthetic peptide, corresponding to aa 408-439 (AEEQKLISEEDLLRKRREQLKHKLEQLRNSCA) from C-terminus of human c-	Preservative	0.05% Sodium Azide
Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Mouse Gene ID 4609 Gene Symbol MYC Species Human, Chicken (Negative), Mouse (Negative), Rat (Negative) Reactivity Notes Does not react with Mouse, Rat or Chicken. Specificity/Sensitivity It recognizes a transcription factor of 64-67kDa, identified as c-myc. Its epitope spans between aa 410-419 (EQKLISEEDL) which is a specific portion of an alpha helical region of human c-myc protein. This monoclonal antibody shows no cross-reaction with v-myc. c-myc is involved in the control of cell proliferation and differentiation and is amplified and/or overexpressed in a variety of tumors. Over-expression of c-myc protein occurs frequently in luminal cells of prostate intraepithelial neoplasia as well as in most primary carcinomas and metastatic disease. Immunogen A synthetic peptide, corresponding to aa 408-439 (AEEQKLISEEDLLRKRREQLKHKLEQLRNSCA) from C-terminus of human c-	Isotype	IgG1 Kappa
Buffer50mM Sodium BorateProduct DescriptionHostMouseGene ID4609Gene SymbolMYCSpeciesHuman, Chicken (Negative), Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse, Rat or Chicken.Specificity/SensitivityIt recognizes a transcription factor of 64-67kDa, identified as c-myc. Its epitope spans between aa 410-419 (EQKLISEEDL) which is a specific portion of an alpha helical region of human c-myc protein. This monoclonal antibody shows no cross-reaction with v-myc. c-myc is involved in the control of cell proliferation and differentiation and is amplified and/or overexpressed in a variety of tumors. Over-expression of c-myc protein occurs frequently in luminal cells of prostate intraepithelial neoplasia as well as in most primary carcinomas and metastatic disease.ImmunogenA synthetic peptide, corresponding to aa 408-439 (AEEQKLISEEDLLRKRREQLKHKLEQLRNSCA) from C-terminus of human c-	Conjugate	Alexa Fluor 350
Product Description Host Mouse Gene ID 4609 Gene Symbol MYC Species Human, Chicken (Negative), Mouse (Negative), Rat (Negative) Reactivity Notes Does not react with Mouse, Rat or Chicken. Specificity/Sensitivity It recognizes a transcription factor of 64-67kDa, identified as c-myc. Its epitope spans between aa 410-419 (EQKLISEEDL) which is a specific portion of an alpha helical region of human c-myc protein. This monoclonal antibody shows no cross-reaction with v-myc. c-myc is involved in the control of cell proliferation and differentiation and is amplified and/or overexpressed in a variety of tumors. Overexpression of c-myc protein occurs frequently in luminal cells of prostate intraepithelial neoplasia as well as in most primary carcinomas and metastatic disease. Immunogen A synthetic peptide, corresponding to aa 408-439 (AEEQKLISEEDLLRKRREQLKHKLEQLRNSCA) from C-terminus of human c-	Purity	Protein A or G purified
HostMouseGene ID4609Gene SymbolMYCSpeciesHuman, Chicken (Negative), Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse, Rat or Chicken.Specificity/SensitivityIt recognizes a transcription factor of 64-67kDa, identified as c-myc. Its epitope spans between aa 410-419 (EQKLISEEDL) which is a specific portion of an alpha helical region of human c-myc protein. This monoclonal antibody shows no cross-reaction with v-myc. c-myc is involved in the control of cell proliferation and differentiation and is amplified and/or overexpressed in a variety of tumors. Overexpression of c-myc protein occurs frequently in luminal cells of prostate intraepithelial neoplasia as well as in most primary carcinomas and metastatic disease.ImmunogenA synthetic peptide, corresponding to aa 408-439 (AEEQKLISEEDLLRKRREQLKHKLEQLRNSCA) from C-terminus of human c-	Buffer	50mM Sodium Borate
Gene ID4609Gene SymbolMYCSpeciesHuman, Chicken (Negative), Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse, Rat or Chicken.Specificity/SensitivityIt recognizes a transcription factor of 64-67kDa, identified as c-myc. Its epitope spans between aa 410-419 (EQKLISEEDL) which is a specific portion of an alpha helical region of human c-myc protein. This monoclonal antibody shows no cross-reaction with v-myc. c-myc is involved in the control of cell proliferation and differentiation and is amplified and/or overexpressed in a variety of tumors. Over- expression of c-myc protein occurs frequently in luminal cells of prostate intraepithelial neoplasia as well as in most primary carcinomas and metastatic disease.ImmunogenA synthetic peptide, corresponding to aa 408-439 (AEEQKLISEEDLLRKRREQLKHKLEQLRNSCA) from C-terminus of human c-	Product Description	
Gene SymbolMYCSpeciesHuman, Chicken (Negative), Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse, Rat or Chicken.Specificity/SensitivityIt recognizes a transcription factor of 64-67kDa, identified as c-myc. Its epitope spans between aa 410-419 (EQKLISEEDL) which is a specific portion of an alpha helical region of human c-myc protein. This monoclonal antibody shows no cross-reaction with v-myc. c-myc is involved in the control of cell proliferation and differentiation and is amplified and/or overexpressed in a variety of tumors. Over- expression of c-myc protein occurs frequently in luminal cells of prostate intraepithelial neoplasia as well as in most primary carcinomas and metastatic disease.ImmunogenA synthetic peptide, corresponding to aa 408-439 (AEEQKLISEEDLLRKRREQLKHKLEQLRNSCA) from C-terminus of human c-	Host	Mouse
SpeciesHuman, Chicken (Negative), Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse, Rat or Chicken.Specificity/SensitivityIt recognizes a transcription factor of 64-67kDa, identified as c-myc. Its epitope spans between aa 410-419 (EQKLISEEDL) which is a specific portion of an alpha helical region of human c-myc protein. This monoclonal antibody shows no cross-reaction with v-myc. c-myc is involved in the control of cell proliferation and differentiation and is amplified and/or overexpressed in a variety of tumors. Over- expression of c-myc protein occurs frequently in luminal cells of prostate intraepithelial neoplasia as well as in most primary carcinomas and metastatic disease.ImmunogenA synthetic peptide, corresponding to aa 408-439 (AEEQKLISEEDLLRKRREQLKHKLEQLRNSCA) from C-terminus of human c-	Gene ID	4609
Reactivity NotesDoes not react with Mouse, Rat or Chicken.Specificity/SensitivityIt recognizes a transcription factor of 64-67kDa, identified as c-myc. Its epitope spans between aa 410-419 (EQKLISEEDL) which is a specific portion of an alpha helical region of human c-myc protein. This monoclonal antibody shows no cross-reaction with v-myc. c-myc is involved in the control of cell proliferation and differentiation and is amplified and/or overexpressed in a variety of tumors. Over- expression of c-myc protein occurs frequently in luminal cells of prostate intraepithelial neoplasia as well as in most primary carcinomas and metastatic disease.ImmunogenA synthetic peptide, corresponding to aa 408-439 (AEEQKLISEEDLLRKRREQLKHKLEQLRNSCA) from C-terminus of human c-	Gene Symbol	MYC
Specificity/SensitivityIt recognizes a transcription factor of 64-67kDa, identified as c-myc. Its epitope spans between aa 410-419 (EQKLISEEDL) which is a specific portion of an alpha helical region of human c-myc protein. This monoclonal antibody shows no cross-reaction with v-myc. c-myc is involved in the control of cell proliferation and differentiation and is amplified and/or overexpressed in a variety of tumors. Over- expression of c-myc protein occurs frequently in luminal cells of prostate intraepithelial neoplasia as well as in most primary carcinomas and metastatic disease.ImmunogenA synthetic peptide, corresponding to aa 408-439 (AEEQKLISEEDLLRKRREQLKHKLEQLRNSCA) from C-terminus of human c-	Species	Human, Chicken (Negative), Mouse (Negative), Rat (Negative)
spans between aa 410-419 (EQKLISEEDL) which is a specific portion of an alpha helical region of human c-myc protein. This monoclonal antibody shows no cross-reaction with v-myc. c-myc is involved in the control of cell proliferation and differentiation and is amplified and/or overexpressed in a variety of tumors. Over- expression of c-myc protein occurs frequently in luminal cells of prostate intraepithelial neoplasia as well as in most primary carcinomas and metastatic disease.ImmunogenA synthetic peptide, corresponding to aa 408-439 (AEEQKLISEEDLLRKRREQLKHKLEQLRNSCA) from C-terminus of human c-	Reactivity Notes	Does not react with Mouse, Rat or Chicken.
(AEEQKLISEEDLLRKRREQLKHKLEQLRNSCA) from C-terminus of human c-	Specificity/Sensitivity	spans between aa 410-419 (EQKLISEEDL) which is a specific portion of an alpha helical region of human c-myc protein. This monoclonal antibody shows no cross-reaction with v-myc. c-myc is involved in the control of cell proliferation and differentiation and is amplified and/or overexpressed in a variety of tumors. Over- expression of c-myc protein occurs frequently in luminal cells of prostate intraepithelial neoplasia as well as in most primary carcinomas and metastatic
	Immunogen	(AEEQKLISEEDLLRKRREQLKHKLEQLRNSCA) from C-terminus of human c-



Notes

Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.

Product Application Details

••	
Applications	Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Immunofluorescence
Recommended Dilutions	Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence, CyTOF-ready
Application Notes	Optimal dilution of this antibody should be experimentally determined.

Images

c-Myc Antibody (SPM237) [Alexa Fluor® 350] [NBP2-86682AF350] - Vial of Alexa Fluor 350 conjugated antibody. Alexa Fluor 350 is optimally excited at 346 nm by the UV laser (350 or 355 nm) and has an emission maximum of 442 nm.

III CEREAL IIII	Alexa Fluo	or® 350	
	LASER (nm)	FILTER	7
Alexa Fluor ^e 3 ⁵⁰	LASER (nm) UV (350)	FILTER 450/45]
Alexa Fluore 350	the second se]







Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-86682AF350

H00004609-P01-10ug	Recombinant Human c-Myc GST (N-Term) Protein
210-TA-005	TNF-alpha [Unconjugated]
NBL1-13414	c-Myc Overexpression Lysate
236-EG-200	EGF [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-86682AF350

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

