

Product Datasheet

CD11b Antibody - BSA Free

NBP2-92978-0.1ml

Unit Size: 0.1 ml

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-92978

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-92978



NBP2-92978-0.1ml

CD11b Antibody - BSA Free

Product Information

Unit Size	0.1 ml
Concentration	Please see the vial label for concentration. If unlisted please contact technical services.
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.09% Sodium Azide
Isotype	IgG
Purity	Affinity purified
Buffer	PBS (pH 7.3), 50% glycerol
Target Molecular Weight	127 kDa

Product Description


Description	Novus Biologicals Rabbit CD11b Antibody - BSA Free (NBP2-92978) is a polyclonal antibody validated for use in IHC, WB and ICC/IF. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	3684
Gene Symbol	ITGAM
Species	Human, Mouse, Rat
Immunogen	Recombinant fusion protein containing a sequence corresponding to amino acids 145-340 of CD11b (NP_000623.2). PQEDSDIAFLIDGSGSIIIPHDFRRMKEFVSTVMEQLKKSKTLFSLMQYSEEFRIH FTFKEFQNNPNPRSLVKPITQLLGRTHATGIRKVVRELFNITNGARKNAFKILVV ITDGEKFGDPLGYEDVIPEADREGVIRYVIGVGDAFRSEKSRQELNTIASKPPRD HVFQVNNFEALKTIQNQLREKIFAIEGTQT

Product Application Details


Applications	Western Blot, Immunohistochemistry-Paraffin, Immunocytochemistry/Immunofluorescence, Immunohistochemistry
Recommended Dilutions	Western Blot 1:100 - 1:500, Immunohistochemistry 1:50-1:200, Immunocytochemistry/ Immunofluorescence 1:50 -1:200, Immunohistochemistry-Paraffin

Images


Western Blot: CD11b Antibody [NBP2-92978] - Analysis of extracts of various cell lines, using CD11b antibody (NBP2-92978) at 1:500 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 90s.




Immunohistochemistry-Paraffin: CD11b Antibody [NBP2-92978] - Mouse spleen using CD11b antibody (NBP2-92978) at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.




Immunohistochemistry-Paraffin: CD11b Antibody [NBP2-92978] - Human tonsil using CD11b antibody (NBP2-92978) at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.




Immunocytochemistry/Immunofluorescence: CD11b Antibody [NBP2-92978] - Immunofluorescence analysis of TF-1 cells using CD11b Rabbit pAb (NBP2-92978) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.




Immunocytochemistry/ Immunofluorescence: CD11b Antibody - Azide and BSA Free [NBP2-92978] - Immunofluorescence analysis of TF-1 cells using CD11b Rabbit pAb at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.




Immunocytochemistry/ Immunofluorescence: CD11b Antibody - Azide and BSA Free [NBP2-92978] - Immunofluorescence analysis of THP-1 cells using CD11b Rabbit pAb at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.




Immunocytochemistry/ Immunofluorescence: CD11b Antibody - Azide and BSA Free [NBP2-92978] - Immunofluorescence analysis of THP-1 cells using CD11b Rabbit pAb at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunocytochemistry/ Immunofluorescence: CD11b Antibody - Azide and BSA Free [NBP2-92978] - Immunofluorescence analysis of THP-1 cells using CD11b Rabbit pAb at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunocytochemistry/ Immunofluorescence: CD11b Antibody - Azide and BSA Free [NBP2-92978] - Immunofluorescence analysis of TF-1 cells using CD11b Rabbit pAb at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-92978-0.1ml

HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control
NB110-89474PEP	CD11b Antibody Blocking Peptide

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-92978

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

