Product Datasheet

SIX2 Antibody - Azide and BSA Free NBP2-94801-0.1ml

Unit Size: 0.1 ml

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-94801

Updated 11/7/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-94801



NBP2-94801-0.1ml


| SIX2 Antibody - Azide and BSA Free | |
|------------------------------------|---|
| Product Information | |
| Unit Size | 0.1 ml |
| Concentration | Please see the vial label for concentration. If unlisted please contact technical services. |
| Storage | Store at -20C. Avoid freeze-thaw cycles. |
| Clonality | Polyclonal |
| Preservative | 0.01% Thimerosal |
| Isotype | IgG |
| Purity | Affinity purified |
| Buffer | PBS (pH 7.3), 50% glycerol |
| Target Molecular Weight | 32 kDa |
| Product Description | |
| Host | Rabbit |
| Gene ID | 10736 |
| Gene Symbol | SIX2 |
| Species | Human, Rat |
| Immunogen | Recombinant fusion protein containing a sequence corresponding to amino acids 120-174 of human SIX2 (NP_058628.3). SIWDGEETSYCFKEKSRSVLREWYAHNPYPSPREKRELAEATGLTTTQVSNW FKN |
| Product Application Details | |
| Applications | Western Blot |

| Product Application Details | |
|------------------------------------|-----------------------------|
| Applications | Western Blot |
| Recommended Dilutions | Western Blot 1:500 - 1:1000 |




Images

Western Blot: SIX2 Antibody [NBP2-94801] - Western blot analysis of extracts of various cell lines, using SIX2 antibody (NBP2-94801) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 10s.



Western Blot: SIX2 Antibody [NBP2-94801] - Western blot analysis of extracts of Rat skeletal muscle cells, using SIX2 antibody (NBP2-94801) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 180s.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Toronto, ON M8Z 4E6

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada

Bio-Techne Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Error: Subreport could not be shown.

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-94801

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

