

Product Datasheet

Daudi Whole Cell Lysate NBP3-05285

Unit Size: 500 ug

Store at -70C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP3-05285

Updated 7/23/2023 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP3-05285



NBP3-05285**Daudi Whole Cell Lysate****Product Information**

Unit Size	500 ug
Concentration	Please contact technical services for concentration.
Storage	Store at -70C. Avoid freeze-thaw cycles.
Preservative	No Preservative
Purity	Multi-step
Buffer	1X SDS-PAGE Sample Buffer (62.5 mM Tris HCl, 2% SDS, 10% (v/v) Glycerol and 0.005% bromophenol blue, pH 6.8)

Product Description

Description	Store vial at -70C or COLDER. For extended storage, aliquot contents to minimize freeze/thaw cycles.
Species	Human
Preparation Method	The cells were grown in Dulbecco's medium supplemented with 10% fetal bovine serum. Cells were washed with PBS and then incubated on ice in modified RIPA buffer, containing 150 mM sodium chloride, 50 mM Tris HCl, pH 7.4, 1 mM EDTA, 1.0% NP-40, 0.5% sodium deoxycholic acid, 0.1% SDS and 0.01% (w/v) sodium azide to lyse the cells. Protein integrity was ensured using a cocktail of protease inhibitors with broad specificity for the inhibition of aspartic, cysteine, and serine proteases as well as aminopeptidases (0.1 mM AEBSF HCl, 0.08 uM Aprotinin, 5 uM Bestatin, 1.5 uM E-64, 2 uM Leupeptin Hemisulfate, 1 uM Pepstatin A). Phosphatase inhibitors 1 mM NaF and 1 mM Na3VO4 were also added. Cell debris was removed by centrifugation. Protein concentration was determined by a modified BCA assay using a commercially available kit. Protein concentration was adjusted to 2 mg/mL and then an equal volume of 2X SDS-PAGE sample buffer was added.
Lysate Type	Cell
Lysate Subcellular Fraction	Whole


Product Application Details

Applications	Western Blot, SDS-Page
Recommended Dilutions	Western Blot, SDS-Page
Application Notes	This product has been tested by SDS-PAGE and western blot. Ready-to-use lysates are especially prepared as positive controls for separation by SDS-PAGE and subsequent western blot analysis. Lysates are prepared in denaturing buffer WITHOUT dissociating agents (i.e. no 2-mercaptoethanol or dithiothreitol has been added). Heat lysate to 95C for 5 minutes and rapidly cool. If dissociating conditions are desired, add reducing agent prior to heating. The recommended loading volume per lane is 10-20 uL depending on the size format of your gel.



Images

SDS-Page: Daudi Whole Cell Lysate [NBP3-05285] - Lane 1: Daudi Whole Cell Lysate (10ug) Reduced. Lane 2: Opal Prestained Molecular Weight Markers. Lane 3: Daudi Whole Cell Lysate (10ug) Non-Reduced. Results show wide range of molecular weight bands with no signs of degradation.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Lysates are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-05285

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

