Product Datasheet CSTF2T Antibody (PCRP-CSTF2T-1A3) NBP3-07486-100ug

Unit Size: 100 ug

Store at 4C.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-07486

Updated 7/16/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-07486



NBP3-07486-100ug


CSTF2T Antibody (PCRP-CSTF2T-1A3)

	•••···	- /
Concentration 0.2 mg/ml Storage Store at 4C. Clonality Monoclonal Clone PCRP-CSTF2T-1A3 Preservative 0.05% Sodium Azide Isotype IgG2b Kappa Purity Protein A or G purified Buffer 10 mM PBS with 0.05% BSA Target Molecular Weight 64 kDa Product Description 200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP3-08807) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -8t C. Host Mouse Gene ID 23283 Gene BD CSTF2T Species Human Species Human Species Human Species Human Species Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Immunofluorescence, Immunohistochemistry, Immunohistochemistry/ Immunofluorescence, Immunohistochemistry, Immunohytochemistry/Immunofluorescence, Immunohistochemistry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunofluorescence 1-2 ug/ml, Inmunohistochemistry, Immunofluorescence 1-2 ug/ml, Immunohistochemistry, Immunofluorescence 1-2 ug/ml, Immunohistochemis	Product Information	
Storage Store at 4C. Clonality Monoclonal Clone PCRP-CSTF2T-1A3 Preservative 0.05% Sodium Azide Isotype IgG2b Kappa Purity Protein A or G purified Buffer 10 mM PBS with 0.05% BSA Target Molecular Weight 64 kDa Product Description 200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP3-08807) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80 C. Gene ID 23283 Gene Symbol CSTF2T Species Human Specificity/Sensitivity This gene encodes a nuclear protein with an RRM (RNA recognition motif) domain. The protein is a member of the cleavage stimulation factor (CSTF) complex that is involved in the 3-untranslated region of mRNAs. Immunogen Recombinant full-length human CSTF2T protein (Uniprot: Q9H0L4) Product Application Details Western Blot, Flow Cytometry, Immunofucrescence, Immunohistochemistry, Immunofucrescence, Immunohistochemistry, Immunofucrescence 1-2 ug/ml, Immunofucrescence 1-2 ug/ml, Immunohistochemistry, Immunofucrescence 1-2 ug/ml, Immunohistochemistry, Immunofucrescence 1-2 ug/ml, Protein Array Application Notes Immunohistochemistry (Formalin-fixed	Unit Size	100 ug
Clonality Monoclonal Clonality Monoclonal Clone PCRP-CSTF2T-1A3 Preservative 0.05% Sodium Azide Isotype IgG2b Kappa Purity Protein A or G purified Buffer 10 mM PBS with 0.05% BSA Target Molecular Weight 64 kDa Product Description 200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (MBP3-08807) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80 C. Gene ID 23283 Gene BD 23283 Gene Symbol CSTF2T Species Human Species Human Specificity/Sensitivity This gene encodes a nuclear protein with an RRM (RNA recognition motif) domain. The protein is a member of the cleavage stimulation factor (CSTF) complex that is involved in the 3-untranstated region of mRNAs. Immunogen Recombinant full-length human CSTF2T protein (Uniprot: Q9H0L4) Product Application Details Western Blot, Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunofluor cells, Immunohistochemistry, Immunofluorescence 1-2 ug/ml, Immunohistochemistry, Inmunofluorescen	Concentration	0.2 mg/ml
Clone PCRP-CSTF2T-1A3 Preservative 0.05% Sodium Azide Isotype IgG2b Kappa Purity Protein A or G purified Buffer 10 mM PBS with 0.05% BSA Target Molecular Weight 64 kDa Product Description 200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP3-08807) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80 C. Host Mouse Gene ID 23283 Gene Symbol CSTF2T Specificity/Sensitivity This gene encodes a nuclear protein with an RRM (RNA recognition motif) domain. The protein is a member of the cleavage stimulation factor (CSTF) complex that is involved in the 3-untranslated region of mRNAs. Immunogen Recombinant full-length human CSTF2T protein (Uniprot: Q9H0L4) Product Application Details Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry/ Paraffin, Protein Array Recommended Dilutions Western Blot 1-2 ug/ml, Flow Cytometry 1-2 ug/million cells, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry, Paraffin 1-2 ug/ml for 30 minutes at RT. Staining of formali	Storage	Store at 4C.
Preservative 0.05% Sodium Azide Isotype IgG2b Kappa Purity Protein A or G purified Buffer 10 mM PBS with 0.05% BSA Target Molecular Weight 64 kDa Product Description 200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP3-08807) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80 C. Host Mouse Gene ID 23283 Gene Symbol CSTF2T Species Human Species Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Im	Clonality	Monoclonal
Isotype IgG2b Kappa Purity Protein A or G purified Buffer 10 mM PBS with 0.05% BSA Target Molecular Weight 64 kDa Product Description 200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP3-08807) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80 C. Host Mouse Gene ID 23283 Gene Symbol CSTF2T Species Human Species Human Specificity/Sensitivity This gene encodes a nuclear protein with an RRM (RNA recognition motif) domain. The protein is a member of the cleavage stimulation factor (CSTF) complex that is involved in the 3-untranslated region of mRNAs. Immunogen Recombinant full-length human CSTF2T protein (Uniprot: Q9H0L4) Product Application Details Western Blot, Flow Cytometry, Immunofucorescence, Immunohistochemistry, Immunohistochemistry-Parafin, Protein Array Recommended Dilutions Western Blot, Flow Cytometry, 1-2 ug/million cells, Immunohistochemistry, Immunohistochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry, Immunofluorescence 1-2 ug/ml, Immunohistochemistry, Immunohistochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry -Parafin 1-2 ug/ml for 30 minutes at RT. Staining of formalin-fixed issues requires heating tissue	Clone	PCRP-CSTF2T-1A3
Purity Protein A or G purified Buffer 10 mM PBS with 0.05% BSA Target Molecular Weight 64 kDa Product Description 200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP3-08807) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80 C. Host Mouse Gene ID 23283 Gene Symbol CSTF2T Species Human Specificity/Sensitivity This gene encodes a nuclear protein with an RRM (RNA recognition motif) domain. The protein is a member of the cleavage stimulation factor (CSTF) complex that is involved in the 3-untranslated region of mRNAs. Immunogen Recombinant full-length human CSTF2T protein (Uniprot: Q9H0L4) Product Application Details Western Blot, Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Protein Array Recommended Dilutions Western Blot 1-2 ug/ml, Flow Cytometry 1-2 ug/million cells, Immunohistochemistry, Immunocytochemistry/Immunofluorescence 1-2 ug/ml, Immunohistochemistry, Immunocytochemistry/Immunefluorescence 1-2 ug/ml, Immunohistochemistry, Immunocytochemistry 1-2 ug/millon cells, Immunohistochemistry, Immunocytochemistry 1-2 ug/millon cells, Immunohistochemistry, Immunocytochemistry Immunefluorescence 1-2 ug/ml, Immunohistochemistry, Immunocytochemist	Preservative	0.05% Sodium Azide
Buffer 10 mM PBS with 0.05% BSA Target Molecular Weight 64 kDa Product Description 200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP3-08807) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -8 C. Host Mouse Gene ID 23283 Gene Symbol CSTF2T Species Human Specificity/Sensitivity This gene encodes a nuclear protein with an RRM (RNA recognition motif) domain. The protein is a member of the cleavage stimulation factor (CSTF) complex that is involved in the 3-untranslated region of mRNAs. Immunogen Recombinant full-length human CSTF2T protein (Uniprot: Q9H0L4) Product Application Details Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Immunofluorescence 1-2 ug/ml, Forein Array Application Notes Immunohistochemistry, Gormalin-fixed) 1-2 ug/ml, Protein Array Application Notes Immunohistochemistry Parafiin 1-2 ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 11mM	Isotype	IgG2b Kappa
Target Molecular Weight 64 kDa Product Description 200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP3-08807) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80 C. Host Mouse Gene ID 23283 Gene Symbol CSTF2T Species Human Specificity/Sensitivity This gene encodes a nuclear protein with an RRM (RNA recognition motif) domain. The protein is a member of the cleavage stimulation factor (CSTF) complex that is involved in the 3-untranslated region of mRNAs. Immunogen Recombinant full-length human CSTF2T protein (Uniprot: Q9H0L4) Product Application Details Applications Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Protein Array Recommended Dilutions Western Blot 1-2 ug/ml, Flow Cytometry 1-2 ug/million cells, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry. Immunofluorescence 1-2 ug/ml, Immunohistochemistry. Foradin fixed: 1-2 ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Purity	Protein A or G purified
Product Description 200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP3-08807) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80 C. Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80 C. Host Mouse Gene ID 23283 Gene Symbol CSTF2T Species Specificity/Sensitivity This gene encodes a nuclear protein with an RRM (RNA recognition motif) domain. The protein is a member of the cleavage stimulation factor (CSTF) complex that is involved in the 3-untranslated region of mRNAs. Immunogen Recombinant full-length human CSTF2T protein (Uniprot: Q9H0L4) Product Application Details Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry/Immunofluorescence 1-2 ug/ml, Immunohistochemistry, Immunocytochemistry/Immunofluorescence 1-2 ug/ml, Immunohistochemistry (Formalin-fixed): 1-2 ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Buffer	10 mM PBS with 0.05% BSA
Description 200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP3-08807) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -8 C. Host Mouse Gene ID 23283 Gene Symbol CSTF2T Species Human Specificity/Sensitivity This gene encodes a nuclear protein with an RRM (RNA recognition motif) domain. The protein is a member of the cleavage stimulation factor (CSTF) complex that is involved in the 3-untranslated region of mRNAs. Immunogen Recombinant full-length human CSTF2T protein (Uniprot: Q9H0L4) Product Application Details Applications Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Paraffin, Protein Array Recommended Dilutions Western Blot 1-2 ug/ml, Flow Cytometry 1-2 ug/million cells, Immunohistochemistry, Immunofluorescence 1-2 ug/ml, Immunohistochemistry, Immunofluorescence 1-2 ug/ml, Immunohistochemistry, Paraffin 1-2 ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Target Molecular Weight	64 kDa
Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP3-08807) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80 C. Host Gene ID 23283 Gene Symbol CSTF2T Species Human Specificity/Sensitivity This gene encodes a nuclear protein with an RRM (RNA recognition motif) domain. The protein is a member of the cleavage stimulation factor (CSTF) complex that is involved in the 3-untranslated region of mRNAs. Immunogen Recombinant full-length human CSTF2T protein (Uniprot: Q9H0L4) Product Application Details Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry/ Paraffin, Protein Array Recommended Dilutions Western Blot 1-2 ug/ml, Flow Cytometry 1-2 ug/million cells, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry, Immunofluorescence 1-2 ug/ml, Immunohistochemistry, Immunofluorescence 1-2 ug/ml, Immunohistochemistry, Immunofluorescence 1-2 ug/ml, Flow Cytometry 1-2 ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Product Description	
C. Mouse Gene ID 23283 Gene Symbol CSTF2T Species Human Specificity/Sensitivity This gene encodes a nuclear protein with an RRM (RNA recognition motif) domain. The protein is a member of the cleavage stimulation factor (CSTF) complex that is involved in the 3-untranslated region of mRNAs. Immunogen Recombinant full-length human CSTF2T protein (Uniprot: Q9H0L4) Product Application Details Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunocytochemistry-Paraffin, Protein Array Recommended Dilutions Western Blot 1-2 ug/ml, Flow Cytometry 1-2 ug/million cells, Immunohistochemistry-Paraffin 1-2 ug/ml, Protein Array Application Notes Immunohistochemistry (Formalin-fixed) 1-2 ug/ml, Protein Array Application Notes Immunohistochemistry (Formalin-fixed) 1-2 ug/ml, For Array	Description	Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP3-08807)
Gene ID 23283 Gene Symbol CSTF2T Species Human Specificity/Sensitivity This gene encodes a nuclear protein with an RRM (RNA recognition motif) domain. The protein is a member of the cleavage stimulation factor (CSTF) complex that is involved in the 3-untranslated region of mRNAs. Immunogen Recombinant full-length human CSTF2T protein (Uniprot: Q9H0L4) Product Application Details Mestern Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Protein Array Recommended Dilutions Western Blot 1-2 ug/ml, Flow Cytometry 1-2 ug/million cells, Immunohistochemistry. Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry.Paraffin 1-2 ug/ml, Protein Array Application Notes Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.		
Gene Symbol CSTF2T Species Human Specificity/Sensitivity This gene encodes a nuclear protein with an RRM (RNA recognition motif) domain. The protein is a member of the cleavage stimulation factor (CSTF) complex that is involved in the 3-untranslated region of mRNAs. Immunogen Recombinant full-length human CSTF2T protein (Uniprot: Q9H0L4) Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Protein Array Recommended Dilutions Western Blot 1-2 ug/ml, Flow Cytometry 1-2 ug/million cells, Immunohistochemistry, Immunofluorescence 1-2 ug/ml, Immunohistochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry/ Paraffin 1-2 ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Host	Mouse
SpeciesHumanSpecificity/SensitivityThis gene encodes a nuclear protein with an RRM (RNA recognition motif) domain. The protein is a member of the cleavage stimulation factor (CSTF) complex that is involved in the 3-untranslated region of mRNAs.ImmunogenRecombinant full-length human CSTF2T protein (Uniprot: Q9H0L4)Product Application DetailsWestern Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Protein ArrayRecommended DilutionsWestern Blot 1-2 ug/ml, Flow Cytometry 1-2 ug/million cells, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry-Paraffin 1-2 ug/ml, Protein ArrayApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Gene ID	23283
Specificity/SensitivityThis gene encodes a nuclear protein with an RRM (RNA recognition motif) domain. The protein is a member of the cleavage stimulation factor (CSTF) complex that is involved in the 3-untranslated region of mRNAs.ImmunogenRecombinant full-length human CSTF2T protein (Uniprot: Q9H0L4)Product Application DetailsWestern Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Protein ArrayRecommended DilutionsWestern Blot 1-2 ug/ml, Flow Cytometry 1-2 ug/million cells, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry Paraffin 1-2 ug/ml, Protein ArrayApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Gene Symbol	CSTF2T
domain. The protein is a member of the cleavage stimulation factor (CSTF) complex that is involved in the 3-untranslated region of mRNAs.ImmunogenRecombinant full-length human CSTF2T protein (Uniprot: Q9H0L4)Product Application DetailsWestern Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Protein ArrayRecommended DilutionsWestern Blot 1-2 ug/ml, Flow Cytometry 1-2 ug/million cells, Immunohistochemistry, Immunohistochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry Paraffin 1-2 ug/ml, Protein ArrayApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Species	Human
Product Application DetailsApplicationsWestern Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Protein ArrayRecommended DilutionsWestern Blot 1-2 ug/ml, Flow Cytometry 1-2 ug/million cells, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry-Paraffin 1-2 ug/ml, Protein ArrayApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Specificity/Sensitivity	domain. The protein is a member of the cleavage stimulation factor (CSTF)
ApplicationsWestern Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Protein ArrayRecommended DilutionsWestern Blot 1-2 ug/ml, Flow Cytometry 1-2 ug/million cells, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry-Paraffin 1-2 ug/ml, Protein ArrayApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Immunogen	Recombinant full-length human CSTF2T protein (Uniprot: Q9H0L4)
Immunohistochemistry, Immunohistochemistry-Paraffin, Protein ArrayRecommended DilutionsWestern Blot 1-2 ug/ml, Flow Cytometry 1-2 ug/million cells, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry-Paraffin 1-2 ug/ml, Protein ArrayApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Product Application Details	
Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml, Immunohistochemistry-Paraffin 1-2 ug/ml, Protein ArrayApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Applications	
of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Recommended Dilutions	Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1-2 ug/ml,
	Application Notes	

www.novusbio.com











Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP3-07486-100ug

HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
NBP1-43317-0.5mg	Mouse IgG2b Kappa Light Chain Isotype Control (MG2b)
H00023283-P01-10ug	Recombinant Human CSTF2T GST (N-Term) Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-07486

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

