Product Datasheet

Cytokeratin 7/17 Antibody (C-46) [Alexa Fluor® 488] NBP3-08343AF488

Unit Size: 100 ul

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-08343AF488

Updated 10/26/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP3-08343AF488



NBP3-08343AF488

Cytokeratin 7/17 Antibody (C-46) [Alexa Fluor® 488]

Cytokeratin 7/17 Antibody (C-46) [Alexa Fluor® 488]		
Product Information		
Unit Size	100 ul	
Concentration	Please see the vial label for concentration. If unlisted please contact technical services.	
Storage	Store at 4C in the dark.	
Clonality	Monoclonal	
Clone	C-46	
Preservative	0.05% Sodium Azide	
Isotype	IgG1 Kappa	
Conjugate	Alexa Fluor 488	
Purity	Protein A or G purified	
Buffer	50mM Sodium Borate	
Product Description		
Host	Mouse	
Gene ID	3855	
Gene Symbol	KRT7	
Species	Human, Bovine, Ferret (Negative), Mouse (Negative), Rabbit (Negative), Rat (Negative)	
Reactivity Notes	Does not react with Mouse, Rat, Ferret or Rabbit.	
Specificity/Sensitivity	It recognizes intermediate filament proteins (IFPs) of 55kDa and 46kDa, which are identified as cytokeratin 7 and Cytokeratin 17 respectively. On formalin-fixed, paraffin embedded sections, C-46 reacts with keratin 7 only. Cytokeratins comprise a diverse group of intermediate filament proteins (IFPs) that are expressed as pairs in both keratinized and non-keratinized epithelial tissue. Cytokeratins play a critical role in differentiation and tissue specialization and function to maintain the overall structural integrity of epithelial cells. Cytokeratins have been found to be useful markers of tissue differentiation, which is directly applicable to the characterization of malignant tumors. Cytokeratin 7 is a basic cytokeratin, which is found in most glandular and transitional epithelia but not in the stratified squamous epithelia. Cytokeratin 7 is expressed in the epithelial cells of ovary, lung, and breast but not of colon, prostate, or gastrointestinal tract. Cytokeratin 17 is expressed in epithelial cells of various origins, such as bronchial epithelial cells and skin appendages. Cytokeratin 17 antibody marks basal cell differentiation.	
Immunogen	Cytoskeletal preparation from HeLa cells (human cervical carcinoma cells)	



	Page 2 of 3 v.20.1 Updated 10/26/2023
Notes	Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.
Product Application Details	

Product Application Details	
Applications	Western Blot, Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin
Recommended Dilutions	Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry-Paraffin
Application Notes	Optimal dilution of this antibody should be experimentally determined.



Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP3-08343AF488

IC002G Mouse IgG1 Isotype Control (11711) [Alexa Fluor® 488]

8184-CK-050 Choline Kinase beta [Unconjugated]

NB300-223 Vimentin Antibody

NB200-103 p53 Antibody (PAb 240) - BSA Free

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-08343AF488

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

