Product Datasheet

Cytokeratin 7/17 Antibody (C-46) [mFluor Violet 610 SE] NBP3-08343MFV610

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-08343MFV610

Updated 10/26/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP3-08343MFV610



NBP3-08343MFV610

Cytokeratin 7/17 Antibody (C-46) [mFluor Violet 610 SE]

	Cytokeratin 7/17 Antibody (C-46) [mindor violet 610 SE]	
Product Information		
Unit Size	0.1 ml	
Concentration	Please see the vial label for concentration. If unlisted please contact technical services.	
Storage	Store at 4C in the dark.	
Clonality	Monoclonal	
Clone	C-46	
Preservative	0.05% Sodium Azide	
Isotype	IgG1 Kappa	
Conjugate	mFluor Violet 610 SE	
Purity	Protein A or G purified	
Buffer	50mM Sodium Borate	
Product Description		
Host	Mouse	
Gene ID	3855	
Gene Symbol	KRT7	
Species	Human, Bovine, Ferret (Negative), Mouse (Negative), Rabbit (Negative), Rat (Negative)	
Reactivity Notes	Does not react with Mouse, Rat, Ferret or Rabbit.	
Specificity/Sensitivity	It recognizes intermediate filament proteins (IFPs) of 55kDa and 46kDa, which are identified as cytokeratin 7 and Cytokeratin 17 respectively. On formalin-fixed,	
	paraffin embedded sections, C-46 reacts with keratin 7 only. Cytokeratins comprise a diverse group of intermediate filament proteins (IFPs) that are expressed as pairs in both keratinized and non-keratinized epithelial tissue. Cytokeratins play a critical role in differentiation and tissue specialization and function to maintain the overall structural integrity of epithelial cells. Cytokeratins have been found to be useful markers of tissue differentiation, which is directly applicable to the characterization of malignant tumors. Cytokeratin 7 is a basic cytokeratin, which is found in most glandular and transitional epithelia but not in the stratified squamous epithelia. Cytokeratin 7 is expressed in the epithelial cells of ovary, lung, and breast but not of colon, prostate, or gastrointestinal tract. Cytokeratin 17 is expressed in epithelial cells of various origins, such as bronchial epithelial cells and skin appendages. Cytokeratin 17 antibody marks basal cell differentiation.	
Immunogen	comprise a diverse group of intermediate filament proteins (IFPs) that are expressed as pairs in both keratinized and non-keratinized epithelial tissue. Cytokeratins play a critical role in differentiation and tissue specialization and function to maintain the overall structural integrity of epithelial cells. Cytokeratins have been found to be useful markers of tissue differentiation, which is directly applicable to the characterization of malignant tumors. Cytokeratin 7 is a basic cytokeratin, which is found in most glandular and transitional epithelia but not in the stratified squamous epithelia. Cytokeratin 7 is expressed in the epithelial cells of ovary, lung, and breast but not of colon, prostate, or gastrointestinal tract. Cytokeratin 17 is expressed in epithelial cells of various origins, such as bronchial epithelial cells and skin appendages. Cytokeratin 17 antibody marks	
Immunogen Notes	comprise a diverse group of intermediate filament proteins (IFPs) that are expressed as pairs in both keratinized and non-keratinized epithelial tissue. Cytokeratins play a critical role in differentiation and tissue specialization and function to maintain the overall structural integrity of epithelial cells. Cytokeratins have been found to be useful markers of tissue differentiation, which is directly applicable to the characterization of malignant tumors. Cytokeratin 7 is a basic cytokeratin, which is found in most glandular and transitional epithelia but not in the stratified squamous epithelia. Cytokeratin 7 is expressed in the epithelial cells of ovary, lung, and breast but not of colon, prostate, or gastrointestinal tract. Cytokeratin 17 is expressed in epithelial cells of various origins, such as bronchial epithelial cells and skin appendages. Cytokeratin 17 antibody marks basal cell differentiation.	
	comprise a diverse group of intermediate filament proteins (IFPs) that are expressed as pairs in both keratinized and non-keratinized epithelial tissue. Cytokeratins play a critical role in differentiation and tissue specialization and function to maintain the overall structural integrity of epithelial cells. Cytokeratins have been found to be useful markers of tissue differentiation, which is directly applicable to the characterization of malignant tumors. Cytokeratin 7 is a basic cytokeratin, which is found in most glandular and transitional epithelia but not in the stratified squamous epithelia. Cytokeratin 7 is expressed in the epithelial cells of ovary, lung, and breast but not of colon, prostate, or gastrointestinal tract. Cytokeratin 17 is expressed in epithelial cells of various origins, such as bronchial epithelial cells and skin appendages. Cytokeratin 17 antibody marks basal cell differentiation. Cytoskeletal preparation from HeLa cells (human cervical carcinoma cells) mFluor(TM) is a trademark of AAT Bioquest, Inc. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The	
Notes	comprise a diverse group of intermediate filament proteins (IFPs) that are expressed as pairs in both keratinized and non-keratinized epithelial tissue. Cytokeratins play a critical role in differentiation and tissue specialization and function to maintain the overall structural integrity of epithelial cells. Cytokeratins have been found to be useful markers of tissue differentiation, which is directly applicable to the characterization of malignant tumors. Cytokeratin 7 is a basic cytokeratin, which is found in most glandular and transitional epithelia but not in the stratified squamous epithelia. Cytokeratin 7 is expressed in the epithelial cells of ovary, lung, and breast but not of colon, prostate, or gastrointestinal tract. Cytokeratin 17 is expressed in epithelial cells of various origins, such as bronchial epithelial cells and skin appendages. Cytokeratin 17 antibody marks basal cell differentiation. Cytoskeletal preparation from HeLa cells (human cervical carcinoma cells) mFluor(TM) is a trademark of AAT Bioquest, Inc. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The	
Notes Product Application Details	comprise a diverse group of intermediate filament proteins (IFPs) that are expressed as pairs in both keratinized and non-keratinized epithelial tissue. Cytokeratins play a critical role in differentiation and tissue specialization and function to maintain the overall structural integrity of epithelial cells. Cytokeratins have been found to be useful markers of tissue differentiation, which is directly applicable to the characterization of malignant tumors. Cytokeratin 7 is a basic cytokeratin, which is found in most glandular and transitional epithelia but not in the stratified squamous epithelia. Cytokeratin 7 is expressed in the epithelial cells of ovary, lung, and breast but not of colon, prostate, or gastrointestinal tract. Cytokeratin 17 is expressed in epithelial cells of various origins, such as bronchial epithelial cells and skin appendages. Cytokeratin 17 antibody marks basal cell differentiation. Cytoskeletal preparation from HeLa cells (human cervical carcinoma cells) mFluor(TM) is a trademark of AAT Bioquest, Inc. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Western Blot, Flow Cytometry, Immunocytochemistry/Immunofluorescence,	



Images

Cytokeratin 7/17 Antibody (C-46) [mFluor Violet 610 SE] [NBP3-08343MFV610] - Vial of mFluor Violet 610 conjugated antibody. mFluor Violet 610 is optimally excited at 421 nm by the Violet laser (405 nm) and has an emission maximum of 613 nm.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP3-08343MFV610

8184-CK-050	Choline Kinase beta [Unconjugated]
NB300-223	Vimentin Antibody
NB200-103	p53 Antibody (PAb 240) - BSA Free
NB100-687	Cytokeratin 19 Antibody - BSA Free

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-08343MFV610

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

