Product Datasheet

MCM7 Antibody (rMCM7/1468) [Alexa Fluor® 700] NBP3-08389AF700

Unit Size: 100 ul

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-08389AF700

Updated 10/26/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-08389AF700



NBP3-08389AF700

MCM7 Antibody (rMCM7/1468) [Alexa Fluor® 700]

Product InformationUnit Size100 ulConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalClonerMCM7/1468Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 700PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMonoclonalGene ID4176Gene ID4176Gene SymbolMCM7SpeciesHuman, CanineMarkerProliferation MarkerSpecificity/SensitivityThe specificity of this monoclonal antibody to its intended target was validated by HuProTIM Array, containing more than 19,000, full-length human proteins, MCM7 is one of the highly conserved mini-chromosome maintenance proteins (MCM) that is essential for the initiation of eukaryotic genome replication. The specificity of this monoclonal antibody to its intended target was validated by HuProTIM Array, containing more than 19,000, full-length human proteins, MCM7 is one of the highly conserved mini-chromosome maintenance proteins (MCM) that is essential for the initiation of eukaryotic genome replication. The hexameric protein complex formed by the MCM proteins is a key component of the per-replication complex formed by the MCM proteins, The MCM complex consisting of this protein and MCM2, 4 and 6 proteins possesses DNA helicase activity, and may act as a DNA unwinding enzyme. Cyclin D1- dependent kinase, CDK4, is found to associate with this protein, and may regulate the binding of this protein fragment (aa195-319) (exact s		······································
ConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalClonerMCM7/1468Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 700PurityProtein A or G purifiedBufferSom Sodium BorateProduct DescriptionMouseGene ID4176Gene ID4176Gene SymbolMCM7SpeciesHuman, CanineMarkerProliferation MarkerSpecificity/SensitivityThe specificity of this monoclonal antibody to its intended target was validated by HuProTTM Array, containing more than ind-chromosome maintenance proteins (MCM) that is essential for the initiation of explication related proteins. The MCM complex and the initiation of explication for Kas and in the recruitment of other DNA replication related proteins. The MCM complex and may be involved in the formation of replication for kas and in the recruitment of other DNA replication related proteins. The MCM protein size as DNA unwinding enzyme. Cyclin D1- dependent kinase, CDK4, is found to associate with this protein, and may regulate the binding of this protein more than flyCes. Superiori. The helicase activity, and may act as a DNA unwinding enzyme. Cyclin D1- dependent kinase, CDK4, is found to associate with this protein, and may regulate the binding of this protein more than flyCes. Superiers and MCM2 is a consisting of this protein marker supersor protein RB1/RB.ImmunogenRecombinant human MCM7 protein fragment (aa195-319) (exact sequence is	Product Information	
services.StorageStore at 4C in the dark.ClonalityMonoclonalClonerMCM7/1468Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 700PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseGene ID4176Gene SymbolMCM7SpeciesHuman, CanineMarkerProliferation MarkerSpecificity/SensitivityThe specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19,000, full-length human proteins. MCM7 is one of the highly conserved mini-chromosome maintenance proteins (MCM) that is essential for the initiation of eukaryotic genome replication. The hexameric protein complex formed by the MCM proteins is a key component of the pre-replication complex and may be involved in the formation of replication forks and in the recruitment of other DNA replication related proteins. The MCM complex consisting of this protein and MCM2, and 6 proteins RB1/RB.ImmunogenRecombinant human MCM7 protein fragment (aa195-319) (exact sequence is	Unit Size	100 ul
ClonalityMonoclonalClonerMCM7/1468Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 700PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description4176HostMouseGene ID4176Gene SymbolMCM7SpeciesHuman, CanineMarkerProliferation MarkerSpecificity/SensitivityThe specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19,000, full-length human proteins. MCM7 is one of the highly conserved mini-chromosome maintenance proteins (MCM) tai to essential for the initiation of eukaryotic genome replication. The hexameric protein complex formed by the MCM proteins is a key component of the pre-replication complex and may be involved in the formation of replication forks and in the recruitment of other DNA replication related proteins. The MCM complex consisting of this protein and MCM2, 4 and 6 proteins proteins. The MCM protein scale activity, and may act as a DNA unwinding enzyme. Cyclin D1- dependent kinase, CDK4, is found to associate with this protein, RB1/RB.ImmunogenRecombinant human MCM7 protein fragment (aa195-319) (exact sequence is	Concentration	
ClonerMCM7/1468Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 700PurityProtein A or G purifiedBufferSomM Sodium BorateProduct DescriptionMouseHostMouseGene ID4176Gene SymbolMCM7SpeciesHuman, CanineMarkerProliferation MarkerSpecificity/SensitivityThe specificity of this monoclonal antibody to its intended target was validated by HUProTTM Array, containing more than 19,000, full-length human proteins. MCM7 is one of the hiphly conserved mini-chromosome maintenance proteins (MCM) that is essential for the initiation of eukaryotic genome replication. The hexameric protein complex formed by the MCM proteins is a key component of the pre-replication complex formed by the MCM proteins. The MCM complex consisting of this protein and MCM2, 4 and 6 proteins possesses DNA helicase activity, and may act as a DNA unwinding enzyme. Cyclin D1- dependent kinase, CDK4, is found to associate with this protein RB1/RB.ImmunogenRecombinant MCM7 protein fragment (aa195-319) (exact sequence is	Storage	Store at 4C in the dark.
Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 700PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID4176Gene SymbolMCM7SpeciesHuman, CanineMarkerProliferation MarkerSpecificity/SensitivityThe specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19,000, full-length human proteins. MCM7 is one of the highly conserved mini-chromosome maintenance proteins (MCM) that is essential for the initiation of eukaryotic genome replication. The hexameric protein complex and may be involved in the formation of replication forks and in the recruitment of other DNA replication related proteins. The MCM complex consisting of this protein and MCM2, 4 and 6 proteins possesses DNA helicase activity, and may act as a DNA unwinding enzyme. Cyclin D1- dependent kinase, CDK4, is found to associate with this protein, RB1/RB.ImmunogenRecombinant human MCM7 protein fragment (aa195-319) (exact sequence is	Clonality	Monoclonal
IsotypeIgG1 KappaConjugateAlexa Fluor 700PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID4176Gene SymbolMCM7SpeciesHuman, CanineMarkerProliferation MarkerSpecificity/SensitivityThe specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19,000, full-length human proteins. MCM7 is one of the highly conserved mini-chromosome maintenance proteins (MCM) that is essential for the initiation of eukaryotic genome replication. The hexameric protein complex formed by the MCM proteins is a key component of the pre-replication complex and may be involved in the formation of replication forks and in the recruitment of other DNA replication related proteins. The MCM complex consisting of this protein and MCM2, 4 and 6 proteins possesses DNA helicase activity, and may act as a DNA unwinding enzyme. Cyclin D1- dependent kinase, CDK4, is found to associate with this protein mad may regulate the binding of this protein fragment (aa195-319) (exact sequence is	Clone	rMCM7/1468
ConjugateAlexa Fluor 700PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID4176Gene SymbolMCM7SpeciesHuman, CanineMarkerProliferation MarkerSpecificity/SensitivityThe specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19,000, full-length human proteins. MCM7 is one of the highly conserved mini-chromosome maintenance proteins (MCM) that is essential for the initiation of eukaryotic genome replication. The hexameric protein complex and may be involved in the formation of replication forks and in the recruitment of other DNA replication related proteins. The MCM complex consisting of this protein and MCM2, 4 and 6 proteins possesses DNA helicase activity, and may act as a DNA unwinding enzyme. Cyclin D1- dependent kinase, CDK4, is found to associate with this protein, and may regulate the binding of this protein mith the tumor suppressor protein RB1/RB.ImmunogenRecombinant human MCM7 protein fragment (aa195-319) (exact sequence is	Preservative	0.05% Sodium Azide
Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 4176 Gene Symbol MCM7 Species Human, Canine Marker Proliferation Marker Specificity/Sensitivity The specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19,000, full-length human proteins. MCM7 is one of the highly conserved mini-chromosome maintenance proteins (MCM) that is essential for the initiation of eukaryotic genome replication. The hexameric protein complex and may be involved in the formation of replication forks and in the recruitment of other DNA replication related proteins. The MCM complex consisting of this protein and MCM2, 4 and 6 proteins possesses DNA helicase activity, and may act as a DNA unwinding enzyme. Cyclin D1-dependent kinase, CDK4, is found to associate with this protein, and may regulate the binding of this protein with the tumor suppressor protein RB1/RB. Immunogen Recombinant human MCM7 protein fragment (aa195-319) (exact sequence is	Isotype	IgG1 Kappa
Buffer50mM Sodium BorateProduct DescriptionHostMouseGene ID4176Gene SymbolMCM7SpeciesHuman, CanineMarkerProliferation MarkerSpecificity/SensitivityThe specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19,000, full-length human proteins. MCM7 is one of the highly conserved mini-chromosome maintenance proteins (MCM) that is essential for the initiation of eukaryotic genome replication. The hexameric protein complex formed by the MCM proteins is a key component of the pre-replication complex and may be involved in the formation of replication forks and in the recruitment of other DNA replication related proteins. The MCM complex consisting of this protein and MCM2, 4 and 6 proteins possesses DNA helicase activity, and may act as a DNA unwinding enzyme. Cyclin D1- dependent kinase, CDK4, is found to associate with this protein, and may regulate the binding of this protein with the tumor suppressor protein RB1/RB.ImmunogenRecombinant human MCM7 protein fragment (aa195-319) (exact sequence is	Conjugate	Alexa Fluor 700
Product DescriptionHostMouseGene ID4176Gene SymbolMCM7SpeciesHuman, CanineMarkerProliferation MarkerSpecificity/SensitivityThe specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19,000, full-length human proteins. MCM7 is one of the highly conserved mini-chromosome maintenance proteins (MCM) that is essential for the initiation of eukaryotic genome replication. The hexameric protein complex formed by the MCM proteins is a key component of the pre-replication complex and may be involved in the formation of replication forks and in the recruitment of other DNA replication related proteins. The MCM complex consisting of this protein and MCM2, 4 and 6 proteins possesses DNA helicase activity, and may act as a DNA unwinding enzyme. Cyclin D1- dependent kinase, CDK4, is found to associate with this protein, and may regulate the binding of this protein with the tumor suppressor protein RB1/RB.ImmunogenRecombinant human MCM7 protein fragment (aa195-319) (exact sequence is	Purity	Protein A or G purified
HostMouseGene ID4176Gene SymbolMCM7SpeciesHuman, CanineMarkerProliferation MarkerSpecificity/SensitivityThe specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19,000, full-length human proteins. MCM7 is one of the highly conserved mini-chromosome maintenance proteins (MCM) that is essential for the initiation of eukaryotic genome replication. The hexameric protein complex formed by the MCM proteins is a key component of the pre-replication complex and may be involved in the formation of replication forks and in the recruitment of other DNA replication related proteins. The MCM complex consisting of this protein and MCM2, 4 and 6 proteins possesses DNA helicase activity, and may act as a DNA unwinding enzyme. Cyclin D1- dependent kinase, CDK4, is found to associate with this protein, and may regulate the binding of this protein with the tumor suppressor protein RB1/RB.ImmunogenRecombinant human MCM7 protein fragment (aa195-319) (exact sequence is	Buffer	50mM Sodium Borate
Gene ID4176Gene SymbolMCM7SpeciesHuman, CanineMarkerProliferation MarkerSpecificity/SensitivityThe specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19,000, full-length human proteins. MCM7 is one of the highly conserved mini-chromosome maintenance proteins (MCM) that is essential for the initiation of eukaryotic genome replication. The hexameric protein complex formed by the MCM proteins is a key component of the pre-replication complex and may be involved in the formation of replication forks and in the recruitment of other DNA replication related proteins. The MCM complex consisting of this protein and MCM2, 4 and 6 proteins possesses DNA helicase activity, and may act as a DNA unwinding enzyme. Cyclin D1- dependent kinase, CDK4, is found to associate with this protein, and may regulate the binding of this protein with the tumor suppressor protein RB1/RB.ImmunogenRecombinant human MCM7 protein fragment (aa195-319) (exact sequence is	Product Description	
Gene SymbolMCM7SpeciesHuman, CanineMarkerProliferation MarkerSpecificity/SensitivityThe specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19,000, full-length human proteins. MCM7 is one of the highly conserved mini-chromosome maintenance proteins (MCM) that is essential for the initiation of eukaryotic genome replication. The hexameric protein complex formed by the MCM proteins is a key component of the pre-replication complex and may be involved in the formation of replication forks and in the recruitment of other DNA replication related proteins. The MCM complex consisting of this protein and MCM2, 4 and 6 proteins possesses DNA helicase activity, and may act as a DNA unwinding enzyme. Cyclin D1- dependent kinase, CDK4, is found to associate with this protein, and may regulate the binding of this protein with the tumor suppressor protein RB1/RB.ImmunogenRecombinant human MCM7 protein fragment (aa195-319) (exact sequence is	Host	Mouse
SpeciesHuman, CanineMarkerProliferation MarkerSpecificity/SensitivityThe specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19,000, full-length human proteins. MCM7 is one of the highly conserved mini-chromosome maintenance proteins (MCM) that is essential for the initiation of eukaryotic genome replication. The hexameric protein complex formed by the MCM proteins is a key component of the pre-replication complex and may be involved in the formation of replication forks and in the recruitment of other DNA replication related proteins. The MCM complex consisting of this protein and MCM2, 4 and 6 proteins possesses DNA helicase activity, and may act as a DNA unwinding enzyme. Cyclin D1- dependent kinase, CDK4, is found to associate with this protein, and may regulate the binding of this protein with the tumor suppressor protein RB1/RB.ImmunogenRecombinant human MCM7 protein fragment (aa195-319) (exact sequence is	Gene ID	4176
MarkerProliferation MarkerSpecificity/SensitivityThe specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19,000, full-length human proteins. MCM7 is one of the highly conserved mini-chromosome maintenance proteins (MCM) that is essential for the initiation of eukaryotic genome replication. The hexameric protein complex formed by the MCM proteins is a key component of the pre-replication complex and may be involved in the formation of replication forks and in the recruitment of other DNA replication related proteins. The MCM complex consisting of this protein and MCM2, 4 and 6 proteins possesses DNA helicase activity, and may act as a DNA unwinding enzyme. Cyclin D1- dependent kinase, CDK4, is found to associate with this protein, and may regulate the binding of this protein with the tumor suppressor protein RB1/RB.ImmunogenRecombinant human MCM7 protein fragment (aa195-319) (exact sequence is	Gene Symbol	MCM7
Specificity/SensitivityThe specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19,000, full-length human proteins. MCM7 is one of the highly conserved mini-chromosome maintenance proteins (MCM) that is essential for the initiation of eukaryotic genome replication. The hexameric protein complex formed by the MCM proteins is a key component of the pre-replication complex and may be involved in the formation of replication forks and in the recruitment of other DNA replication related proteins. The MCM complex consisting of this protein and MCM2, 4 and 6 proteins possesses DNA helicase activity, and may act as a DNA unwinding enzyme. Cyclin D1- dependent kinase, CDK4, is found to associate with this protein, and may regulate the binding of this protein with the tumor suppressor protein RB1/RB.ImmunogenRecombinant human MCM7 protein fragment (aa195-319) (exact sequence is	Species	Human, Canine
HuProtTM Array, containing more than 19,000, full-length human proteins. MCM7 is one of the highly conserved mini-chromosome maintenance proteins (MCM) that is essential for the initiation of eukaryotic genome replication. The hexameric protein complex formed by the MCM proteins is a key component of the pre-replication complex and may be involved in the formation of replication forks and in the recruitment of other DNA replication related proteins. The MCM complex consisting of this protein and MCM2, 4 and 6 proteins possesses DNA helicase activity, and may act as a DNA unwinding enzyme. Cyclin D1- dependent kinase, CDK4, is found to associate with this protein, and may regulate the binding of this protein with the tumor suppressor protein RB1/RB.ImmunogenRecombinant human MCM7 protein fragment (aa195-319) (exact sequence is	Marker	Proliferation Marker
	Specificity/Sensitivity	HuProtTM Array, containing more than 19,000, full-length human proteins. MCM7 is one of the highly conserved mini-chromosome maintenance proteins (MCM) that is essential for the initiation of eukaryotic genome replication. The hexameric protein complex formed by the MCM proteins is a key component of the pre-replication complex and may be involved in the formation of replication forks and in the recruitment of other DNA replication related proteins. The MCM complex consisting of this protein and MCM2, 4 and 6 proteins possesses DNA helicase activity, and may act as a DNA unwinding enzyme. Cyclin D1- dependent kinase, CDK4, is found to associate with this protein, and may regulate the binding of this protein with the tumor suppressor protein RB1/RB.
	Immunogen	



	Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.
Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Protein Array
Recommended Dilutions	Western Blot, Immunohistochemistry-Paraffin, Protein Array
Application Notes	Optimal dilution of this antibody should be experimentally determined.

Notes





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP3-08389AF700

IC002N	Mouse IgG1 Isotype Control (11711) [Alexa Fluor® 700]
NBP2-51540-0.1mg	Recombinant Human MCM7 His Protein
NBL1-12956	MCM7 Overexpression Lysate
NB500-106	PCNA Antibody (PC10)

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-08389AF700

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

