Product Datasheet

StAR Antibody (STAR/2077) [DyLight 680] NBP3-08547FR

Unit Size: 100 ul

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-08547FR

Updated 10/26/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-08547FR



NBP3-08547FR

Application Notes

Product Information	StAR Antibody (STAR/2077) [DyLight 680]	
Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonality Monoclonal Clone STAR/2077 Preservative 0.05% Sodium Azide Isotype IgG2b Kappa Conjugate DyLight 680 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 6770 Gene Symbol STAR Species Human Marker Leydig Cell Marker Specificity/Sensitivity Steroidogenic Acute Regulatory Protein (STAR) controls the rate-limiting step of steroidegenesis by translocating cholesterol from the outer mitochondrial membrane to the inner membrane where it is later cleaved to pregnenolone. It is primarily present in steroid-producing cells, including Leydig cells in the testis, theca cells and luteal cells in the ovary and adrenal cells in the adrenal cortex. Due to low levels of pregnenolone, seminomas and Leydig cell tumors display no specific STAR staining. Therefore, STAR antibody may assist in differentiating sex cord stromal tumors (SCST), seminomas and embryonal carcinomas. Immunogen Recombinant fragment (acround as 39-108) of human StAR protein (exact sequence is proprietary) (Uniprot: P49675) Notes Immunofication Details Applications Immunohistochemistry-Paraffin	Product Information	
Storage Store at 4C in the dark. Clonality Monoclonal Clone STAR/2077 Preservative 0.05% Sodium Azide Isotype IgG2b Kappa Conjugate DyLight 680 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 6770 Gene Symbol STAR Species Human Marker Leydig Cell Marker Specificity/Sensitivity Steroidogenic Acute Regulatory Protein (STAR) controls the rate-limiting step of steroidegenesis by translocating cholesterol from the outer mitochondrial membrane to the inner membrane where it is later cleaved to pregnenolone. It is primarily present in steroid-producing cells, including Leydig cells in the testis, theca cells and luteal cells in the ovary and adrenal cells in the testis, theca cells and luteal cells in the ovary and adrenal cells in the testis, theca cells and luteal cells in the ovary and adrenal cells in the testis, theca cells and luteal cells in the ovary and adrenal cells in the faternal cortex. Due to low levels of pregnenolone, seminomas and embryonal carcinomas. Immunogen Recombinant fragment (around aa 39–108) of human StAR protein (exact sequence is proprietary) (Uniprot: P49675) Notes Immunofistochemistry-Paraffin Immunohistochemistry-Paraffin	Unit Size	100 ul
Clone STAR/2077 Preservative 0.05% Sodium Azide Isotype IgG2b Kappa Conjugate DyLight 680 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 6770 Gene Symbol STAR Species Human Marker Leydig Cell Marker Specificity/Sensitivity Steroidogenic Acute Regulatory Protein (STAR) controls the rate-limiting step of steroidegenesis by translocating cholesterol from the outer mitochondrial membrane to the inner membrane where it is later cleaved to pregnenolone. It is primarily present in steroid-producing cells, including Leydig cells in the testis, theca cells and luteal cells in the ovary and adrenal cells in the adrenal cortex. Due to low levels of pregnenolone, seminomas and Leydig cell tumors display no specific STAR staining. Therefore, STAR antibody may assist in differentiating sex cord stromal tumors (SCST), seminomas and embryonal carcinomas. Immunogen Recombinant fragment (around aa 39-108) of human StAR protein (exact sequence is proprietary) (Uniprot: P49675) Notes Immunoptication Details Applications Immunoshistochemistry-Paraffin	Concentration	· ·
Clone STAR/2077 Preservative 0.05% Sodium Azide Isotype IgG2b Kappa Conjugate DyLight 680 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 6770 Gene Symbol STAR Species Human Marker Leydig Cell Marker Specificity/Sensitivity Steroidogenic Acute Regulatory Protein (STAR) controls the rate-limiting step of steroidegenesis by translocating cholesterol from the outer mitochondrial membrane to the inner membrane where it is later cleaved to pregnenolone. It is primarily present in steroid-producing cells, including Leydig cells in the testis, theca cells and Iuteal cells in the ovary and adrenal cells in the adrenal cortex. Due to low levels of pregnenolone, seminomas and Leydig cell tumors display no specific STAR staining. Therefore, STAR antibody may assist in differentiating sex cord stromal tumors (SCST), seminomas and embryonal carcinomas. Immunogen Recombinant fragment (around aa 39-108) of human StAR protein (exact sequence is proprietary) (Uniprot: P49675) Notes Product Application Details Applications Immunohistochemistry-Paraffin	Storage	Store at 4C in the dark.
Preservative 0.05% Sodium Azide Isotype IgG2b Kappa Conjugate DyLight 680 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 6770 Gene Symbol STAR Species Human Marker Leydig Cell Marker Specificity/Sensitivity Steroidogenic Acute Regulatory Protein (STAR) controls the rate-limiting step of steroidegenesis by translocating cholesterol from the outer mitochondrial membrane to the inner membrane where it is later cleaved to pregnenolone. It is primarily present in steroid-producing cells, including Leydig cells in the testis, theca cells and luteal cells in the oavray and adreal cells in the adrenal cortex. Due to low levels of pregnenolone, seminomas and Leydig cell tumors display no specific STAR staining. Therefore, STAR antibody may assist in differentiating sex cord stormal tumors (SCST), seminomas and embryonal carcinomas. Immunogen Recombinant fragment (around aa 39-108) of human StAR protein (exact sequence is proprietary) (Uniprot: P49675) Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. Product Application Details Applications Immunohistochemistry-Paraffin	Clonality	Monoclonal
Isotype	Clone	STAR/2077
Conjugate DyLight 680 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 6770 Gene Symbol STAR Species Human Marker Leydig Cell Marker Specificity/Sensitivity Steroidogenic Acute Regulatory Protein (STAR) controls the rate-limiting step of steroidegenesis by translocating cholesterol from the outer mitochondrial membrane to the inner membrane where it is later cleaved to pregnenolone. It is primarily present in steroid-producing cells, including Leydig cells in the testis, theca cells and luteal cells in the ovary and adrenal cells in the adrenal cortex. Due to low levels of pregnenolone, seminomas and Leydig cell tumors display no specific STAR staining. Therefore, STAR antibody may assist in differentiating sex cord stromal tumors (SCST), seminomas and embryonal carcinomas. Immunogen Recombinant fragment (around aa 39-108) of human StAR protein (exact sequence is proprietary) (Uniprot: P49675) Notes Product Application Details Applications Immunohistochemistry-Paraffin	Preservative	0.05% Sodium Azide
Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 6770 Gene Symbol STAR Species Human Marker Leydig Cell Marker Specificity/Sensitivity Steroidogenic Acute Regulatory Protein (STAR) controls the rate-limiting step of steroidegenesis by translocating cholesterol from the outer mitochondrial membrane to the inner membrane where it is later cleaved to pregnenolone. It is primarily present in steroid-producing cells, including Leydig cells in the testis, theca cells and luteal cells in the ovary and adrenal cortex. Due to low levels of pregnenolone, seminomas and Leydig cell tumors display no specific STAR staining. Therefore, STAR antibody may assist in differentiating sex cord stromal tumors (SCST), seminomas and embryonal carcinomas. Immunogen Recombinant fragment (around aa 39-108) of human StAR protein (exact sequence is proprietary) (Uniprot: P49675) Notes Product Application Details Applications Immunohistochemistry-Paraffin	Isotype	IgG2b Kappa
Product Description	Conjugate	DyLight 680
Product Description Host Mouse Gene ID 6770 Gene Symbol STAR Species Human Marker Leydig Cell Marker Specificity/Sensitivity Steroidogenic Acute Regulatory Protein (STAR) controls the rate-limiting step of steroidegenesis by translocating cholesterol from the outer mitochondrial membrane to the inner membrane where it is later cleaved to pregnenolone. It is primarily present in steroid-producing cells, including Leydig cells in the testis, theca cells and luteal cells in the ovary and adrenal cells in the adrenal cortex. Due to low levels of pregnenolone, seminomas and Leydig cell tumors display no specific STAR staining. Therefore, STAR antibody may assist in differentiating sex cord stromal tumors (SCST), seminomas and embryonal carcinomas. Immunogen Recombinant fragment (around aa 39-108) of human StAR protein (exact sequence is proprietary) (Uniprot: P49675) Notes Product Application Details Applications Immunohistochemistry-Paraffin	Purity	Protein A or G purified
Host Mouse Gene ID 6770 Gene Symbol STAR Species Human Marker Leydig Cell Marker Specificity/Sensitivity Steroidogenic Acute Regulatory Protein (STAR) controls the rate-limiting step of steroidogenesis by translocating cholesterol from the outer mitochondrial membrane to the inner membrane where it is later cleaved to pregnenolone. It is primarily present in steroid-producing cells, including Leydig cells in the testis, theca cells and luteal cells in the ovary and adrenal cells in the adrenal cortex. Due to low levels of pregnenolone, seminomas and Leydig cell tumors display no specific STAR staining. Therefore, STAR antibody may assist in differentiating sex cord stromal tumors (SCST), seminomas and embryonal carcinomas. Immunogen Recombinant fragment (around aa 39-108) of human StAR protein (exact sequence is proprietary) (Uniprot: P49675) Notes Product Application Details Applications Immunohistochemistry-Paraffin	Buffer	50mM Sodium Borate
Gene Symbol STAR Species Human Marker Leydig Cell Marker Specificity/Sensitivity Steroidogenic Acute Regulatory Protein (STAR) controls the rate-limiting step of steroidegenesis by translocating cholesterol from the outer mitochondrial membrane to the inner membrane where it is later cleaved to pregnenolone. It is primarily present in steroid-producing cells, including Leydig cells in the testis, theca cells and luteal cells in the ovary and adrenal cells in the adrenal cortex. Due to low levels of pregnenolone, seminomas and Leydig cell tumors display no specific STAR staining. Therefore, STAR antibody may assist in differentiating sex cord stromal tumors (SCST), seminomas and embryonal carcinomas. Immunogen Recombinant fragment (around aa 39-108) of human StAR protein (exact sequence is proprietary) (Uniprot: P49675) Notes Product Application Details Applications Immunohistochemistry-Paraffin	Product Description	
Star	Host	Mouse
Marker Leydig Cell Marker	Gene ID	6770
Marker Specificity/Sensitivity Steroidogenic Acute Regulatory Protein (STAR) controls the rate-limiting step of steroidegenesis by translocating cholesterol from the outer mitochondrial membrane to the inner membrane where it is later cleaved to pregnenolone. It is primarily present in steroid-producing cells, including Leydig cells in the testis, theca cells and luteal cells in the ovary and adrenal cells in the adrenal cortex. Due to low levels of pregnenolone, seminomas and Leydig cell tumors display no specific STAR staining. Therefore, STAR antibody may assist in differentiating sex cord stromal tumors (SCST), seminomas and embryonal carcinomas. Immunogen Recombinant fragment (around aa 39-108) of human StAR protein (exact sequence is proprietary) (Uniprot: P49675) Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. Product Application Details Applications Immunohistochemistry-Paraffin	Gene Symbol	STAR
Specificity/Sensitivity Steroidogenic Acute Regulatory Protein (STAR) controls the rate-limiting step of steroidegenesis by translocating cholesterol from the outer mitochondrial membrane to the inner membrane where it is later cleaved to pregnenolone. It is primarily present in steroid-producing cells, including Leydig cells in the testis, theca cells and luteal cells in the ovary and adrenal cells in the adrenal cortex. Due to low levels of pregnenolone, seminomas and Leydig cell tumors display no specific STAR staining. Therefore, STAR antibody may assist in differentiating sex cord stromal tumors (SCST), seminomas and embryonal carcinomas. Immunogen Recombinant fragment (around aa 39-108) of human StAR protein (exact sequence is proprietary) (Uniprot: P49675) Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. Product Application Details Applications Immunohistochemistry-Paraffin	Species	Human
steroidegenesis by translocating cholesterol from the outer mitochondrial membrane to the inner membrane where it is later cleaved to pregnenolone. It is primarily present in steroid-producing cells, including Leydig cells in the testis, theca cells and luteal cells in the ovary and adrenal cells in the adrenal cortex. Due to low levels of pregnenolone, seminomas and Leydig cell tumors display no specific STAR staining. Therefore, STAR antibody may assist in differentiating sex cord stromal tumors (SCST), seminomas and embryonal carcinomas. Immunogen Recombinant fragment (around aa 39-108) of human StAR protein (exact sequence is proprietary) (Uniprot: P49675) Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. Product Application Details Immunohistochemistry-Paraffin	Marker	Leydig Cell Marker
Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. Product Application Details Applications Immunohistochemistry-Paraffin	Specificity/Sensitivity	steroidegenesis by translocating cholesterol from the outer mitochondrial membrane to the inner membrane where it is later cleaved to pregnenolone. It is primarily present in steroid-producing cells, including Leydig cells in the testis, theca cells and luteal cells in the ovary and adrenal cells in the adrenal cortex. Due to low levels of pregnenolone, seminomas and Leydig cell tumors display no specific STAR staining. Therefore, STAR antibody may assist in differentiating
DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. Product Application Details Applications Immunohistochemistry-Paraffin	Immunogen	
Applications Immunohistochemistry-Paraffin	Notes	DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.
·	Product Application Details	
Recommended Dilutions Immunohistochemistry-Paraffin	Applications	Immunohistochemistry-Paraffin
·	Recommended Dilutions	Immunohistochemistry-Paraffin



Optimal dilution of this antibody should be experimentally determined.



Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP3-08547FR

NBP1-43317FR Mouse IgG2b Kappa Light Chain Isotype Control (MG2b) [DyLight 680]

NBP1-72531-100ug Recombinant Human StAR His Protein

291-G1-200 IGF-I/IGF-1 [Unconjugated]
NBP2-09031 StAR Overexpression Lysate

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-08547FR

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

