Product Datasheet

Cytokeratin 3 Antibody (KRT3/2542) [CoraFluor™ 1] NBP3-08582CL1

Unit Size: 0.1 ml

Store at 4C in the dark. Do not freeze.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-08582CL1

Updated 8/13/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP3-08582CL1



NBP3-08582CL1

Cytokeratin 3 Antibody (KRT3/2542) [CoraFluor™ 1]

| Droduct Information | Cytokeratin 5 Antibody (KK 13/2 | | |
|--|---------------------------------|--|--|
| Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Do not freeze. Clonality Monoclonal Clone KRT3/2542 Preservative No Preservative Isotype IgS1 Kappa Conjugate CoraFluor 1 Purity Protein A or G purified Buffer PBS Product Description Description CoraFluor(TM) 1 is a high performance terbium-based TR-FRET (Time-Resolved Fluorescence) Resonance Energy Transfer) or TRF (Time-Resolved Fluorescence) donor for high throughput assay development. CoraFluor(TM) 1 absorbs UV light at approximately 340 nm, and emits at approximately 490 nm, 545 nm, 355 nm and 620 nm. It is compatible with common acceptor dyse that absorb at the emission wavelengins of CoraFluor(TM) 1. CoraFluor(TM) 1 corafl | Product Information | | |
| Storage Store at 4C in the dark. Do not freeze. Clonality Monoclonal Clone KRT3/2542 Preservative No Preservative Isotype IgG1 Kappa Conjugate CoraFluor 1 Purity Protein A or G purified Buffer PBS Product Description Description CoraFluor(TM) 1 is a high performance terbium-based TR-FRET (Time-Resolved Fluorescence Resonance Energy Transfer) or TRF (Time-Resolved Fluorescence) donor for high throughput assay development. CoraFluor(TM) 1 absorbs UV light at approximately 340 nm, and emits at approximately 490 nm, 545 nm, 585 nm and 620 nm. It is compatible with common acceptor dyes that absorb at the emission wavelengths of CoraFluor(TM) 1. CoraFluor(TM) 1 can be used for the development of robust and scalable TR-FRET binding assays such as target engagement, ternary complex, protein-protein interaction and protein quantification assays. CoraFluor(TM) 1, amine reactive CoraFluor(TM) 1, amine reactive CoraFluor(TM) 1, thiol reactive For more information, please see our CoraFluor(TM) TR-FRET technology flyer. Host Mouse Gene ID 3850 Gene Symbol KRT3 Species Human, Bovine, Rabbit Marker Corneal Epithelial Marker Specificity/Sensitivity The protein encoded by this gene is a member of the keratin gene family. The type II cytokeratins consists of basic or neutral proteins, which are arranged in pairs of heterotypic keratin chains co-expressed during differentiation of simple and stratified epithelial tissues. This type II cytokeratin is specifically expressed in the corneal epithelium with family member KRT12 and mutations in these genes have been associated with Meesmanns Corneal Dystrophy. KRT3/25542 recognizes the 646t D polypeptic/elveratin K2) of palate epithelium as a rabbit corneal epithelium and keratin K76 (cytokeratin 3) of rabbit corneal epithelium and keratin K76 (cytokeratin 3) of rabbit corneal epithelium and keratin K76 (cytokeratin 3) of rabbit corneal epithelium. | Unit Size | 0.1 ml | |
| Clone KRT3/2542 Preservative No Preservative Isotype IgG1 Kappa Conjugate CoraFluor 1 Purity Protein A or G purified Buffer PBS Product Description CoraFluor(TM) 1 is a high performance terbium-based TR-FRET (Time-Resolved Fluorescence) donor for high throughput assay development. CoraFluor(TM) 1 absorbs UV light at approximately 490 nm, and emits at approximately 490 nm, 545 nm, 585 nm and 520 nm. It is compatible with common acceptor dyes that absorb at the emission wavelengths of CoraFluor(TM) 1 can be used for the development of robust and scalable TR-FRET binding assays such as larget engagement, ternary complex, protein-protein interaction and protein quantification assays. CoraFluor(TM) 1. amine reactive CoraFluor(TM) 1. thiol reactive For more information, please see our CoraFluor(TM) TR-FRET technology flyer. Host Mouse Gene ID 3850 Gene Symbol KRT3 Species Human, Bovine, Rabbit Marker Corneal Epithelial Marker Specificity/Sensitivity The protein encoded by this gene is a member of the keratin gene family. The type II cytokeratins consists of basic or neutral proteins, which are arranged in pairs of heterotypic keratin chains co-expressed during differentiation of simple and stratified epithelial tissues. This type II cytokeratin is specifically expressed in the corneal epithelium with family member KRT12 and mustors in these genes have been associated with Meesmanns Corneal Dystrophy.KRT3/2542 recognizes the 64kD polypeptic/keratin K2p) of palate epithelium. Also recognizes be ovine snout and lip epithelium, as well as a rabbit corneal cell line. Immunogen | Concentration | · · | |
| Clone KRT3/2542 Preservative No Preservative Isotype IgG1 Kappa Conjugate CoraFluor 1 Purity Protein A or G purified Buffer PBS Product Description CoraFluor(TM) 1 is a high performance terbium-based TR-FRET (Time-Resolved Fluorescence Resonance Energy Transfer) or TRF (Time-Resolved Fluorescence) donor for high throughput assay development. CoraFluor(TM) 1 absorbs UV light at approximately 340 nm, and emits at approximately 430 nm, S45 nm, S45 nm, S65 nm and 620 nm. It is compatible with common acceptor dyes that absorb at the emission wavelengths of CoraFluor(TM) 1. CoraFluor(TM) 1 can be used for the development of robust and scalable TR-FRET binding assays such as target engagement, ternary complex, protein-protein interaction and protein quantification assays. CoraFluor(TM) 1. amine reactive CoraFluor(TM) 1. amine reactive CoraFluor(TM) 1. amine reactive For more information, please see our CoraFluor(TM) TR-FRET technology flyer. Host Mouse Gene ID 3850 Gene Symbol KRT3 Species Human, Bovine, Rabbit Marker Corneal Epithelial Marker Specificity/Sensitivity The protein encoded by this gene is a member of the keratin gene family. The type II cytokeratin consist of basic or neutral proteins, which are arranged in pairs of heterotypic keratin chains co-expressed during differentiation of simple and stratified epithelial tissues. This type II cytokeratin is specifically expressed in the corneal epithelium with family member KRT12 and mutations in these genes have been associated with Meesmanns Corneal Dystrophy. KRT3/2542 recognizes the 64kD polypepticle/keratin K3 (cytokeratin K3 pod ribbit corneal epithelium and keratin K76 (cytokeratin K20) of palate epithelium. As such represents an excellent marker for studying corneal epithelial-type differentiation. Also recognizes bovine snout and lip epithelium, as well | Storage | Store at 4C in the dark. Do not freeze. | |
| Preservative Isotype IgG1 Kappa | Clonality | Monoclonal | |
| IgG1 Kappa IgG1 Kappa | Clone | KRT3/2542 | |
| Conjugate CoraFluor 1 Purity Protein A or G purified Buffer PBS Product Description CoraFluor(TM) 1 is a high performance terbium-based TR-FRET (Time-Resolved Fluorescence) donor for high throughput assay development. CoraFluor(TM) 1 absorbs UV light at approximately 340 mn, and emits at approximately 490 mn, 545 mn, 585 mm and 620 mn. It is compatible with common acceptor dyes that absorb at the emission wavelengths of CoraFluor(TM) 1. CoraFluor(TM) 1 can be used for the development of robust and scalable TR-FRET binding assays such as target engagement, ternary complex, protein-protein interaction and protein quantification assays. CoraFluor(TM) 1, amine reactive CoraFluor(TM) 1, thiol reactive For more information, please see our CoraFluor(TM) TR-FRET technology flyer. Host Mouse Gene ID 3850 Gene Symbol KRT3 Species Human, Bovine, Rabbit Marker Corneal Epithelial Marker Specificity/Sensitivity The protein encoded by this gene is a member of the keratin gene family. The type II cytokeratins consist of basic or neutral proteins, which are arranged in pairs of heterotypic keratin chains co-expressed during differentiation of simple and stratified epithelial itsues. This type II cytokeratin sis specifically expressed in the corneal epithelium with family member KRT12 and mutations in these genes have been associated with Meesmanns Corneal Dystrophy. KRT3/2542 recognizes the 64kD polypeptide/keratin K2p) of palate epithelium. As such represents an excellent marker for studying corneal epithelium. As such represents an excellent marker for studying corneal epithelium, as well as a rabbit corneal cell line. Immunogen | Preservative | No Preservative | |
| Purity Protein A or G purified Buffer PBS Product Description CoraFluor(TM) 1 is a high performance terbium-based TR-FRET (Time-Resolved Fluorescence) donor for high throughput assay development. CoraFluor(TM) 1 absorbs UV light at approximately 340 nm, and emits at approximately 490 nm, 545 nm, 585 nm and 620 nm, it is compatible with common acceptor dyes that absorb at the emission wavelengths of CoraFluor(TM) 1. CoraFluor(TM) 1 can be used for the development of robust and scalable TR-FRET binding assays such as target engagement, ternary complex, protein-protein interaction and protein quantification assays. CoraFluor(TM) 1. amine reactive CoraFluor(TM) 1. thiol reactive For more information, please see our CoraFluor(TM) TR-FRET technology flyer. Host Mouse Gene ID 3850 Gene Symbol KRT3 Species Human, Bovine, Rabbit Marker Comeal Epithelial Marker The protein encoded by this gene is a member of the keratin gene family. The type II cytokeratins consist of basic or neutral proteins, which are arranged in pairs of heterotypic keratin chains co-expressed during differentiation of simple and stratified epithelial tissues. This type II cytokeratins is specifically expressed in the corneal epithelium with family member KRT12 and mutations in these genes have been associated with Meesmanns Corneal polysrophy. KRT3/2542 recognizes the 64kD polypeptide/keratin K2p) of palate epithelium. As such represents an excellent marker for studying corneal epithelium. As such represents an excellent marker for studying corneal epithelium, as well as a rabbit corneal cell line. Immunogen | Isotype | IgG1 Kappa | |
| PBS | Conjugate | CoraFluor 1 | |
| Product Description CoraFluor(TM) 1 is a high performance terbium-based TR-FRET (Time-Resolved Fluorescence) donor for high throughput assay development. CoraFluor(TM) 1 absorbs UV light at approximately 340 nm, and emits at approximately 490 nm, 545 nm, 585 nm and 620 nm. It is compatible with common acceptor dyes that absorb at the emission wavelengths of CoraFluor(TM) 1. CoraFluor(TM) 1 can be used for the development of robust and scalable TR-FRET binding assays such as target engagement, ternary complex, protein-protein interaction and protein quantification assays. CoraFluor(TM) 1, amine reactive CoraFluor(TM) 1, thiol reactive For more information, please see our CoraFluor(TM) TR-FRET technology flyer. Host Mouse Gene ID 3850 Gene Symbol KRT3 Species Human, Bovine, Rabbit Marker Corneal Epithelial Marker Specificity/Sensitivity The protein encoded by this gene is a member of the keratin gene family. The type II cytokeratins consist of basic or neutral proteins, which are arranged in pairs of heterotypic keratin chains co-expressed during differentiation of simple and stratified epithelial tissues. This type II cytokeratin is specifically expressed in the corneal epithelium with family member KRT12 and mutations in these genes have been associated with Meesmanns Corneal Dystrophy. KRT3/2542 recognizes the 64kD polypeptide/keratin K3 (cytokeratin 3) of rabbit corneal epithelium and keratin K76 (cytokeratin K2p) of palate epithelium. As such represents an excellent marker for studying corneal epithelium. As such represents an excellent marker for studying corneal epithelium. As such represents an excellent marker for studying corneal epithelium. As such represents an excellent marker for studying corneal epithelium. As such represents an excellent marker for studying corneal epithelium. As such | Purity | Protein A or G purified | |
| CoraFluor(TM) 1 is a high performance terbium-based TR-FRET (Time-Resolved Fluorescence Resonance Energy Transfer) or TRF (Time-Resolved Fluorescence) donor for high throughput assay development. CoraFluor(TM) 1 absorbs UV light at approximately 340 nm, and emits at approximately 490 nm, 545 nm, 585 nm and 620 nm. It is compatible with common acceptor dyes that absorb at the emission wavelengths of CoraFluor(TM) 1. CoraFluor(TM) 1 can be used for the development of robust and scalable TR-FRET binding assays such as target engagement, ternary complex, protein-protein interaction and protein quantification assays. CoraFluor(TM) 1, amine reactive | Buffer | PBS | |
| CoraFluor(TM) 1 is a high performance terbium-based TR-FRET (Time-Resolved Fluorescence Resonance Energy Transfer) or TRF (Time-Resolved Fluorescence) donor for high throughput assay development. CoraFluor(TM) 1 absorbs UV light at approximately 340 nm, and emits at approximately 490 nm, 545 nm, 585 nm and 620 nm. It is compatible with common acceptor dyes that absorb at the emission wavelengths of CoraFluor(TM) 1. CoraFluor(TM) 1 can be used for the development of robust and scalable TR-FRET binding assays such as target engagement, ternary complex, protein-protein interaction and protein quantification assays. CoraFluor(TM) 1, amine reactive | Product Description | | |
| Gene Symbol KRT3 Species Human, Bovine, Rabbit Corneal Epithelial Marker Specificity/Sensitivity The protein encoded by this gene is a member of the keratin gene family. The type II cytokeratins consist of basic or neutral proteins, which are arranged in pairs of heterotypic keratin chains co-expressed during differentiation of simple and stratified epithelial tissues. This type II cytokeratin is specifically expressed in the corneal epithelium with family member KRT12 and mutations in these genes have been associated with Meesmanns Corneal Dystrophy.KRT3/2542 recognizes the 64kD polypeptide/keratin K3 (cytokeratin 3) of rabbit corneal epithelium and keratin K76 (cytokeratin K2p) of palate epithelium. As such represents an excellent marker for studying corneal epithelial-type differentiation. Also recognizes bovine snout and lip epithelium, as well as a rabbit corneal cell line. Immunogen Tissue, cells or virus corresponding to Rabbit Cytokeratin 3/CK-3. Rabbit corneal | Description | Fluorescence Resonance Energy Transfer) or TRF (Time-Resolved Fluorescence) donor for high throughput assay development. CoraFluor(TM) 1 absorbs UV light at approximately 340 nm, and emits at approximately 490 nm, 545 nm, 585 nm and 620 nm. It is compatible with common acceptor dyes that absorb at the emission wavelengths of CoraFluor(TM) 1. CoraFluor(TM) 1 can be used for the development of robust and scalable TR-FRET binding assays such as target engagement, ternary complex, protein-protein interaction and protein quantification assays. CoraFluor(TM) 1, amine reactive CoraFluor(TM) 1, thiol reactive | |
| Species | Host | | |
| Species Human, Bovine, Rabbit Corneal Epithelial Marker Specificity/Sensitivity The protein encoded by this gene is a member of the keratin gene family. The type II cytokeratins consist of basic or neutral proteins, which are arranged in pairs of heterotypic keratin chains co-expressed during differentiation of simple and stratified epithelial tissues. This type II cytokeratin is specifically expressed in the corneal epithelium with family member KRT12 and mutations in these genes have been associated with Meesmanns Corneal Dystrophy.KRT3/2542 recognizes the 64kD polypeptide/keratin K3 (cytokeratin 3) of rabbit corneal epithelium and keratin K76 (cytokeratin K2p) of palate epithelium. As such represents an excellent marker for studying corneal epithelial-type differentiation. Also recognizes bovine snout and lip epithelium, as well as a rabbit corneal cell line. Immunogen Tissue, cells or virus corresponding to Rabbit Cytokeratin 3/CK-3. Rabbit corneal | Gene ID | 3850 | |
| Marker Corneal Epithelial Marker The protein encoded by this gene is a member of the keratin gene family. The type II cytokeratins consist of basic or neutral proteins, which are arranged in pairs of heterotypic keratin chains co-expressed during differentiation of simple and stratified epithelial tissues. This type II cytokeratin is specifically expressed in the corneal epithelium with family member KRT12 and mutations in these genes have been associated with Meesmanns Corneal Dystrophy.KRT3/2542 recognizes the 64kD polypeptide/keratin K3 (cytokeratin 3) of rabbit corneal epithelium and keratin K76 (cytokeratin K2p) of palate epithelium. As such represents an excellent marker for studying corneal epithelial-type differentiation. Also recognizes bovine snout and lip epithelium, as well as a rabbit corneal cell line. Immunogen Tissue, cells or virus corresponding to Rabbit Cytokeratin 3/CK-3. Rabbit corneal | Gene Symbol | KRT3 | |
| The protein encoded by this gene is a member of the keratin gene family. The type II cytokeratins consist of basic or neutral proteins, which are arranged in pairs of heterotypic keratin chains co-expressed during differentiation of simple and stratified epithelial tissues. This type II cytokeratin is specifically expressed in the corneal epithelium with family member KRT12 and mutations in these genes have been associated with Meesmanns Corneal Dystrophy.KRT3/2542 recognizes the 64kD polypeptide/keratin K3 (cytokeratin 3) of rabbit corneal epithelium and keratin K76 (cytokeratin K2p) of palate epithelium. As such represents an excellent marker for studying corneal epithelial-type differentiation. Also recognizes bovine snout and lip epithelium, as well as a rabbit corneal cell line. Immunogen Tissue, cells or virus corresponding to Rabbit Cytokeratin 3/CK-3. Rabbit corneal | Species | Human, Bovine, Rabbit | |
| type II cytokeratins consist of basic or neutral proteins, which are arranged in pairs of heterotypic keratin chains co-expressed during differentiation of simple and stratified epithelial tissues. This type II cytokeratin is specifically expressed in the corneal epithelium with family member KRT12 and mutations in these genes have been associated with Meesmanns Corneal Dystrophy.KRT3/2542 recognizes the 64kD polypeptide/keratin K3 (cytokeratin 3) of rabbit corneal epithelium and keratin K76 (cytokeratin K2p) of palate epithelium. As such represents an excellent marker for studying corneal epithelial-type differentiation. Also recognizes bovine snout and lip epithelium, as well as a rabbit corneal cell line. Immunogen Tissue, cells or virus corresponding to Rabbit Cytokeratin 3/CK-3. Rabbit corneal | Marker | Corneal Epithelial Marker | |
| | Specificity/Sensitivity | The protein encoded by this gene is a member of the keratin gene family. The type II cytokeratins consist of basic or neutral proteins, which are arranged in pairs of heterotypic keratin chains co-expressed during differentiation of simple and stratified epithelial tissues. This type II cytokeratin is specifically expressed in the corneal epithelium with family member KRT12 and mutations in these genes have been associated with Meesmanns Corneal Dystrophy.KRT3/2542 recognizes the 64kD polypeptide/keratin K3 (cytokeratin 3) of rabbit corneal epithelium and keratin K76 (cytokeratin K2p) of palate epithelium. As such represents an excellent marker for studying corneal epithelial-type differentiation. Also recognizes bovine snout and lip epithelium, as well as a rabbit corneal cell | |
| | Immunogen | | |



| Notes | CoraFluor (TM) is a trademark of Bio-Techne Corp. Sold for research purposes only under agreement from Massachusetts General Hospital. US patent 2022/0025254 |
|------------------------------------|---|
| Product Application Details | |
| Applications | Western Blot, Immunohistochemistry-Paraffin |
| Recommended Dilutions | Western Blot, Immunohistochemistry-Paraffin |
| Application Notes | Optimal dilution of this antibody should be experimentally determined. |





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP3-08582CL1

H00003850-Q01-10ug Recombinant Human Cytokeratin 3 GST (N-Term) Protein

236-EG-200 EGF [Unconjugated]

NB100-355 RPE65 Antibody (401.8B11.3D9) - BSA Free

NBP2-29429 Cytokeratin, pan Antibody (AE-1/AE-3)

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-08582CL1

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

