# **Product Datasheet**

# Bcl-10 Antibody (BL10/2988R) [Alexa Fluor® 488] NBP3-08803AF488

Unit Size: 100 ul

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-08803AF488

Updated 10/26/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP3-08803AF488



# NBP3-08803AF488

Bcl-10 Antibody (BL10/2988R) [Alexa Fluor® 488]

Product Information         Unit Size         100 ul           Concentration         Please see the vial label for concentration. If unlisted please contact technical services.           Storage         Store at 4C in the dark.           Clonality         Monoclonal           Clone         BL10/2988R           Preservative         0.05% Sodium Azide           Isotype         IgG           Conjugate         Alexa Fluor 488           Purity         Protein A or G purified           Buffer         50mM Sodium Borate           Product Description           Host         Rabbit           Gene ID         8915           Gene Symbol         BCL10           Specific         Human           Marker         MALT-Lymphoma Marker           Specificity/Sensitivity         BCL10, with an N-terminal caspase recruitment domain (CARD), is found in a number of apoptotic regulatory molecules. It was identified through its direct involvement in t(1;14) of mucosa-associated lymphoid tissue (MALT) lymphoma. Expression of BCL10 was shown to induce NFB activation in a NIK-dependent pathway. This monoclonal antibody labels subpopulations of normal B and T cells and is a useful tool for the sub-classification of lymphomas. In MALT lymphomas in MALT lymphomas with the t(1;14) thranslocation, while 55% of MALT lymphomas lacking this translocation exhibited the same labeling pattern, although at a much lower level. <th< th=""><th>DCI-10 Antibody (DL 10/29001X) [A</th><th colspan="3">BCI-10 Altibudy (BE10/2900K) [Alexa Fluot® 400]</th></th<>	DCI-10 Antibody (DL 10/29001X) [A	BCI-10 Altibudy (BE10/2900K) [Alexa Fluot® 400]		
Concentration  Please see the vial label for concentration. If unlisted please contact technical services.  Storage  Store at 4C in the dark.  Clonality  Monoclonal  Clone  BL10/2988R  Preservative  0.05% Sodium Azide  Isotype  IgG  Conjugate  Alexa Fluor 488  Purity  Protein A or G purified  Buffer  50mM Sodium Borate  Product Description  Host  Rabbit  Gene ID  8915  Gene Symbol  BCL10  Species  Human  Marker  MALT-Lymphoma Marker  Specificity/Sensitivity  BCL10, with an N-terminal caspase recruitment domain (CARD), is found in a number of apoptotic regulatory molecules. It was identified through its direct involvement in t(1;14) of mucosa-associated hymphoid tissue (MALT) lymphoma. Expression of BCL10 was shown to induce NF B activation in a NIK-dependent pathway. This monoclonal antibody labels subpopulations of normal B and T cells and is a useful tool for the sub-classification of lymphomas. In MALT lymphomas with the t(1;14) translocation, while 55% of MALT lymphomas lacking this translocation exhibited the same labeling pattern, although at a much lower level.  Immunogen	Product Information	Product Information		
Storage Store at 4C in the dark.  Clonality Monoclonal  Clone BL10/2988R  Preservative 0.05% Sodium Azide  Isotype IgG  Conjugate Alexa Fluor 488  Purity Protein A or G purified  Buffer 50mM Sodium Borate  Product Description  Host Rabbit Rabbit  Gene ID 8915  Gene Symbol BCL10  Species Human  Marker MALT-Lymphoma Marker  Specificity/Sensitivity BCL10, with an N-terminal caspase recruitment domain (CARD), is found in a number of apoptotic regulatory molecules. It was identified through its direct involvement in t(1;14) of mucosa-associated lymphoid tissue (MALT) lymphoma. Expression of BCL10 was shown to induce NF B activation in a NIK-dependent pathway. This monoclonal antibody labels subpolutations of normal B and T cells and is a useful tool for the sub-classification of lymphomas. In MALT lymphomas with the t(1;14) translocation, while 55% of MALT lymphomas lacking this translocation exhibited the same labeling pattern, although at a much lower level.  Immunogen Recombinant human Bcl-10 protein fragment (around aa122-168) (exact	Unit Size	100 ul		
Clone BL10/2988R  Preservative 0.05% Sodium Azide  Isotype IgG  Conjugate Alexa Fluor 488  Purity Protein A or G purified  Buffer 50mM Sodium Borate  Product Description  Host Rabbit  Gene ID 8915  Gene Symbol BCL10  Species Human  Marker MALT-Lymphoma Marker  Specificity/Sensitivity  BCL10, with an N-terminal caspase recruitment domain (CARD), is found in a number of apoptotic regulatory molecules. It was identified through its direct involvement in t(1;14) of mucosa-associated lymphoid tissue (MALT) lymphoma. Expression of BCL10 was shown to induce NF B activation in a NIK-dependent pathway. This monoclonal antibody labels subpopulations of normal B and T cells and is a useful tool for the sub-classification of lymphomas. In MALT lymphomas with the t(1;14) translocation, while 55% of MALT lymphomas lacking this translocation exhibited the same labeling pattern, although at a much lower level.  Immunogen  Recombinant human Bcl-10 protein fragment (around aa122-168) (exact	Concentration	· ·		
Clone BL10/2988R  Preservative 0.05% Sodium Azide  Isotype IgG  Conjugate Alexa Fluor 488  Purity Protein A or G purified  Buffer 50mM Sodium Borate  Product Description  Host Rabbit Rabbit  Gene ID 8915  Gene Symbol BCL10  Species Human  Marker MALT-Lymphoma Marker  Specificity/Sensitivity BCL10, with an N-terminal caspase recruitment domain (CARD), is found in a number of apoptotic regulatory molecules. It was identified through its direct involvement in t(1;14) of mucosa-associated lymphoid tissue (MALT) lymphoma. Expression of BCL10 was shown to induce NF B activation in a NIK-dependent pathway. This monoclonal antibody labels subpopulations of normal B and T cells and is a useful tool for the sub-classification of lymphomas. In MALT lymphomas with the t(1;14) translocation, while 55% of MALT lymphomas lacking this translocation exhibited the same labeling pattern, although at a much lower level.  Immunogen Recombinant human Bcl-10 protein fragment (around aa122-168) (exact	Storage	Store at 4C in the dark.		
Preservative 0.05% Sodium Azide  Isotype IgG  Conjugate Alexa Fluor 488  Purity Protein A or G purified  Buffer 50mM Sodium Borate  Product Description  Host Rabbit Rabbit  Gene ID 8915  Gene Symbol BCL10  Species Human  Marker MALT-Lymphoma Marker  Specificity/Sensitivity BCL10, with an N-terminal caspase recruitment domain (CARD), is found in a number of apoptotic regulatory molecules. It was identified through its direct involvement in t(1;14) of mucosa-associated lymphoid tissue (MALT) lymphoma. Expression of BCL10 was shown to induce NF B activation in a NIK-dependent pathway. This monoclonal antibody labels subpopulations of normal B and T cells and is a useful tool for the sub-classification of lymphomas. In MALT lymphomas with the t(1;14) translocation, while 55% of MALT lymphomas lacking this translocation exhibited the same labeling pattern, although at a much lower level.  Immunogen Recombinant human Bcl-10 protein fragment (around aa122-168) (exact	Clonality	Monoclonal		
Isotype IgG Conjugate Alexa Fluor 488 Purity Protein A or G purified Buffer 50mM Sodium Borate  Product Description Host Rabbit Gene ID 8915 Gene Symbol BCL10 Species Human Marker MALT-Lymphoma Marker  Specificity/Sensitivity BCL10, with an N-terminal caspase recruitment domain (CARD), is found in a number of apoptotic regulatory molecules. It was identified through its direct involvement in t(1;14) of mucosa-associated lymphoid tissue (MALT) lymphoma. Expression of BCL10 was shown to induce NF B activation in a NIK-dependent pathway. This monoclonal antibody labels subpopulations of normal B and T cells and is a useful tool for the sub-classification of lymphomas. In MALT lymphomas with the t(1;14) translocation, while 55% of MALT lymphomas lacking this translocation exhibited the same labeling pattern, although at a much lower level.  Immunogen Recombinant human Bcl-10 protein fragment (around aa122-168) (exact	Clone	BL10/2988R		
Conjugate Alexa Fluor 488  Purity Protein A or G purified Buffer 50mM Sodium Borate  Product Description Host Rabbit Gene ID 8915 Gene Symbol BCL10 Species Human Marker MALT-Lymphoma Marker Specificity/Sensitivity BCL10, with an N-terminal caspase recruitment domain (CARD), is found in a number of apoptotic regulatory molecules. It was identified through its direct involvement in t(1;14) of mucosa-associated lymphoid tissue (MALT) lymphoma. Expression of BCL10 was shown to induce NF B activation in a NIK-dependent pathway. This monoclonal antibody labels subpopulations of normal B and T cells and is a useful tool for the sub-classification of lymphomas. In MALT lymphomas with the t(1;14) translocation, while 55% of MALT lymphomas lacking this translocation exhibited the same labeling pattern, although at a much lower level.  Immunogen Recombinant human Bcl-10 protein fragment (around aa122-168) (exact	Preservative	0.05% Sodium Azide		
Purity Protein A or G purified  Buffer 50mM Sodium Borate  Product Description  Host Rabbit  Gene ID 8915  Gene Symbol BCL10  Species Human  Marker MALT-Lymphoma Marker  Specificity/Sensitivity BCL10, with an N-terminal caspase recruitment domain (CARD), is found in a number of apoptotic regulatory molecules. It was identified through its direct involvement in t(1;14) of mucosa-associated lymphoid tissue (MALT) lymphoma. Expression of BCL10 was shown to induce NF B activation in a NIK-dependent pathway. This monoclonal antibody labels subpopulations of normal B and T cells and is a useful tool for the sub-classification of lymphomas. In MALT lymphomas with the t(1;14) translocation, while 55% of MALT lymphomas lacking this translocation exhibited the same labeling pattern, although at a much lower level.  Immunogen Recombinant human Bcl-10 protein fragment (around aa122-168) (exact	Isotype	IgG		
Buffer 50mM Sodium Borate  Product Description Host Rabbit Gene ID 8915 Gene Symbol BCL10 Species Human  Marker MALT-Lymphoma Marker Specificity/Sensitivity BCL10, with an N-terminal caspase recruitment domain (CARD), is found in a number of apoptotic regulatory molecules. It was identified through its direct involvement in t(1;14) of mucosa-associated lymphoid tissue (MALT) lymphoma. Expression of BCL10 was shown to induce NF B activation in a NIK-dependent pathway. This monoclonal antibody labels subpopulations of normal B and T cells and is a useful tool for the sub-classification of lymphomas. In MALT lymphomas with the t(1;14) translocation, while 55% of MALT lymphomas lacking this translocation exhibited the same labeling pattern, although at a much lower level.  Immunogen Recombinant human Bcl-10 protein fragment (around aa122-168) (exact	Conjugate	Alexa Fluor 488		
Product Description  Host Rabbit  Gene ID 8915  Gene Symbol BCL10  Species Human  Marker MALT-Lymphoma Marker  Specificity/Sensitivity BCL10, with an N-terminal caspase recruitment domain (CARD), is found in a number of apoptotic regulatory molecules. It was identified through its direct involvement in t(1;14) of mucosa-associated lymphoid tissue (MALT) lymphoma. Expression of BCL10 was shown to induce NF B activation in a NIK-dependent pathway. This monoclonal antibody labels subpopulations of normal B and T cells and is a useful tool for the sub-classification of lymphomas. In MALT lymphomas with the t(1;14) translocation, while 55% of MALT lymphomas lacking this translocation exhibited the same labeling pattern, although at a much lower level.  Immunogen Recombinant human Bcl-10 protein fragment (around aa122-168) (exact	Purity	Protein A or G purified		
Host Rabbit  Gene ID 8915  Gene Symbol BCL10  Species Human  Marker MALT-Lymphoma Marker  Specificity/Sensitivity BCL10, with an N-terminal caspase recruitment domain (CARD), is found in a number of apoptotic regulatory molecules. It was identified through its direct involvement in t(1;14) of mucosa-associated lymphoid tissue (MALT) lymphoma. Expression of BCL10 was shown to induce NF B activation in a NIK-dependent pathway. This monoclonal antibody labels subpopulations of normal B and T cells and is a useful tool for the sub-classification of lymphomas. In MALT lymphomas with the t(1;14) translocation, while 55% of MALT lymphomas lacking this translocation exhibited the same labeling pattern, although at a much lower level.  Immunogen Recombinant human Bcl-10 protein fragment (around aa122-168) (exact	Buffer	50mM Sodium Borate		
Gene Symbol  BCL10  Species  Human  Marker  MALT-Lymphoma Marker  Specificity/Sensitivity  BCL10, with an N-terminal caspase recruitment domain (CARD), is found in a number of apoptotic regulatory molecules. It was identified through its direct involvement in t(1;14) of mucosa-associated lymphoid tissue (MALT) lymphoma. Expression of BCL10 was shown to induce NF B activation in a NIK-dependent pathway. This monoclonal antibody labels subpopulations of normal B and T cells and is a useful tool for the sub-classification of lymphomas. In MALT lymphomas with the t(1;14) translocation, while 55% of MALT lymphomas lacking this translocation exhibited the same labeling pattern, although at a much lower level.  Immunogen  Recombinant human Bcl-10 protein fragment (around aa122-168) (exact	Product Description			
Gene Symbol  BCL10  Species  Human  Marker  MALT-Lymphoma Marker  Specificity/Sensitivity  BCL10, with an N-terminal caspase recruitment domain (CARD), is found in a number of apoptotic regulatory molecules. It was identified through its direct involvement in t(1;14) of mucosa-associated lymphoid tissue (MALT) lymphoma. Expression of BCL10 was shown to induce NF B activation in a NIK-dependent pathway. This monoclonal antibody labels subpopulations of normal B and T cells and is a useful tool for the sub-classification of lymphomas. In MALT lymphomas with the t(1;14) translocation, while 55% of MALT lymphomas lacking this translocation exhibited the same labeling pattern, although at a much lower level.  Immunogen  Recombinant human Bcl-10 protein fragment (around aa122-168) (exact	Host	Rabbit		
Specificity/Sensitivity  BCL10, with an N-terminal caspase recruitment domain (CARD), is found in a number of apoptotic regulatory molecules. It was identified through its direct involvement in t(1;14) of mucosa-associated lymphoid tissue (MALT) lymphoma. Expression of BCL10 was shown to induce NF B activation in a NIK-dependent pathway. This monoclonal antibody labels subpopulations of normal B and T cells and is a useful tool for the sub-classification of lymphomas. In MALT lymphomas with the t(1;14) translocation, while 55% of MALT lymphomas lacking this translocation exhibited the same labeling pattern, although at a much lower level.  Immunogen  Recombinant human Bcl-10 protein fragment (around aa122-168) (exact	Gene ID	8915		
Marker  Specificity/Sensitivity  BCL10, with an N-terminal caspase recruitment domain (CARD), is found in a number of apoptotic regulatory molecules. It was identified through its direct involvement in t(1;14) of mucosa-associated lymphoid tissue (MALT) lymphoma. Expression of BCL10 was shown to induce NF B activation in a NIK-dependent pathway. This monoclonal antibody labels subpopulations of normal B and T cells and is a useful tool for the sub-classification of lymphomas. In MALT lymphomas with the t(1;14) translocation, while 55% of MALT lymphomas lacking this translocation exhibited the same labeling pattern, although at a much lower level.  Immunogen  Recombinant human Bcl-10 protein fragment (around aa122-168) (exact	Gene Symbol	BCL10		
Specificity/Sensitivity  BCL10, with an N-terminal caspase recruitment domain (CARD), is found in a number of apoptotic regulatory molecules. It was identified through its direct involvement in t(1;14) of mucosa-associated lymphoid tissue (MALT) lymphoma. Expression of BCL10 was shown to induce NF B activation in a NIK-dependent pathway. This monoclonal antibody labels subpopulations of normal B and T cells and is a useful tool for the sub-classification of lymphomas. In MALT lymphomas with the t(1;14) translocation, while 55% of MALT lymphomas lacking this translocation exhibited the same labeling pattern, although at a much lower level.  Immunogen  Recombinant human Bcl-10 protein fragment (around aa122-168) (exact	Species	Human		
number of apoptotic regulatory molecules. It was identified through its direct involvement in t(1;14) of mucosa-associated lymphoid tissue (MALT) lymphoma. Expression of BCL10 was shown to induce NF B activation in a NIK-dependent pathway. This monoclonal antibody labels subpopulations of normal B and T cells and is a useful tool for the sub-classification of lymphomas. In MALT lymphomas with the t(1;14) translocation, while 55% of MALT lymphomas lacking this translocation exhibited the same labeling pattern, although at a much lower level.  Immunogen  Recombinant human Bcl-10 protein fragment (around aa122-168) (exact	Marker	MALT-Lymphoma Marker		
	Specificity/Sensitivity	number of apoptotic regulatory molecules. It was identified through its direct involvement in t(1;14) of mucosa-associated lymphoid tissue (MALT) lymphoma. Expression of BCL10 was shown to induce NF B activation in a NIK-dependent pathway. This monoclonal antibody labels subpopulations of normal B and T cells and is a useful tool for the sub-classification of lymphomas. In MALT lymphomas with the t(1;14) translocation, while 55% of MALT lymphomas lacking this translocation exhibited the same labeling pattern, although at a much lower level.		
	Immunogen			



	9
Notes	Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.
Product Application Details	

Product Application Details	
Applications	Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin
Recommended Dilutions	Flow Cytometry, Immunohistochemistry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry-Paraffin
Application Notes	Optimal dilution of this antibody should be experimentally determined.



## Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

### **Products Related to NBP3-08803AF488**

NBP2-24982 Rabbit IgG Isotype Control [Alexa Fluor® 488]

NBP1-99064-10uq Recombinant Human Bcl-10 His Protein

210-TA-005 TNF-alpha [Unconjugated]
NBP2-07935 Bcl-10 Overexpression Lysate

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-08803AF488

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

