Product Datasheet

Bcl-10 Antibody (BL10/2988R) [Alexa Fluor® 594] NBP3-08803AF594

Unit Size: 100 ul

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-08803AF594

Updated 10/26/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP3-08803AF594



NBP3-08803AF594

Bcl-10 Antibody (BL10/2988R) [Alexa Fluor® 594]

Unit Size 100 ul Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonality Monoclonal Clone BL10/2988R Preservative 0.05% Sodium Azide Isotype IgG Conjugate Alexa Fluor 594 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Rabbit Gene ID 8915 Gene Symbol BCL10 Species Human Marker MALT-Lymphoma Marker Specificity/Sensitivity BCL10, with an N-terminal caspase recruitment domain (CARD), is found in a number of apoptotic regulatory molecules. It was identified through its direct involvement in t(1;14) of mucosa-associated lymphoid tissue (MALT) lymphoma. Expression of BCL10 was shown to induce NFB a activation in a NIK-dependent pathway. This monoclonal antibody labels subpopulations of normal B and T cells and is a useful tool for the sub-classification of lymphomas. In MALT lymphomas with the t(1;14) translocation, while 55% of MALT lymphomas lacking this translocation exhibited the same labeling pattern, although at a much lower level. Immunogen	DOI 10 / WILLDOUG (DE 10/20001) [/ WOXA / WOXA /		
Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonality Monoclonal Clone BL10/2988R Preservative 0.05% Sodium Azide Isotype IgG Conjugate Alexa Fluor 594 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Rabbit Gene ID 8915 Gene Symbol BCL10 Species Human Marker MALT-Lymphoma Marker Specificity/Sensitivity BCL10, with an N-terminal caspase recruitment domain (CARD), is found in a number of apoptotic regulatory molecules. It was identified through its direct involvement in 1(1;14) of mucosa-associated lymphoid tissue (MALT) lymphoma. Expression of BCL10 was shown to induce NF B activation in a NIK-dependent pathway. This monoclonal antibody labels subpopulations of normal B and T cells and is a useful tool for the sub-classification of lymphomas. In MALT lymphomas with the t(1;14) translocation, while 55% of MALT lymphomas lacking this translocation exhibited the same labeling pattern, although at a much lower level. Immunogen Recombinant human Bcl-10 protein fragment (around aa122-168) (exact	Product Information		
Storage Store at 4C in the dark. Clonality Monoclonal Clone BL10/2988R Preservative 0.05% Sodium Azide Isotype IgG Conjugate Alexa Fluor 594 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Rabbit Gene ID 8915 Gene Symbol BCL10 Species Human Marker MALT-Lymphoma Marker Specificity/Sensitivity BCL10, with an N-terminal caspase recruitment domain (CARD), is found in a number of apoptotic regulatory molecules. It was identified through its direct involvement in 1(1;14) of mucosa-associated lymphoid tissue (MALT) lymphoma. Expression of BCL10 was shown to induce NF B activation in a NIK-dependent pathway. This monoclonal antibody labels subpopulations of normal B and T cells and is a useful tool for the sub-classification of lymphomas. In MALT lymphomas with the 1(1;14) translocation, while 55% of MALT lymphomas lacking this translocation exhibited the same labeling pattern, although at a much lower level. Immunogen Recombinant human Bcl-10 protein fragment (around aa122-168) (exact	Unit Size	100 ul	
Clone BL10/2988R Preservative 0.05% Sodium Azide Isotype IgG Conjugate Alexa Fluor 594 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Rabbit Gene ID 8915 Gene Symbol BCL10 Species Human Marker MALT-Lymphoma Marker Specificity/Sensitivity BCL10, with an N-terminal caspase recruitment domain (CARD), is found in a number of apoptotic regulatory molecules. It was identified through its direct involvement in t(1;14) of mucosa-associated lymphoid tissue (MALT) lymphoma. Expression of BCL10 was shown to induce NF B activation in a NIK-dependent pathway. This monoclonal antibody labels subpopulations of normal B and T cells and is a useful tool for the sub-classification of lymphomas. In MALT lymphomas with the t(1;14) translocation, while 55% of MALT lymphomas lacking this translocation exhibited the same labeling pattern, although at a much lower level. Immunogen Recombinant human Bcl-10 protein fragment (around aa122-168) (exact	Concentration	·	
Clone BL10/2988R Preservative 0.05% Sodium Azide Isotype IgG Conjugate Alexa Fluor 594 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Rabbit Rabbit Gene ID 8915 Gene Symbol BCL10 Species Human Marker MALT-Lymphoma Marker Specificity/Sensitivity BCL10, with an N-terminal caspase recruitment domain (CARD), is found in a number of apoptotic regulatory molecules. It was identified through its direct involvement in t(1;14) of mucosa-associated lymphoid tissue (MALT) lymphoma. Expression of BCL10 was shown to induce NF B activation in a NIK-dependent pathway. This monoclonal antibody labels subpopulations of normal B and T cells and is a useful tool for the sub-classification of lymphomas. In MALT lymphomas with the t(1;14) translocation, while 55% of MALT lymphomas lacking this translocation exhibited the same labeling pattern, although at a much lower level. Immunogen Recombinant human Bcl-10 protein fragment (around aa122-168) (exact	Storage	Store at 4C in the dark.	
Preservative 0.05% Sodium Azide Isotype IgG Conjugate Alexa Fluor 594 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Rabbit Rabbit Gene ID 8915 Gene Symbol BCL10 Species Human Marker MALT-Lymphoma Marker Specificity/Sensitivity BCL10, with an N-terminal caspase recruitment domain (CARD), is found in a number of apoptotic regulatory molecules. It was identified through its direct involvement in t(1;14) of mucosa-associated lymphoid tissue (MALT) lymphoma. Expression of BCL10 was shown to induce NF B activation in a NIK-dependent pathway. This monoclonal antibody labels subpopulations of normal B and T cells and is a useful tool for the sub-classification of lymphomas. In MALT lymphomas with the t(1;14) translocation, while 55% of MALT lymphomas lacking this translocation exhibited the same labeling pattern, although at a much lower level. Immunogen Recombinant human Bcl-10 protein fragment (around aa122-168) (exact	Clonality	Monoclonal	
Isotype IgG Conjugate Alexa Fluor 594 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Rabbit Gene ID 8915 Gene Symbol BCL10 Species Human Marker MALT-Lymphoma Marker Specificity/Sensitivity BCL10, with an N-terminal caspase recruitment domain (CARD), is found in a number of apoptotic regulatory molecules. It was identified through its direct involvement in t(1;14) of mucosa-associated lymphoid tissue (MALT) lymphoma. Expression of BCL10 was shown to induce NF B activation in a NIK-dependent pathway. This monoclonal antibody labels subpopulations of normal B and T cells and is a useful tool for the sub-classification of lymphomas. In MALT lymphomas with the t(1;14) translocation, while 55% of MALT lymphomas lacking this translocation exhibited the same labeling pattern, although at a much lower level. Immunogen Recombinant human Bcl-10 protein fragment (around aa122-168) (exact	Clone	BL10/2988R	
Conjugate Alexa Fluor 594 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Rabbit Gene ID 8915 Gene Symbol BCL10 Species Human Marker MALT-Lymphoma Marker Specificity/Sensitivity BCL10, with an N-terminal caspase recruitment domain (CARD), is found in a number of apoptotic regulatory molecules. It was identified through its direct involvement in t(1;14) of mucosa-associated lymphoid tissue (MALT) lymphoma. Expression of BCL10 was shown to induce NF B activation in a NIK-dependent pathway. This monoclonal antibody labels subpopulations of normal B and T cells and is a useful tool for the sub-classification of lymphomas. In MALT lymphomas with the t(1;14) translocation, while 55% of MALT lymphomas lacking this translocation exhibited the same labeling pattern, although at a much lower level. Immunogen Recombinant human Bcl-10 protein fragment (around aa122-168) (exact	Preservative	0.05% Sodium Azide	
Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Rabbit Gene ID 8915 Gene Symbol BCL10 Species Human Marker MALT-Lymphoma Marker Specificity/Sensitivity BCL10, with an N-terminal caspase recruitment domain (CARD), is found in a number of apoptotic regulatory molecules. It was identified through its direct involvement in t(1;14) of mucosa-associated lymphoid tissue (MALT) lymphoma. Expression of BCL10 was shown to induce NF B activation in a NIK-dependent pathway. This monoclonal antibody labels subpopulations of normal B and T cells and is a useful tool for the sub-classification of lymphomas. In MALT lymphomas with the t(1;14) translocation, while 55% of MALT lymphomas lacking this translocation exhibited the same labeling pattern, although at a much lower level. Immunogen Recombinant human Bcl-10 protein fragment (around aa122-168) (exact	Isotype	IgG	
Product Description	Conjugate	Alexa Fluor 594	
Product Description Host Rabbit Gene ID 8915 Gene Symbol BCL10 Species Human Marker MALT-Lymphoma Marker Specificity/Sensitivity BCL10, with an N-terminal caspase recruitment domain (CARD), is found in a number of apoptotic regulatory molecules. It was identified through its direct involvement in t(1;14) of mucosa-associated lymphoid tissue (MALT) lymphoma. Expression of BCL10 was shown to induce NF B activation in a NIK-dependent pathway. This monoclonal antibody labels subpopulations of normal B and T cells and is a useful tool for the sub-classification of lymphomas. In MALT lymphomas with the t(1;14) translocation, while 55% of MALT lymphomas lacking this translocation exhibited the same labeling pattern, although at a much lower level. Immunogen Recombinant human Bcl-10 protein fragment (around aa122-168) (exact	Purity	Protein A or G purified	
Rabbit R	Buffer	50mM Sodium Borate	
Gene ID 8915 Gene Symbol BCL10 Species Human Marker MALT-Lymphoma Marker Specificity/Sensitivity BCL10, with an N-terminal caspase recruitment domain (CARD), is found in a number of apoptotic regulatory molecules. It was identified through its direct involvement in t(1;14) of mucosa-associated lymphoid tissue (MALT) lymphoma. Expression of BCL10 was shown to induce NF B activation in a NIK-dependent pathway. This monoclonal antibody labels subpopulations of normal B and T cells and is a useful tool for the sub-classification of lymphomas. In MALT lymphomas with the t(1;14) translocation, while 55% of MALT lymphomas lacking this translocation exhibited the same labeling pattern, although at a much lower level. Immunogen Recombinant human Bcl-10 protein fragment (around aa122-168) (exact	Product Description		
Gene Symbol BCL10 Species Human Marker MALT-Lymphoma Marker Specificity/Sensitivity BCL10, with an N-terminal caspase recruitment domain (CARD), is found in a number of apoptotic regulatory molecules. It was identified through its direct involvement in t(1;14) of mucosa-associated lymphoid tissue (MALT) lymphoma. Expression of BCL10 was shown to induce NF B activation in a NIK-dependent pathway. This monoclonal antibody labels subpopulations of normal B and T cells and is a useful tool for the sub-classification of lymphomas. In MALT lymphomas with the t(1;14) translocation, while 55% of MALT lymphomas lacking this translocation exhibited the same labeling pattern, although at a much lower level. Immunogen Recombinant human Bcl-10 protein fragment (around aa122-168) (exact	Host	Rabbit	
Species Marker MALT-Lymphoma Marker Specificity/Sensitivity BCL10, with an N-terminal caspase recruitment domain (CARD), is found in a number of apoptotic regulatory molecules. It was identified through its direct involvement in t(1;14) of mucosa-associated lymphoid tissue (MALT) lymphoma. Expression of BCL10 was shown to induce NF B activation in a NIK-dependent pathway. This monoclonal antibody labels subpopulations of normal B and T cells and is a useful tool for the sub-classification of lymphomas. In MALT lymphomas with the t(1;14) translocation, while 55% of MALT lymphomas lacking this translocation exhibited the same labeling pattern, although at a much lower level. Immunogen Recombinant human Bcl-10 protein fragment (around aa122-168) (exact	Gene ID	8915	
MALT-Lymphoma Marker Specificity/Sensitivity BCL10, with an N-terminal caspase recruitment domain (CARD), is found in a number of apoptotic regulatory molecules. It was identified through its direct involvement in t(1;14) of mucosa-associated lymphoid tissue (MALT) lymphoma. Expression of BCL10 was shown to induce NF B activation in a NIK-dependent pathway. This monoclonal antibody labels subpopulations of normal B and T cells and is a useful tool for the sub-classification of lymphomas. In MALT lymphomas with the t(1;14) translocation, while 55% of MALT lymphomas lacking this translocation exhibited the same labeling pattern, although at a much lower level. Immunogen Recombinant human Bcl-10 protein fragment (around aa122-168) (exact	Gene Symbol	BCL10	
Specificity/Sensitivity BCL10, with an N-terminal caspase recruitment domain (CARD), is found in a number of apoptotic regulatory molecules. It was identified through its direct involvement in t(1;14) of mucosa-associated lymphoid tissue (MALT) lymphoma. Expression of BCL10 was shown to induce NF B activation in a NIK-dependent pathway. This monoclonal antibody labels subpopulations of normal B and T cells and is a useful tool for the sub-classification of lymphomas. In MALT lymphomas with the t(1;14) translocation, while 55% of MALT lymphomas lacking this translocation exhibited the same labeling pattern, although at a much lower level. Immunogen Recombinant human Bcl-10 protein fragment (around aa122-168) (exact	Species	Human	
number of apoptotic regulatory molecules. It was identified through its direct involvement in t(1;14) of mucosa-associated lymphoid tissue (MALT) lymphoma. Expression of BCL10 was shown to induce NF B activation in a NIK-dependent pathway. This monoclonal antibody labels subpopulations of normal B and T cells and is a useful tool for the sub-classification of lymphomas. In MALT lymphomas with the t(1;14) translocation, while 55% of MALT lymphomas lacking this translocation exhibited the same labeling pattern, although at a much lower level. Immunogen Recombinant human Bcl-10 protein fragment (around aa122-168) (exact	Marker	MALT-Lymphoma Marker	
		number of apoptotic regulatory molecules. It was identified through its direct involvement in t(1;14) of mucosa-associated lymphoid tissue (MALT) lymphoma. Expression of BCL10 was shown to induce NF B activation in a NIK-dependent pathway. This monoclonal antibody labels subpopulations of normal B and T cells and is a useful tool for the sub-classification of lymphomas. In MALT lymphomas with the t(1;14) translocation, while 55% of MALT lymphomas lacking this translocation exhibited the same labeling pattern, although at a much lower level.	
	Immunogen		



	9
Notes	Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.
Product Application Details	

Product Application Details	
Applications	Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin
Recommended Dilutions	Flow Cytometry, Immunohistochemistry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry-Paraffin
Application Notes	Optimal dilution of this antibody should be experimentally determined.



Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP3-08803AF594

IC1051T Rabbit IgG Isotype Control (60024B) [Alexa Fluor® 594]

NBP1-99064-10ug Recombinant Human Bcl-10 His Protein

210-TA-005 TNF-alpha [Unconjugated]
NBP2-07935 Bcl-10 Overexpression Lysate

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-08803AF594

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

