# **Product Datasheet**

## Alkaline Phosphatase/ALPP Antibody (ALPP/2899R) [Alexa Fluor® 405] NBP3-08817AF405

Unit Size: 100 ul

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-08817AF405

Updated 10/26/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP3-08817AF405



#### NBP3-08817AF405

Alkaline Phosphatase/ALPP Antibody (ALPP/2899R) [Alexa Fluor® 405]

diagnosis of gestational trophoblastic disease.ImmunogenRecombinant full-length human Alkaline Phosphatase/ALPP protein (Uniprot:	•	
ConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneALPP/2899RPreservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 405PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionKabbitGene ID250Gene SymbolALPPSpeciesHumanMarkerGerm Cell Tumor MarkerSpecificity/SensitivityReacts with a 70kDa membrane-bound isozyme (Regan and Nagao type) of Placental Alkaline Phosphatase (PLAP) occurring in the placenta during the 3rd trimester of gestation. It is lighly specific for PLAP and shows no cross-reaction with other isozymes of alkaline phosphatase. Anti-PLAP leasts with germ cell tumors and can discriminate between these and other neoplasms. Somatic neoplasms e.g. breast, gastrointestinal, prostatic, and urinary cancers may also immunoreact with antibedies to PLAP. Anti-PLAP positivity in conjunction with anti-keratin negativity favors seminoma over carcinoma. Germ cell tumors are usually anti-keratin negativity favors seminoma over carcinoma. Germ cell unter diagnosis of gestational trophoblastic disease.ImmunogenRecombinant full-length human Alkaline Phosphatase/ALPP protein (Uniprot:	Product Information	
services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneALPP/2899RPreservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 405PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionKabbitHostRabbitGene ID250Gene SymbolALPPSpeciesHumanMarkerGerm Cell Tumor MarkerSpecificity/SensitivityPlacental Akaline Phosphatase (PLAP) occurring in the placenta during the 3rd trimester of gestation. It is highly specific for PLAP and shows no cross-reaction with other isozymes of alkaline phosphatase. Anti-PLAP reacts with germ cell tumors and can discriminate between these and other neoplasms. Somatic neoplasms e.g. breast, gastrointestinal, prostatic, and urinary cancers may also immunoreact with antibodies to PLAP. Anti-PLAP has been useful in the diagnosis of gestational trophoblastic disease.ImmunogenRecombinant full-length human Alkaline Phosphatase./ALPP protein (Uniprot:	Unit Size	100 ul
ClonalityMonoclonalCloneALPP/2899RPreservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 405PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionEabbitHostRabbitGene ID250Gene SymbolALPPSpeciesHumanMarkerGerm Cell Tumor MarkerSpecificity/SensitivityReacts with a 70kDa membrane-bound isozyme (Regan and Nagao type) of Placental Alkaline Phosphatase (PLAP) occurring in the placenta during the 3rd trimester of gestation. It is highly specific for PLAP and shows no cross-reaction with other isozymes of alkaline phosphatase. Anti-PLAP reacts with germ cell tumors and can discriminate between these and other neoplasms. Somatic neoplasms e.g. breast, gastrointestinal, prostatic, and urinary cancers may also immunoreact with antibodies to PLAP. Anti-PLAP positivity in conjunction with anti-Keratin negativity favors seminoma over carcinoma. Germ cell tumors are usually anti-keratin positive, but they regularly fail to stain with anti-EMA, whereas most carcinomas stain with anti-EMA, Anti-PLAP has been useful in the diagnosis of gestational trophoblastic disease.ImmunogenRecombinant full-length human Alkaline Phosphatase/ALPP protein (Uniprot:	Concentration	
CloneALPP/2899RPreservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 405PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionExampleHostRabbitGene ID250Gene SymbolALPPSpeciesHumanMarkerGerm Cell Tumor MarkerSpecificity/SensitivityReacts with a 70kDa membrane-bound isozyme (Regan and Nagao type) of Placental Alkaline Phosphatase (PLAP) occurring in the placenta during the 3rd trimester of gestation. It is highly specific for PLAP and shows no cross-reaction with other isozymes of alkaline phosphatase. Anti-PLAP neacts with germ cell tumors and can discriminate between these and other neoplasms. Somatic neoplasms e.g. breast, gastrointestinal, prostatic, and urinary cancers may also immunoreact with antibodies to PLAP. Anti-PLAP positivity in conjunction with anti-Keratin negativity favors seminoma over carcinoma. Germ cell tumors are usually anti-keratin positive, but they regularly fail to stain with anti-EMA, whereas most carcinomas stain with anti-EMA. Anti-PLAP protein (Uniprot:ImmunogenRecombinant full-length human Alkaline Phosphatase/ALPP protein (Uniprot:	Storage	Store at 4C in the dark.
Preservative 0.05% Sodium Azide   Isotype IgG   Conjugate Alexa Fluor 405   Purity Protein A or G purified   Buffer 50mM Sodium Borate   Product Description Host   Host Rabbit   Gene ID 250   Gene Symbol ALPP   Species Human   Marker Germ Cell Tumor Marker   Specificity/Sensitivity Reacts with a 70kDa membrane-bound isozyme (Regan and Nagao type) of Placental Alkaline Phosphatase (PLAP) occurring in the placenta during the 3rd trimester of gestation. It is highly specific for PLAP neacts with germ cell tumors and can discriminate between these and other neoplasms. Somatic neoplasms e.g. breast, gastrointestinal, prostatic, and urinary cancers may also immunoreact with antib-elise to PLAP. Anti-PLAP positivity in conjunction with anti-keratin negativity favors seminoma over carcinoma. Germ cell tumors are usually anti-keratin positive, but they regularly fail to stain with anti-EMA, whereas most carcinomas stain with anti-PLAP has been useful in the diagnosis of gestational trophoblastic disease.   Immunogen Recombinant full-length human Alkaline Phosphatase/ALPP protein (Uniprot:	Clonality	Monoclonal
IsotypeIgGConjugateAlexa Fluor 405PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID250Gene SymbolALPPSpeciesHumanMarkerGerm Cell Tumor MarkerSpecificity/SensitivityReacts with a 70kDa membrane-bound isozyme (Regan and Nagao type) of Placental Alkaline Phosphatase (PLAP) occurring in the placenta during the 3rd trimester of gestation. It is highly specific for PLAP and shows no cross-reaction with other isozymes of alkaline phosphatase. Anti-PLAP reacts with germ cell tumors and can discriminate between these and other neoplasms. Somatic neoplasms e.g. breast, gastrointestinal, prostitiv, in conjunction with anti-keratin negativity favors seminoma over carcinoma. Germ cell tumors are usually anti-keratin positive, but they regularly fail to stain with anti-EMA, whereas most carcinomas stain with anti-PLAP protein (Uniprot:ImmunogenRecombinant full-length human Alkaline Phosphatase/ALPP protein (Uniprot:	Clone	ALPP/2899R
ConjugateAlexa Fluor 405PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID250Gene SymbolALPPSpeciesHumanMarkerGerm Cell Tumor MarkerSpecificity/SensitivityReacts with a 70kDa membrane-bound isozyme (Regan and Nagao type) of Placental Alkaline Phosphatase (PLAP) occurring in the placenta during the 3rd trimester of gestation. It is highly specific for PLAP and shows no cross-reaction with other isozymes of alkaline phosphatase. Anti-PLAP reacts with germ cell tumors and can discriminate between these and other neoplasms. Somatic neoplasms e.g. breast, gastrointestinal, prostatic, and urinary cancers may also immunoreact with anti-bodies to PLAP. Anti-PLAP positivity in conjunction with anti-keratin negativity favors seminoma over carcinoma. Germ cell tumors are usually anti-keratin positive, but they regularly fail to stain with anti-EMA, whereas most carcinomas stain with anti-EMA. Anti-PLAP has been useful in the diagnosis of gestational trophoblastic disease.ImmunogenRecombinant full-length human Alkaline Phosphatase/ALPP protein (Uniprot:	Preservative	0.05% Sodium Azide
Purity Protein A or G purified   Buffer 50mM Sodium Borate   Product Description    Host Rabbit   Gene ID 250   Gene Symbol ALPP   Species Human   Marker Germ Cell Tumor Marker   Specificity/Sensitivity Reacts with a 70kDa membrane-bound isozyme (Regan and Nagao type) of Placental Alkaline Phosphatase (PLAP) occurring in the placenta during the 3rd trimester of gestation. It is highly specific for PLAP and shows no cross-reaction with other isozymes of alkaline phosphatase. Anti-PLAP reacts with germ cell tumors and can discriminate between these and other neoplasms. Somatic neoplasms e.g. breast, gastrointestinal, prostatic, and urinary cancers may also immunoreact with antibodies to PLAP. Anti-PLAP positivity in conjunction with anti-keratin negativity favors seminoma over carcinoma. Germ cell tumors are usually anti-keratin positive, but they regularly fail to stain with anti-EMA, whereas most carcinomas stain with anti-EMA. Anti-PLAP has been useful in the diagnosis of gestational trophoblastic disease.   Immunogen Recombinant full-length human Alkaline Phosphatase/ALPP protein (Uniprot:	Isotype	IgG
Buffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID250Gene SymbolALPPSpeciesHumanMarkerGerm Cell Tumor MarkerSpecificity/SensitivityReacts with a 70kDa membrane-bound isozyme (Regan and Nagao type) of Placental Alkaline Phosphatase (PLAP) occurring in the placenta during the 3rd trimester of gestation. It is highly specific for PLAP and shows no cross-reaction with other isozymes of alkaline phosphatase. Anti-PLAP neacts with germ cell tumors and can discriminate between these and other neoplasms. Somatic neoplasms e.g. breast, gastrointestinal, prostatic, and urinary cancers may also immunoreact with antibodies to PLAP. Anti-PLAP positivity in conjunction with anti-keratin negativity favors seminoma over carcinoma. Germ cell tumors are usually anti-keratin positive, but they regularly fail to stain with anti-EMA, whereas most carcinomas stain with anti-EMA. Anti-PLAP has been useful in the diagnosis of gestational trophoblastic disease.ImmunogenRecombinant full-length human Alkaline Phosphatase/ALPP protein (Uniprot:	Conjugate	Alexa Fluor 405
Product DescriptionHostRabbitGene ID250Gene SymbolALPPSpeciesHumanMarkerGerm Cell Tumor MarkerSpecificity/SensitivityReacts with a 70kDa membrane-bound isozyme (Regan and Nagao type) of Placental Alkaline Phosphatase (PLAP) occurring in the placenta during the 3rd trimester of gestation. It is highly specific for PLAP and shows no cross-reaction with other isozymes of alkaline phosphatase. Anti-PLAP reacts with germ cell tumors and can discriminate between these and other neoplasms. Somatic neoplasms e.g. breast, gastrointestinal, prostatic, and urinary cancers may also immunoreact with antibodies to PLAP. Anti-PLAP positivity in conjunction with anti-keratin negativity favors seminoma over carcinoma. Germ cell tumors are usually anti-keratin positive, but they regularly fail to stain with anti-EMA, whereas most carcinomas stain with anti-EMA, Anti-PLAP has been useful in the diagnosis of gestational trophoblastic disease.ImmunogenRecombinant full-length human Alkaline Phosphatase/ALPP protein (Uniprot:	Purity	Protein A or G purified
HostRabbitGene ID250Gene SymbolALPPSpeciesHumanMarkerGerm Cell Tumor MarkerSpecificity/SensitivityReacts with a 70kDa membrane-bound isozyme (Regan and Nagao type) of Placental Alkaline Phosphatase (PLAP) occurring in the placenta during the 3rd trimester of gestation. It is highly specific for PLAP and shows no cross-reaction with other isozymes of alkaline phosphatase. Anti-PLAP reacts with germ cell tumors and can discriminate between these and other neoplasms. Somatic neoplasms e.g. breast, gastrointestinal, prostatic, and urinary cancers may also immunoreact with anti-bedies to PLAP. Anti-PLAP positivity in conjunction with anti-keratin negativity favors seminoma over carcinoma. Germ cell tumors are usually anti-keratin postive, but they regularly fail to stain with anti-EMA, whereas most carcinomas stain with anti-FLAP has been useful in the diagnosis of gestational trophoblastic disease.ImmunogenRecombinant full-length human Alkaline Phosphatase/ALPP protein (Uniprot:	Buffer	50mM Sodium Borate
Gene ID250Gene SymbolALPPSpeciesHumanMarkerGerm Cell Tumor MarkerSpecificity/SensitivityReacts with a 70kDa membrane-bound isozyme (Regan and Nagao type) of Placental Alkaline Phosphatase (PLAP) occurring in the placenta during the 3rd trimester of gestation. It is highly specific for PLAP and shows no cross-reaction with other isozymes of alkaline phosphatase. Anti-PLAP reacts with germ cell tumors and can discriminate between these and other neoplasms. Somatic neoplasms e.g. breast, gastrointestinal, prostatic, and urinary cancers may also immunoreact with anti-bodies to PLAP. Anti-PLAP positivity in conjunction with anti-keratin negativity favors seminoma over carcinoma. Germ cell tumors are usually anti-keratin positive, but they regularly fail to stain with anti-EMA, whereas most carcinomas stain with anti-EMA. Anti-PLAP has been useful in the diagnosis of gestational trophoblastic disease.ImmunogenRecombinant full-length human Alkaline Phosphatase/ALPP protein (Uniprot:	Product Description	
Gene SymbolALPPSpeciesHumanMarkerGerm Cell Tumor MarkerSpecificity/SensitivityReacts with a 70kDa membrane-bound isozyme (Regan and Nagao type) of Placental Alkaline Phosphatase (PLAP) occurring in the placenta during the 3rd trimester of gestation. It is highly specific for PLAP and shows no cross-reaction with other isozymes of alkaline phosphatase. Anti-PLAP reacts with germ cell tumors and can discriminate between these and other neoplasms. Somatic neoplasms e.g. breast, gastrointestinal, prostatic, and urinary cancers may also immunoreact with antibodies to PLAP. Anti-PLAP positivity in conjunction with anti-keratin negativity favors seminoma over carcinoma. Germ cell tumors are usually anti-keratin positive, but they regularly fail to stain with anti-EMA, whereas most carcinomas stain with anti-EMA. Anti-PLAP has been useful in the diagnosis of gestational trophoblastic disease.ImmunogenRecombinant full-length human Alkaline Phosphatase/ALPP protein (Uniprot:	Host	Rabbit
SpeciesHumanMarkerGerm Cell Tumor MarkerSpecificity/SensitivityReacts with a 70kDa membrane-bound isozyme (Regan and Nagao type) of Placental Alkaline Phosphatase (PLAP) occurring in the placenta during the 3rd trimester of gestation. It is highly specific for PLAP and shows no cross-reaction with other isozymes of alkaline phosphatase. Anti-PLAP reacts with germ cell tumors and can discriminate between these and other neoplasms. Somatic neoplasms e.g. breast, gastrointestinal, prostatic, and urinary cancers may also immunoreact with antibodies to PLAP. Anti-PLAP positivity in conjunction with anti-keratin negativity favors seminoma over carcinoma. Germ cell tumors are usually anti-keratin positive, but they regularly fail to stain with anti-EMA, whereas most carcinomas stain with anti-EMA. Anti-PLAP has been useful in the diagnosis of gestational trophoblastic disease.ImmunogenRecombinant full-length human Alkaline Phosphatase/ALPP protein (Uniprot:	Gene ID	250
MarkerGerm Cell Tumor MarkerSpecificity/SensitivityReacts with a 70kDa membrane-bound isozyme (Regan and Nagao type) of Placental Alkaline Phosphatase (PLAP) occurring in the placenta during the 3rd trimester of gestation. It is highly specific for PLAP and shows no cross-reaction with other isozymes of alkaline phosphatase. Anti-PLAP reacts with germ cell tumors and can discriminate between these and other neoplasms. Somatic neoplasms e.g. breast, gastrointestinal, prostatic, and urinary cancers may also immunoreact with antibodies to PLAP. Anti-PLAP positivity in conjunction with anti-keratin negativity favors seminoma over carcinoma. Germ cell tumors are usually anti-keratin positive, but they regularly fail to stain with anti-EMA, whereas most carcinomas stain with anti-EMA. Anti-PLAP has been useful in the diagnosis of gestational trophoblastic disease.ImmunogenRecombinant full-length human Alkaline Phosphatase/ALPP protein (Uniprot:	Gene Symbol	ALPP
Specificity/SensitivityReacts with a 70kDa membrane-bound isozyme (Regan and Nagao type) of Placental Alkaline Phosphatase (PLAP) occurring in the placenta during the 3rd trimester of gestation. It is highly specific for PLAP and shows no cross-reaction with other isozymes of alkaline phosphatase. Anti-PLAP reacts with germ cell tumors and can discriminate between these and other neoplasms. Somatic neoplasms e.g. breast, gastrointestinal, prostatic, and urinary cancers may also immunoreact with antibodies to PLAP. Anti-PLAP positivity in conjunction with anti-keratin negativity favors seminoma over carcinoma. Germ cell tumors are usually anti-keratin positive, but they regularly fail to stain with anti-EMA, whereas most carcinomas stain with anti-EMA. Anti-PLAP has been useful in the diagnosis of gestational trophoblastic disease.ImmunogenRecombinant full-length human Alkaline Phosphatase/ALPP protein (Uniprot:	Species	Human
Placental Alkaline Phosphatase (PLAP) occurring in the placenta during the 3rd trimester of gestation. It is highly specific for PLAP and shows no cross-reaction with other isozymes of alkaline phosphatase. Anti-PLAP reacts with germ cell tumors and can discriminate between these and other neoplasms. Somatic neoplasms e.g. breast, gastrointestinal, prostatic, and urinary cancers may also immunoreact with antibodies to PLAP. Anti-PLAP positivity in conjunction with anti-keratin negativity favors seminoma over carcinoma. Germ cell tumors are usually anti-keratin positive, but they regularly fail to stain with anti-EMA, whereas most carcinomas stain with anti-EMA. Anti-PLAP has been useful in the diagnosis of gestational trophoblastic disease.ImmunogenRecombinant full-length human Alkaline Phosphatase/ALPP protein (Uniprot:	Marker	Germ Cell Tumor Marker
	Specificity/Sensitivity	Placental Alkaline Phosphatase (PLAP) occurring in the placenta during the 3rd trimester of gestation. It is highly specific for PLAP and shows no cross-reaction with other isozymes of alkaline phosphatase. Anti-PLAP reacts with germ cell tumors and can discriminate between these and other neoplasms. Somatic neoplasms e.g. breast, gastrointestinal, prostatic, and urinary cancers may also immunoreact with antibodies to PLAP. Anti-PLAP positivity in conjunction with anti-keratin negativity favors seminoma over carcinoma. Germ cell tumors are usually anti-keratin positive, but they regularly fail to stain with anti-EMA, whereas most carcinomas stain with anti-EMA. Anti-PLAP has been useful in the
	Immunogen	Recombinant full-length human Alkaline Phosphatase/ALPP protein (Uniprot: P05187)

www.novusbio.com



	Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.
Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin
<b>Recommended Dilutions</b>	Western Blot, Immunohistochemistry-Paraffin
Application Notes	Optimal dilution of this antibody should be experimentally determined.

Notes





### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

#### Products Related to NBP3-08817AF405

NBP2-24891AF405	Rabbit IgG Isotype Control [Alexa Fluor® 405]
NBP2-34056PEP	Alkaline Phosphatase/ALPP Recombinant Protein Antigen
210-TA-005	TNF-alpha [Unconjugated]
NBL1-07488	Alkaline Phosphatase/ALPP Overexpression Lysate

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-08817AF405

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

