

Product Datasheet

Fc epsilon RI alpha Antibody (Mar-1) - Azide and BSA Free NBP3-09039-0.2mg

Unit Size: 0.2 mg

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP3-09039

Updated 11/7/2023 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP3-09039



NBP3-09039-0.2mg

Fc epsilon RI alpha Antibody (Mar-1) - Azide and BSA Free

Product Information	
Unit Size	0.2 mg
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	Mar-1
Preservative	0.02% Proclin 300
Isotype	IgG Lambda
Purity	Protein A purified
Buffer	PBS
Product Description	
Host	Armenian Hamster
Gene Symbol	FCER1A
Species	Mouse
Specificity/Sensitivity	This antibody specifically targets the non-signalling alpha chain of the FcεR1 complex.
Immunogen	FcεR1 alpha
Product Application Details	
Applications	Flow Cytometry, Immunohistochemistry, Immunoprecipitation
Recommended Dilutions	Flow Cytometry, Immunohistochemistry, Immunoprecipitation
Application Notes	This antibody has been used in flow cytometric analysis of FcεR1 alpha cell surface expression by basophils (Min et al, 2004; Chen et al, 2009), and immunohistochemical analysis of epidermal sheets (Holzmann et al, 2004). It has also been used in immunoprecipitation analysis of cell lysates from lung cDCs and MC/9 mast cells (Grayson et al, 2007). In vitro incubation of basophils with this antibody blocks subsequent MAR-1 staining, but does not activate basophil IL-4 production (Sokol et al, 2008). Intravenous treatment of mice with this antibody results in basophil depletion in peripheral blood, spleen, bone marrow and liver, but not skin or intraperitoneal mast cell depletion (Sokol et al, 2008). Following treatment with this antibody, basophil-depleted mice display impaired Th2 differentiation upon papain immunisation (Sokol et al, 2008). However, MAR-1 treated mice respond normally to high-dose intravenous Ig in both the K/BxN serum transfer arthritis and collagen-induced arthritis models, despite basophil depletion (Campbell et al, 2014).





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP3-09039-0.2mg

NB120-5738	Goat anti-Armenian Hamster IgG (H+L) Secondary Antibody
6678-FC-050	Fc epsilon RI alpha [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-09039

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

