Product Datasheet

MUC5AC Antibody (9-13M1) [DyLight 650] NBP3-11484C

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-11484C

Updated 10/26/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP3-11484C



NBP3-11484C

MUC5AC Antibody (9-13M1) [DyLight 650]

Product Information	ical
Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonality Monoclonal Clone 9-13M1 Preservative 10.05% Sodium Azide Isotype IgG1 Kappa Conjugate DyLight 650 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 4586 Gene Symbol MUC5AC Species Human Specificity/Sensitivity This monoclonal antibody recognizes the peptide core of gastric mucin M1 (recently identified as Mucin 5AC). Its epitope is located in the N-terminal cysteine rich part of the peptide core of MUC5AC, which is heavily glycosylater. Its epitope is destroyed by beta-mercaptoethalo but not by periodate treatmen monoclonal antibody 2-11M1 reacts with the protein backbone exclusively; it or reacts with fully deglycosylated. This can be achieved with standard periodate oxidation method. The success of the deglycosylation, the preparation should no longer be stainable with PAS reagent. Only then 2-11M1 will react should MUC5AC be present. This mucin is present in primary ovariar mucinous cancer but usually absent in colorectal adenocarcinoma, thus showil an expression pattern opposite to MUC2. Together with a panel of antibodies,	ical
Storage Store at 4C in the dark. Clonality Monoclonal Clone 9-13M1 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate DyLight 650 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 4586 Gene Symbol MUC5AC Species Human Specificity/Sensitivity This monoclonal antibody recognizes the peptide core of gastric mucin M1 (recently identified as Mucin 5AC). Its epitope is located in the N-terminal cysteine rich part of the peptide core of MUC5AC, which is heavily glycosylated Its epitope is destroyed by beta-mercaptoethanol but not by periodate treatmen monoclonal antibody 2-11M1 reacts with the protein backbone exclusively; it o reacts with fully deglycosylated. This can be achieved with standard periodate oxidation method. The success of the deglycosylation can be checke with routine PAS (Periodic Acid Shiff) staning. After deglycosylation, the preparation should no longer be stainable with PAS reagent. Only then 2-11M1 will react should MUC5AC be present. This mucin is present in primary ovariar mucinous cancer but usually absent in colorectal adenocarcinoma, thus showin an expression pattern opposite to MUC2. Together with a panel of antibodies,	ical
Clone 9-13M1 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate DyLight 650 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 4586 Gene Symbol MUC5AC Species Human Specificity/Sensitivity This monoclonal antibody recognizes the peptide core of gastric mucin M1 (recently identified as Mucin 5AC). Its epitope is located in the N-terminal cysteine rich part of the peptide core of MUC5AC, which is heavily glycosylated its epitope is destroyed by beta-mercaptoethanol but not by periodate treatment monoclonal antibody 2-11M1 reacts with the protein backbone exclusively; it or eracts with fully deglycosylated. This can be achieved with standard periodate oxidation method. The success of the deglycosylation, the preparation should no longer be stainable with PAS reagent. Only then 2-11M1 will react should MUC5AC be present. This mucin is present in primary ovariar mucinous cancer but usually absent in colorectal adenocarcinoma, thus showing an expression pattern opposite to MUC2. Together with a panel of antibodies,	
Clone 9-13M1 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate DyLight 650 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 4586 Gene Symbol MUC5AC Species Human Specificity/Sensitivity This monoclonal antibody recognizes the peptide core of gastric mucin M1 (recently identified as Mucin 5AC). Its epitope is located in the N-terminal cysteine rich part of the peptide core of MUC5AC, which is heavily glycosylated its epitope is destroyed by beta-mercaptoethanol but not by periodate treatment monoclonal antibody 2-11M1 reacts with the protein backbone exclusively; it or exacts with fully deglycosylated. This can be achieved with standard periodate oxidation method. The success of the deglycosylation, the preparation should no longer be stainable with PAS reagent. Only then 2-11M1 will react should MUC5AC be present. This mucin is present in primary ovariar mucinous cancer but usually absent in colorectal adenocarcinoma, thus showing an expression pattern opposite to MUC2. Together with a panel of antibodies,	
Preservative Jostype	
Isotype IgG1 Kappa DyLight 650 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 4586 Gene Symbol MUC5AC Species Human Specificity/Sensitivity This monoclonal antibody recognizes the peptide core of gastric mucin M1 (recently identified as Mucin 5AC). Its epitope is located in the N-terminal cysteine rich part of the peptide core of MUC5AC, which is heavily glycosylate. Its epitope is destroyed by beta-mercaptoethanol but not by periodate treatmen monoclonal antibody 2-11M1 reacts with the protein backbone exclusively; it o reacts with fully deglycosylated. This can be achieved with standard periodate oxidation method. The success of the deglycosylation, can be checke with routine PAS (Periodic Acid Shiff) staining. After deglycosylation, the preparation should no longer be stainable with PAS reagent. Only then 2-11M1 will react should MUC5AC be present. This mucin is present in primary ovariar mucinous cancer but usually absent in colorectal adenocarcinoma, thus showing an expression pattern opposite to MUC2. Together with a panel of antibodies,	
Conjugate DyLight 650 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 4586 Gene Symbol MUC5AC Species Human This monoclonal antibody recognizes the peptide core of gastric mucin M1 (recently identified as Mucin 5AC). Its epitope is located in the N-terminal cysteine rich part of the peptide core of MUC5AC, which is heavily glycosylated. Its epitope is destroyed by beta-mercaptoethanol but not by periodate treatment monoclonal antibody 2-11M1 reacts with the protein backbone exclusively; it or reacts with fully deglycosylated. This can be achieved with standard periodate oxidation method. The success of the deglycosylation can be checked with routine PAS (Periodic Acid Shiff) staining. After deglycosylation, the preparation should no longer be stainable with PAS reagent. Only then 2-11M1 will react should MUC5AC be present. This mucin is present in primary ovariar mucinous cancer but usually absent in colorectal adenocarcinoma, thus showing an expression pattern opposite to MUC2. Together with a panel of antibodies,	
Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 4586 Gene Symbol MUC5AC Species Human Specificity/Sensitivity This monoclonal antibody recognizes the peptide core of gastric mucin M1 (recently identified as Mucin 5AC). Its epitope is located in the N-terminal cysteine rich part of the peptide core of MUC5AC, which is heavily glycosylated Its epitope is destroyed by beta-mercaptoethanol but not by periodate treatment monoclonal antibody 2-11M1 reacts with the protein backbone exclusively; it or reacts with fully deglycosylated. This can be achieved with standard periodate oxidation method. The success of the deglycosylation can be checked with routine PAS (Periodic Acid Shiff) staining. After deglycosylation, the preparation should no longer be stainable with PAS reagent. Only then 2-11M1 will react should MUC5AC be present. This mucin is present in primary ovariar mucinous cancer but usually absent in colorectal adenocarcinoma, thus showing an expression pattern opposite to MUC2. Together with a panel of antibodies,	
Product Description	
Product Description Host Mouse Gene ID 4586 Gene Symbol MUC5AC Species Human This monoclonal antibody recognizes the peptide core of gastric mucin M1 (recently identified as Mucin 5AC). Its epitope is located in the N-terminal cysteine rich part of the peptide core of MUC5AC, which is heavily glycosylated Its epitope is destroyed by beta-mercaptoethanol but not by periodate treatment monoclonal antibody 2-11M1 reacts with the protein backbone exclusively; it or reacts with fully deglycosylated MUC5AC. Therefore, the material under test should also be fully deglycosylated. This can be achieved with standard periodate oxidation method. The success of the deglycosylation can be checked with routine PAS (Periodic Acid Shiff) staining. After deglycosylation, the preparation should no longer be stainable with PAS reagent. Only then 2-11M2 will react should MUC5AC be present. This mucin is present in primary ovariar mucinous cancer but usually absent in colorectal adenocarcinoma, thus showing an expression pattern opposite to MUC2. Together with a panel of antibodies,	
Host Gene ID 4586 Gene Symbol MUC5AC Species Human This monoclonal antibody recognizes the peptide core of gastric mucin M1 (recently identified as Mucin 5AC). Its epitope is located in the N-terminal cysteine rich part of the peptide core of MUC5AC, which is heavily glycosylated Its epitope is destroyed by beta-mercaptoethanol but not by periodate treatmen monoclonal antibody 2-11M1 reacts with the protein backbone exclusively; it or reacts with fully deglycosylated MUC5AC. Therefore, the material under test should also be fully deglycosylated. This can be achieved with standard periodate oxidation method. The success of the deglycosylation can be checked with routine PAS (Periodic Acid Shiff) staining. After deglycosylation, the preparation should no longer be stainable with PAS reagent. Only then 2-11M2 will react should MUC5AC be present. This mucin is present in primary ovariar mucinous cancer but usually absent in colorectal adenocarcinoma, thus showing an expression pattern opposite to MUC2. Together with a panel of antibodies,	
Gene Symbol MUC5AC Species Human Specificity/Sensitivity This monoclonal antibody recognizes the peptide core of gastric mucin M1 (recently identified as Mucin 5AC). Its epitope is located in the N-terminal cysteine rich part of the peptide core of MUC5AC, which is heavily glycosylated Its epitope is destroyed by beta-mercaptoethanol but not by periodate treatmen monoclonal antibody 2-11M1 reacts with the protein backbone exclusively; it o reacts with fully deglycosylated MUC5AC. Therefore, the material under test should also be fully deglycosylated. This can be achieved with standard periodate oxidation method. The success of the deglycosylation can be checked with routine PAS (Periodic Acid Shiff) staining. After deglycosylation, the preparation should no longer be stainable with PAS reagent. Only then 2-11M2 will react should MUC5AC be present. This mucin is present in primary ovariar mucinous cancer but usually absent in colorectal adenocarcinoma, thus showing an expression pattern opposite to MUC2. Together with a panel of antibodies,	
Gene Symbol MUC5AC Species Human This monoclonal antibody recognizes the peptide core of gastric mucin M1 (recently identified as Mucin 5AC). Its epitope is located in the N-terminal cysteine rich part of the peptide core of MUC5AC, which is heavily glycosylated Its epitope is destroyed by beta-mercaptoethanol but not by periodate treatment monoclonal antibody 2-11M1 reacts with the protein backbone exclusively; it or reacts with fully deglycosylated MUC5AC. Therefore, the material under test should also be fully deglycosylated. This can be achieved with standard periodate oxidation method. The success of the deglycosylation can be checked with routine PAS (Periodic Acid Shiff) staining. After deglycosylation, the preparation should no longer be stainable with PAS reagent. Only then 2-11M2 will react should MUC5AC be present. This mucin is present in primary ovariar mucinous cancer but usually absent in colorectal adenocarcinoma, thus showing an expression pattern opposite to MUC2. Together with a panel of antibodies,	
Specificity/Sensitivity This monoclonal antibody recognizes the peptide core of gastric mucin M1 (recently identified as Mucin 5AC). Its epitope is located in the N-terminal cysteine rich part of the peptide core of MUC5AC, which is heavily glycosylated Its epitope is destroyed by beta-mercaptoethanol but not by periodate treatment monoclonal antibody 2-11M1 reacts with the protein backbone exclusively; it or reacts with fully deglycosylated MUC5AC. Therefore, the material under test should also be fully deglycosylated. This can be achieved with standard periodate oxidation method. The success of the deglycosylation can be checked with routine PAS (Periodic Acid Shiff) staining. After deglycosylation, the preparation should no longer be stainable with PAS reagent. Only then 2-11M2 will react should MUC5AC be present. This mucin is present in primary ovariar mucinous cancer but usually absent in colorectal adenocarcinoma, thus showing an expression pattern opposite to MUC2. Together with a panel of antibodies,	
This monoclonal antibody recognizes the peptide core of gastric mucin M1 (recently identified as Mucin 5AC). Its epitope is located in the N-terminal cysteine rich part of the peptide core of MUC5AC, which is heavily glycosylated Its epitope is destroyed by beta-mercaptoethanol but not by periodate treatment monoclonal antibody 2-11M1 reacts with the protein backbone exclusively; it or reacts with fully deglycosylated MUC5AC. Therefore, the material under test should also be fully deglycosylated. This can be achieved with standard periodate oxidation method. The success of the deglycosylation can be checked with routine PAS (Periodic Acid Shiff) staining. After deglycosylation, the preparation should no longer be stainable with PAS reagent. Only then 2-11M2 will react should MUC5AC be present. This mucin is present in primary ovariar mucinous cancer but usually absent in colorectal adenocarcinoma, thus showing an expression pattern opposite to MUC2. Together with a panel of antibodies,	
(recently identified as Mucin 5AC). Its epitope is located in the N-terminal cysteine rich part of the peptide core of MUC5AC, which is heavily glycosylated Its epitope is destroyed by beta-mercaptoethanol but not by periodate treatment monoclonal antibody 2-11M1 reacts with the protein backbone exclusively; it or reacts with fully deglycosylated MUC5AC. Therefore, the material under test should also be fully deglycosylated. This can be achieved with standard periodate oxidation method. The success of the deglycosylation can be checked with routine PAS (Periodic Acid Shiff) staining. After deglycosylation, the preparation should no longer be stainable with PAS reagent. Only then 2-11M2 will react should MUC5AC be present. This mucin is present in primary ovariar mucinous cancer but usually absent in colorectal adenocarcinoma, thus showing an expression pattern opposite to MUC2. Together with a panel of antibodies,	
Anti-MUC5AC may be useful for differential identification of primary mucinous ovarian tumors from colon adenocarcinoma metastatic to the ovary. MUC5AC antibodies may also be useful for identification of intestinal metaplasia as well in the identification of pancreatic carcinoma and pre-cancerous changes vs. normal pancreas.	lated. tment. it only st ecked 1M1 arian howing ies, bus
Immunogen M1 mucin preparation from the fluid of an ovarian mucinous cyst belonging to a O Le(a-b) patient	
Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries	S.
Product Application Details	to an
Applications ELISA, Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunofluorescence	to an
Recommended Dilutions Flow Cytometry, ELISA, Immunocytochemistry/Immunofluorescence, Immunofluorescence	to an
Application Notes Optimal dilution of this antibody should be experimentally determined.	to an
ovarian tumors from colon adenocarcinoma metastatic to the ovary. MUC5AC antibodies may also be useful for identification of intestinal metaplasia as well in the identification of pancreatic carcinoma and pre-cancerous changes vs. normal pancreas. Immunogen M1 mucin preparation from the fluid of an ovarian mucinous cyst belonging to a O Le(a-b) patient	owing ies, ous





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom

Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP3-11484C

NBP1-43319C Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1) [DyLight 650]

H00004586-Q01-10ug Recombinant Human MUC5AC GST (N-Term) Protein

210-TA-005 TNF-alpha [Unconjugated]

NBP2-76703 Human MUC5AC ELISA Kit (Colorimetric)

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-11484C

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications



