# **Product Datasheet**

## Myeloid-Associated Differentiation Marker Antibody (MYADM/971) [Alexa Fluor® 700] NBP3-11489AF700

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-11489AF700

Updated 10/26/2023 v.20.1

Earn rewards for product reviews and publications.



Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-11489AF700

#### NBP3-11489AF700

Myeloid-Associated Differentiation Marker Antibody (MYADM/971) [Alexa Fluor® 700]

Product InformationUnit Size0.1 mlConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneMYADM/971Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 700PurityProtein A or G purifiedBuffer500m Modium BorateProduct Description91663Gene ID91663Gene SymbolMYADMSpeciesHuman, Rhesus MacaqueReactivity NotesMacaque Monkey.MarkerMyeloid-Associated Differentiation MarkerSpecificity/SensitivityRecognizes a myeloid associated differentiation antigen in the cytoplasm of mature granulocytes. It shows no reactivity with any other cell type in human tissues. Markers of myeloid cells are useful in the identification of different levels of cellular differentiation. It reacts with early precursor and mature forms of human and monkey myeloid cells. This monoclonal antibody to the detection of myeloid leukemias and granulocytic sarcomas. It can be used as a marker of granulocytes in normal tissues or inflammatory processes.ImmunogenRecombinant full-length human Myeloid-Associated Differentiation Marker	-	
ConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneMYADM/971Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 700PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description91663HostMoseGene SymbolMYADMSpeciesHuman, Rhesus MacaqueReactivity NotesMacaque Monkey.MarkerMyeloid-Associated Differentiation MarkerSpecificity/SensitivityRecognizes a myeloid cells are useful in the identification of different levels of cellular differentiation. It reacts with early processes.ImmunogenRecombinant full-length human Myeloid-Associated Differentiation Marker	Product Information	
services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneMYADM/971Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 700PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description91663Gene ID91663Gene SymbolMYADMSpeciesHuman, Rhesus MacaqueReactivity NotesMacaque Monkey.MarkerMyeloid-Associated Differentiation MarkerSpecificity/SensitivityRecognizes a myeloid associated differentiation of different levels of cellular differentiation. It reacts with early precursor and mature forms of human and monkey myeloid cells. This monoclonal antibody is useful for the detection of myeloid leukemias and granulocytics arcomas. It can be used as a marker of granulocytes in normal tissues or inflammatory processes.ImmunogenRecombinant full-length human Myeloid-Associated Differentiation Marker	Unit Size	0.1 ml
ClonalityMonoclonalCloneMYADM/971Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 700PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description91663Gene ID91663Gene SymbolMYADMSpeciesHuman, Rhesus MacaqueReactivity NotesMacaque Monkey.MarkerMyeloid-Associated Differentiation MarkerSpecificity/SensitivityRecognizes a myeloid associated differentiation antigen in the cytoplasm of mature granulocytes. It shows no reactivity with any other cell type in human tissues. Markers of myeloid cells are useful in the identification of different levels of cellular differentiation. It reacts with early precursor and mature forms of human and monkey myeloid cells. This monoclonal antibody is useful for the detection of myeloid leukemias and granulocytic sarcomas. It can be used as a marker of granulocytes in normal tissues or inflammatory processes.ImmunogenRecombinant full-length human Myeloid-Associated Differentiation Marker	Concentration	•
CloneMYADM/971Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 700PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseGene ID91663Gene SymbolMYADMSpeciesHuman, Rhesus MacaqueReactivity NotesMacaque Monkey.MarkerMyeloid-Associated Differentiation MarkerSpecificity/SensitivityRecognizes a myeloid associated differentiation antigen in the cytoplasm of mature granulocytes. It shows no reactivity with any other cell type in human tissue. Markers of myeloid cells are useful in the identification of different levels of cellular differentiation. It reacts with early procursor and mature forms of human and monkey myeloid cells. This monoclonal antibody is useful for the detection of myeloid leukemias and granulocytic sarcomas. It can be used as a marker of granulocytes in normal tissues or inflammatory processes.ImmunogenRecombinant full-length human Myeloid-Associated Differentiation Marker	Storage	Store at 4C in the dark.
Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 700PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID91663Gene SymbolMYADMSpeciesHuman, Rhesus MacaqueReactivity NotesMacaque Monkey.MarkerMyeloid-Associated Differentiation MarkerSpecificity/SensitivityRecognizes a myeloid associated differentiation antigen in the cytoplasm of mature granulocytes. It shows no reactivity with any other cell type in human tissues. Markers of myeloid cells are useful in the identification of different levels of cellular differentiation. It reacts with early precursor and mature forms of human and monkey myeloid cells. This monoclonal antibody is useful for the detection of myeloid leukemias and granulocytes in normal tissues or inflammatory processes.ImmunogenRecombinant full-length human Myeloid-Associated Differentiation Marker	Clonality	Monoclonal
IsotypeIgG1 KappaConjugateAlexa Fluor 700PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID91663Gene SymbolMYADMSpeciesHuman, Rhesus MacaqueReactivity NotesMacaque Monkey.MarkerMyeloid-Associated Differentiation MarkerSpecificity/SensitivityRecognizes a myeloid associated differentiation antigen in the cytoplasm of mature granulocytes. It shows no reactivity with any other cell type in human tissues. Markers of myeloid cells are useful in the identification of different levels of cellular differentiation. It reacts with early precursor and mature forms of human and monkey myeloid cells. This monoclonal antibody is useful for the detection of myeloid leukemias and granulocytic sarcomas. It can be used as a marker of granulocytes in normal tissues or inflammatory processes.ImmunogenRecombinant full-length human Myeloid-Associated Differentiation Marker	Clone	MYADM/971
ConjugateAlexa Fluor 700PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID91663Gene SymbolMYADMSpeciesHuman, Rhesus MacaqueReactivity NotesMacaque Monkey.MarkerMyeloid-Associated Differentiation MarkerSpecificity/SensitivityRecognizes a myeloid associated differentiation antigen in the cytoplasm of mature granulocytes. It shows no reactivity with any other cell type in human tissues. Markers of myeloid cells are useful in the identification of different levels of cellular differentiation. It reacts with early precursor and mature forms of human and monkey myeloid cells. This monoclonal antibody is useful for the detection of myeloid leukemias and granulocytic sarcomas. It can be used as a marker of granulocytes in normal tissues or inflammatory processes.ImmunogenRecombinant full-length human Myeloid-Associated Differentiation Marker	Preservative	0.05% Sodium Azide
PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID91663Gene SymbolMYADMSpeciesHuman, Rhesus MacaqueReactivity NotesMacaque Monkey.MarkerMyeloid-Associated Differentiation MarkerSpecificity/SensitivityRecognizes a myeloid associated differentiation antigen in the cytoplasm of mature granulocytes. It shows no reactivity with any other cell type in human tissues. Markers of myeloid cells are useful in the identification of different levels of cellular differentiation. It reacts with early precursor and mature forms of human and monkey myeloid cells. This monoclonal antibody is useful for the detection of myeloid leukemias and granulocyte sarcomas. It can be used as a marker of granulocytes in normal tissues or inflammatory processes.ImmunogenRecombinant full-length human Myeloid-Associated Differentiation Marker	Isotype	IgG1 Kappa
Buffer50mM Sodium BorateProduct DescriptionHostMouseGene ID91663Gene SymbolMYADMSpeciesHuman, Rhesus MacaqueReactivity NotesMacaque Monkey.MarkerMyeloid-Associated Differentiation MarkerSpecificity/SensitivityRecognizes a myeloid associated differentiation antigen in the cytoplasm of mature granulocytes. It shows no reactivity with any other cell type in human tissues. Markers of myeloid cells are useful in the identification of different levels of cellular differentiation. It reacts with early precursor and mature forms of human and monkey myeloid cells. This monoclonal antibody is useful for the detection of myeloid leukemias and granulocytic sarcomas. It can be used as a marker of granulocytes in normal tissues or inflammatory processes.ImmunogenRecombinant full-length human Myeloid-Associated Differentiation Marker	Conjugate	Alexa Fluor 700
Product DescriptionHostMouseGene ID91663Gene SymbolMYADMSpeciesHuman, Rhesus MacaqueReactivity NotesMacaque Monkey.MarkerMyeloid-Associated Differentiation MarkerSpecificity/SensitivityRecognizes a myeloid associated differentiation antigen in the cytoplasm of mature granulocytes. It shows no reactivity with any other cell type in human tissues. Markers of myeloid cells are useful in the identification of different levels of cellular differentiation. It reacts with early precursor and mature forms of human and monkey myeloid cells. This monoclonal antibody is useful for the detection of myeloid leukemias and granulocytic sarcomas. It can be used as a marker of granulocytes in normal tissues or inflammatory processes.ImmunogenRecombinant full-length human Myeloid-Associated Differentiation Marker	Purity	Protein A or G purified
HostMouseGene ID91663Gene SymbolMYADMSpeciesHuman, Rhesus MacaqueReactivity NotesMacaque Monkey.MarkerMyeloid-Associated Differentiation MarkerSpecificity/SensitivityRecognizes a myeloid associated differentiation antigen in the cytoplasm of mature granulocytes. It shows no reactivity with any other cell type in human tissues. Markers of myeloid cells are useful in the identification of different levels of cellular differentiation. It reacts with early precursor and mature forms of human and monkey myeloid cells. This monoclonal antibody is useful for the detection of myeloid leukemias and granulocytic sarcomas. It can be used as a marker of granulocytes in normal tissues or inflammatory processes.ImmunogenRecombinant full-length human Myeloid-Associated Differentiation Marker	Buffer	50mM Sodium Borate
Gene ID91663Gene SymbolMYADMSpeciesHuman, Rhesus MacaqueReactivity NotesMacaque Monkey.MarkerMyeloid-Associated Differentiation MarkerSpecificity/SensitivityRecognizes a myeloid associated differentiation antigen in the cytoplasm of mature granulocytes. It shows no reactivity with any other cell type in human tissues. Markers of myeloid cells are useful in the identification of different levels of cellular differentiation. It reacts with early precursor and mature forms of human and monkey myeloid cells. This monoclonal antibody is useful for the detection of myeloid leukemias and granulocytic sarcomas. It can be used as a marker of granulocytes in normal tissues or inflammatory processes.ImmunogenRecombinant full-length human Myeloid-Associated Differentiation Marker	Product Description	
Gene SymbolMYADMSpeciesHuman, Rhesus MacaqueReactivity NotesMacaque Monkey.MarkerMyeloid-Associated Differentiation MarkerSpecificity/SensitivityRecognizes a myeloid associated differentiation antigen in the cytoplasm of mature granulocytes. It shows no reactivity with any other cell type in human tissues. Markers of myeloid cells are useful in the identification of different levels of cellular differentiation. It reacts with early precursor and mature forms of human and monkey myeloid cells. This monoclonal antibody is useful for the detection of myeloid leukemias and granulocytic sarcomas. It can be used as a marker of granulocytes in normal tissues or inflammatory processes.ImmunogenRecombinant full-length human Myeloid-Associated Differentiation Marker	Host	Mouse
SpeciesHuman, Rhesus MacaqueReactivity NotesMacaque Monkey.MarkerMyeloid-Associated Differentiation MarkerSpecificity/SensitivityRecognizes a myeloid associated differentiation antigen in the cytoplasm of mature granulocytes. It shows no reactivity with any other cell type in human tissues. Markers of myeloid cells are useful in the identification of different levels of cellular differentiation. It reacts with early precursor and mature forms of human and monkey myeloid cells. This monoclonal antibody is useful for the detection of myeloid leukemias and granulocytic sarcomas. It can be used as a marker of granulocytes in normal tissues or inflammatory processes.ImmunogenRecombinant full-length human Myeloid-Associated Differentiation Marker	Gene ID	91663
Reactivity NotesMacaque Monkey.MarkerMyeloid-Associated Differentiation MarkerSpecificity/SensitivityRecognizes a myeloid associated differentiation antigen in the cytoplasm of mature granulocytes. It shows no reactivity with any other cell type in human tissues. Markers of myeloid cells are useful in the identification of different levels of cellular differentiation. It reacts with early precursor and mature forms of human and monkey myeloid cells. This monoclonal antibody is useful for the detection of myeloid leukemias and granulocytic sarcomas. It can be used as a marker of granulocytes in normal tissues or inflammatory processes.ImmunogenRecombinant full-length human Myeloid-Associated Differentiation Marker	Gene Symbol	MYADM
MarkerMyeloid-Associated Differentiation MarkerSpecificity/SensitivityRecognizes a myeloid associated differentiation antigen in the cytoplasm of mature granulocytes. It shows no reactivity with any other cell type in human tissues. Markers of myeloid cells are useful in the identification of different levels of cellular differentiation. It reacts with early precursor and mature forms of human and monkey myeloid cells. This monoclonal antibody is useful for the detection of myeloid leukemias and granulocytic sarcomas. It can be used as a marker of granulocytes in normal tissues or inflammatory processes.ImmunogenRecombinant full-length human Myeloid-Associated Differentiation Marker	Species	Human, Rhesus Macaque
Specificity/SensitivityRecognizes a myeloid associated differentiation antigen in the cytoplasm of mature granulocytes. It shows no reactivity with any other cell type in human tissues. Markers of myeloid cells are useful in the identification of different levels of cellular differentiation. It reacts with early precursor and mature forms of human and monkey myeloid cells. This monoclonal antibody is useful for the detection of myeloid leukemias and granulocytic sarcomas. It can be used as a marker of granulocytes in normal tissues or inflammatory processes.ImmunogenRecombinant full-length human Myeloid-Associated Differentiation Marker	Reactivity Notes	Macaque Monkey.
mature granulocytes. It shows no reactivity with any other cell type in human tissues. Markers of myeloid cells are useful in the identification of different levels of cellular differentiation. It reacts with early precursor and mature forms of human and monkey myeloid cells. This monoclonal antibody is useful for the detection of myeloid leukemias and granulocytic sarcomas. It can be used as a marker of granulocytes in normal tissues or inflammatory processes.ImmunogenRecombinant full-length human Myeloid-Associated Differentiation Marker	Marker	Myeloid-Associated Differentiation Marker
	Specificity/Sensitivity	mature granulocytes. It shows no reactivity with any other cell type in human tissues. Markers of myeloid cells are useful in the identification of different levels of cellular differentiation. It reacts with early precursor and mature forms of human and monkey myeloid cells. This monoclonal antibody is useful for the detection of myeloid leukemias and granulocytic sarcomas. It can be used as a marker of granulocytes in normal tissues or inflammatory processes.
	Immunogen	



	Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.
Product Application Details	
Applications	Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunofluorescence
Recommended Dilutions	Flow Cytometry, Immunohistochemistry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence
Application Notes	Optimal dilution of this antibody should be experimentally determined.

Notes





#### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

### **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

#### Products Related to NBP3-11489AF700

IC002NMouse IgG1 Isotype Control (11711) [Alexa Fluor® 700]7954-GM-010/CFGM-CSF [Unconjugated]203-IL-010IL-3 [Unconjugated]214-CS-005G-CSF [Unconjugated]

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-11489AF700

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

