# **Product Datasheet**

# ICAM-1/CD54 Antibody (W-CAM-1 (same as Wehi-CAM-1 or 1H4)) [mFluor Violet 500 SE] NBP3-11490MFV500

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-11490MFV500

Updated 10/26/2023 v.20.1

Earn rewards for product reviews and publications.



Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-11490MFV500

#### NBP3-11490MFV500

ICAM-1/CD54 Antibody (W-CAM-1 (same as Wehi-CAM-1 or 1H4)) [mFluor Violet 500 SE]

| Product Information         |  |
|-----------------------------|--|
| Unit Size                   | 0.1 ml   |
| Concentration               | Please see the vial label for concentration. If unlisted please contact technical services.  |
| Storage                     | Store at 4C in the dark.   |
| Clonality                   | Monoclonal   |
| Clone                       | W-CAM-1 (same as Wehi-CAM-1 or 1H4)  |
| Preservative                | 0.05% Sodium Azide   |
| Isotype                     | IgG2b Kappa  |
| Conjugate                   | mFluor Violet 500 SE   |
| Purity                      | Protein A or G purified  |
| Buffer                      | 50mM Sodium Borate   |
| Product Description         |  |
| Host                        | Mouse  |
| Gene ID                     | 3383   |
| Gene Symbol                 | ICAM1  |
| Species                     | Human  |
| Specificity/Sensitivity     | Recognizes an 85-115kDa protein (variation with cell type), identified as intercellular adhesion molecule (ICAM-1) (Workshop IV). It has 7 potential N-linked glycosylation sites. ICAM-1 is a single chain glycoprotein of Ig supergene family, present on unstimulated endothelial cells (EC) and on a variety of other cell types including activated fibroblasts, EC, macrophages, and lymphocytes. ICAM-1 mediates cell adhesion by binding to integrins CD11a/CD18 (leukocyte adhesion molecule, LFA-1) and to CD11b/CD18 (Mac-1). This interaction enhances antigen-specific T-cell activation. ICAM-1 also binds to CD43 and to Plasmodium falciparum infected RBCs. W-CAM-1 monoclonal antibody blocks aggregation of cell lines mediated by the ICAM-1 and blocks homotypic binding of Purified populations of activated T- and B-lymphocytes and also aggregation of mixed T- and B-cell blasts. It inhibits T-cell adhesion to normal human endothelial cells. Activation induced by cell-cell contact (mixed lymphocyte reaction, T-cell mediated B-cell activation) is significantly inhibited. This monoclonal antibody blocks elements of both effector arms of immune system (cytotoxic cell function and Ig production). |
| Immunogen                   | Raji Burkitt lymphoma cells  |
| Notes                       | mFluor(TM) is a trademark of AAT Bioquest, Inc. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.  |
| Product Application Details |  |
| Applications                | Flow Cytometry, Immunocytochemistry/Immunofluorescence,<br>Immunohistochemistry, Immunohistochemistry-Paraffin, Immunofluorescence   |
| Recommended Dilutions       | Flow Cytometry, Immunohistochemistry,<br>Immunocytochemistry/Immunofluorescence, Immunohistochemistry-Paraffin,<br>Immunofluorescence  |
| Application Notes           | Optimal dilution of this antibody should be experimentally determined.   |
|                             |  |



#### Images

ICAM-1/CD54 Antibody (W-CAM-1 (same as Wehi-CAM-1 or 1H4)) [mFluor Violet 500 SE] - Vial of mFluor Violet 500 conjugated antibody. mFluor Violet 500 is optimally excited at 410 nm by the Violet laser (405 nm) and has an emission maximum of 501 nm.







#### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

# **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

# **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

### Products Related to NBP3-11490MFV500

| NBP2-52268-0.05mg | Recombinant Mouse ICAM-1/CD54 His Protein |
|-------------------|---|
| 210-TA-005        | TNF-alpha [Unconjugated]                  |
| 796-IC-050        | ICAM-1/CD54 [Unconjugated]                |
| D6050             | IL-6 [HRP]                                |

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-11490MFV500

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

