Product Datasheet

WT1 Antibody (WT1/1434R) [CoraFluor™ 1] NBP3-11622CL1

Unit Size: 0.1 ml

Store at 4C in the dark. Do not freeze.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-11622CL1

Updated 8/13/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP3-11622CL1



NBP3-11622CL1

WT1 Antibody (WT1/1434R) [CoraFluor™ 1]

Will Allibody (Will/1454R) [Colai Idol 1]		
Product Information		
Unit Size	0.1 ml	
Concentration	Please see the vial label for concentration. If unlisted please contact technical services.	
Storage	Store at 4C in the dark. Do not freeze.	
Clonality	Monoclonal	
Clone	WT1/1434R	
Preservative	No Preservative	
Isotype	IgG	
Conjugate	CoraFluor 1	
Purity	Protein A or G purified	
Buffer	PBS	
Product Description		
Description	CoraFluor(TM) 1 is a high performance terbium-based TR-FRET (Time-Resolved Fluorescence Resonance Energy Transfer) or TRF (Time-Resolved Fluorescence) donor for high throughput assay development. CoraFluor(TM) 1 absorbs UV light at approximately 340 nm, and emits at approximately 490 nm, 545 nm, 585 nm and 620 nm. It is compatible with common acceptor dyes that absorb at the emission wavelengths of CoraFluor(TM) 1. CoraFluor(TM) 1 can be used for the development of robust and scalable TR-FRET binding assays such as target engagement, ternary complex, protein-protein interaction and protein quantification assays. CoraFluor(TM) 1, amine reactive CoraFluor(TM) 1, thiol reactive For more information, please see our CoraFluor(TM) TR-FRET technology flyer.	
Host	Rabbit	
Gene ID	7490	
Gene Symbol	WT1	
Species	Human	
Marker	Wilm's Tumor & Mesothelial Marker	
Specificity/Sensitivity	Recognizes a 47-55kDa-tumor suppressor protein, identified as Wilms Tumor (WT1) protein. The antibody reacts with all isoforms of the full-length WT1 and also identifies WT1 lacking exon 2-encoded amino acids, frequently found in subsets of sporadic Wilm s tumors.WT1, a sporadic and familial pediatric kidney tumor, is genetically heterogeneous. Wilm s tumor is associated with mutations of WT1, a zinc-finger transcription factor that is essential for the development of the metanephric kidney and the urogenital system. The WT1 gene is normally expressed in fetal kidney and mesothelium, and its expression has been suggested as a marker for Wilm s tumor and mesothelioma. WT1 protein has been identified in proliferative mesothelial cells, malignant mesothelioma, ovarian carcinoma, gonadoblastoma, nephroblastoma, and desmoplastic small round cell tumor. Lung adenocarcinomas rarely stain positive with this antibody. WT1 protein expression in mesothelial cells has become a reliable marker for the diagnosis of mesotheliomas.	



ImmunogenRecombinant human full-length WT1 protein (Uniprot: P19544)NotesCoraFluor (TM) is a trademark of Bio-Techne Corp. Sold for research purposes only under agreement from Massachusetts General Hospital. US patent 2022/0025254Product Application DetailsImmunohistochemistry-Paraffin, ImmunohistochemistryApplicationsImmunohistochemistry, Immunohistochemistry-ParaffinApplication NotesOptimal dilution of this antibody should be experimentally determined.		
only under agreement from Massachusetts General Hospital. US patent 2022/0025254 Product Application Details Applications Immunohistochemistry-Paraffin, Immunohistochemistry Recommended Dilutions Immunohistochemistry, Immunohistochemistry-Paraffin	Immunogen	Recombinant human full-length WT1 protein (Uniprot: P19544)
Applications Immunohistochemistry-Paraffin, Immunohistochemistry Recommended Dilutions Immunohistochemistry, Immunohistochemistry-Paraffin	Notes	only under agreement from Massachusetts General Hospital. US patent
Recommended Dilutions Immunohistochemistry, Immunohistochemistry-Paraffin	Product Application Details	
	Applications	Immunohistochemistry-Paraffin, Immunohistochemistry
Application Notes Optimal dilution of this antibody should be experimentally determined.	Recommended Dilutions	Immunohistochemistry, Immunohistochemistry-Paraffin
	Application Notes	Optimal dilution of this antibody should be experimentally determined.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP3-11622CL1

H00007490-Q01-10ug Recombinant Human WT1 GST (N-Term) Protein

292-G2-050 IGF-II/IGF2 [Unconjugated]
NBP2-11141 WT1 Overexpression Lysate
AF3075 SOX9 Antibody [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-11622CL1

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

