Product Datasheet

Mouse Pure-Blot anti-Rabbit IgG (H+L) Secondary Antibody (eB182) [HRP] NBP3-11668

Unit Size: 200 ul

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-11668

Updated 8/8/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-11668



NBP3-11668


Mouse Pure-Blot anti-Rabbit IgG (H+L) Secondary Antibody (eB182) [HRP]

Storage Store at -20C. Avoid freeze-thaw cycles. Clonality Monoclonal Clone eB182 Preservative No Preservative Isotype IgG Conjugate HRP Purity Protein G purified Buffer 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 0.1 mg/ml Bovine Serum Albumin (BSA, IgG and Protease free), 50% (v/v) Glycerol Product Description Prepared from tissue culture supernatant by Protein G affinity chromatograp Assay by Immunoelectrophoresis resulted in a single precipitin arc against A Rabbit Serum. Host Mouse Species Rabbit Specificity/Sensitivity Reactivity is observed against native Rabbit IgG by both Western blot and ELISA. Immunogen Rabbit IgG Product Application Details Applications Western Blot, ELISA, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation, In-situ Hybridization, Chromatin Immunoprecipitation (ChIP) Recommended Dilutions Western Blot 1:1000, ELISA, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation, In-situ Hybridization, Chromatin Immunoprecipitation (ChIP) Application Notes This secondary antibody may also be used for detection in immunoblotting assays that do not employ immunoprecipitation. It is provided as 1000X solutor occurrence are three key procedural considerations: 1. Protein A or G should not be used for the immunoprecipitation. Use of processing the search of the immunoprecipi	Mouse Pure-Blot anti-Rabbit IgG (H+L) Secondary Antibody (eB182) [HRP]	
Concentration Please see the vial label for concentration. If unlisted please contact technic services. Storage Store at -20C. Avoid freeze-thaw cycles. Clonality Monoclonal Clone B182 Preservative No Preservative Isotype IgG Conjugate HRP Purity Protein G purified Buffer 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 0.1 mg/ml Bovine Serum Albumin (BSA, IgG and Protease free), 50% (v/v) Glycerol Product Description Prepared from tissue culture supernatant by Protein G affinity chromatograp Assay by Immunoelectrophoresis resulted in a single precipitin arc against A Rabbit Serum. Host Mouse Species Rabbit Specificity/Sensitivity Reactivity is observed against native Rabbit IgG by both Western blot and ELISA. Immunogen Rabbit IgG Product Application Details Applications Western Blot, ELISA, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation, In-situ Hybridization, Chromatin Immunoprecipitation (ChIP) Recommended Dilutions Western Blot 1:1000, ELISA, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation, In-situ Hybridization, Chromatin Immunoprecipitation (ChIP) Application Notes This secondary antibody may also be used for detection in immunoblotting assays that do not employ immunoprecipitation. It is provided as 1000X solutor onesered the plots from mininges in sealed bags (removing as much air as possible) with minimal volume (2-3 m used conservatively at 2.5mls/blot will yield enough reagent for 80 blots. Note that there are three key procedural considerations: 1. Protein A or G should not be used for the immunoprecipitation. Use of procedural considerations: 1. Protein A or G should not be used for the immunoprecipitation. Use of procedural considerations: 1. Protein A or G should not be used for the immunoprecipitation. Use of procedural considerations.	Product Information	
Storage Stora at -20C. Avoid freeze-thaw cycles. Clonality Monoclonal Clone eB182 Preservative No Preservative Isotype IgG Conjugate HRP Purity Protein G purified Buffer 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 0.1 mg/ml Bovine Serum Albumin (BSA, IgG and Protease free), 50% (v/v) Glycerol Product Description Prepared from tissue culture supernatant by Protein G affinity chromatograp Assay by Immunoelectrophoresis resulted in a single precipitin arc against A Rabbit Serum. Host Mouse Species Rabbit Specificity/Sensitivity Reactivity is observed against native Rabbit IgG by both Western blot and ELISA. Immunogen Rabbit IgG Product Application Details Applications Western Blot, ELISA, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation, In-situ Hybridization, Chromatin Immunoprecipitation (ChIP) Recommended Dilutions Western Blot 1:1000, ELISA, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation, In-situ Hybridization, Chromatin Immunoprecipitation (ChIP) Application Notes This secondary antibody may also be used for detection in immunoblotting assays that do not employ immunoprecipitation. It is provided as 1000X solutor occurrence are three key procedural considerations: 1. Protein A or G should not be used for the immunoprecipitation. Use of processing the search of the immunoprecipi	Unit Size	200 ul
Clone	Concentration	Please see the vial label for concentration. If unlisted please contact technical services.
Clone eB182 Preservative No Preservative Isotype IgG Conjugate HRP Purity Protein G purified Buffer 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 0.1 mg/ml Bovine Serum Albumin (BSA, IgG and Protease free), 50% (v/v) Glycerol Product Description Prepared from tissue culture supernatant by Protein G affinity chromatograp Assay by Immunoelectrophoresis resulted in a single precipitin arc against A Rabbit Serum. Host Mouse Species Rabbit Specificity/Sensitivity Reactivity is observed against native Rabbit IgG by both Western blot and ELISA. Immunogen Rabbit IgG Product Application Details Applications Western Blot, ELISA, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation, In-situ Hybridization, Chromatin Immunoprecipitation (ChIP) Recommended Dilutions Western Blot 1:1000, ELISA, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation, In-situ Hybridization, Chromatin Immunoprecipitation (ChIP) Application Notes This secondary antibody may also be used for detection in immunoblotting assays that do not employ immunoprecipitation. It is provided as 1000X soluto to conserve reagent, we recommend incubating the blots from minigels in sealed bags (removing as much air as possible) with minimal volume (2-3 m used conservatively at 2.5mls/blot will yield enough reagent for 80 blots. Note that there are three key procedural considerations: 1. Protein A or G should not be used for the immunoprecipitation. Use of processors.	Storage	Store at -20C. Avoid freeze-thaw cycles.
Preservative No Preservative Isotype IgG Conjugate HRP Purity Protein G purified Buffer 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 0.1 mg/ml Bovine Serum Albumin (BSA, IgG and Protease free), 50% (v/v) Glycerol Product Description Prepared from tissue culture supernatant by Protein G affinity chromatograp Assay by Immunoelectrophoresis resulted in a single precipitin arc against A Rabbit Serum. Host Mouse Species Rabbit Specificity/Sensitivity Reactivity is observed against native Rabbit IgG by both Western blot and ELISA. Immunogen Rabbit IgG Product Application Details Applications Western Blot, ELISA, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation, In-situ Hybridization, Chromatin Immunoprecipitation (ChIP) Recommended Dilutions Western Blot 1:1000, ELISA, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation, In-situ Hybridization, Chromatin Immunoprecipitation (ChIP) Application Notes This secondary antibody may also be used for detection in immunoblotting assays that do not employ immunoprecipitation. It is provided as 1000X solu To conserve reagent, we recommend incubating the blots from minigels in sealed bags (removing as much air as possible) with minimal volume (2-3 m used conservatively at 2.5mls/blot will yield enough reagent for 80 blots. Note that there are three key procedural considerations: 1. Protein A or G should not be used for the immunoprecipitation. Use of processing the procedural considerations.	Clonality	Monoclonal
Isotype Conjugate HRP Purity Protein G purified Buffer 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 0.1 mg/ml Bovine Serum Albumin (BSA, IgG and Protease free), 50% (v/v) Glycerol Product Description Prepared from tissue culture supernatant by Protein G affinity chromatograp Assay by Immunoelectrophoresis resulted in a single precipitin arc against A Rabbit Serum. Host Mouse Species Rabbit Specificity/Sensitivity Reactivity is observed against native Rabbit IgG by both Western blot and ELISA. Immunogen Rabbit IgG Product Application Details Applications Western Blot, ELISA, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation, In-situ Hybridization, Chromatin Immunoprecipitation (ChIP) Recommended Dilutions Western Blot 1:1000, ELISA, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation, In-situ Hybridization, Chromatin Immunoprecipitation (ChIP) Application Notes This secondary antibody may also be used for detection in immunoblotting assays that do not employ immunoprecipitation. It is provided as 1000X soluto conserve reagent, we recommend incubating the blots from minigels in sealed bags (removing as much air as possible) with minimal volume (2-3 mused conservatively at 2.5mls/blot will yield enough reagent for 80 blots. Note that there are three key procedural considerations: 1. Protein A or G should not be used for the immunoprecipitation. Use of procedural considerations: 1. Protein A or G should not be used for the immunoprecipitation. Use of procedural considerations: 1. Protein A or G should not be used for the immunoprecipitation. Use of procedural considerations: 1. Protein A or G should not be used for the immunoprecipitation. Use of procedural considerations: 1. Protein A or G should not be used for the immunoprecipitation. Use of procedural considerations:	Clone	eB182
Conjugate HRP Purity Protein G purified Buffer 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 0.1 mg/ml Bovine Serum Albumin (BSA, IgG and Protease free), 50% (v/v) Glycerol Product Description Prepared from tissue culture supernatant by Protein G affinity chromatograp Assay by Immunoelectrophoresis resulted in a single precipitin arc against A Rabbit Serum. Host Mouse Species Rabbit Specificity/Sensitivity Reactivity is observed against native Rabbit IgG by both Western blot and ELISA. Immunogen Rabbit IgG Product Application Details Applications Western Blot, ELISA, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation, In-situ Hybridization, Chromatin Immunoprecipitation (ChIP) Recommended Dilutions Western Blot 1:1000, ELISA, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation, In-situ Hybridization, Chromatin Immunoprecipitation (ChIP) Application Notes This secondary antibody may also be used for detection in immunoblotting assays that do not employ immunoprecipitation. It is provided as 1000X solutor oconserve reagent, we recommend incubating the blots from minigels in sealed bags (removing as much air as possible) with minimal volume (2-3 mused conservatively at 2.5mls/blot will yield enough reagent for 80 blots. Note that there are three key procedural considerations: 1. Protein A or G should not be used for the immunoprecipitation. Use of procedural considerations.	Preservative	No Preservative
Purity Protein G purified Buffer 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 0.1 mg/ml Bovine Serum Albumin (BSA, IgG and Protease free), 50% (v/v) Glycerol Product Description Prepared from tissue culture supernatant by Protein G affinity chromatograp Assay by Immunoelectrophoresis resulted in a single precipitin arc against A Rabbit Serum. Host Mouse Species Rabbit Specificity/Sensitivity Reactivity is observed against native Rabbit IgG by both Western blot and ELISA. Immunogen Rabbit IgG Product Application Details Applications Western Blot, ELISA, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation, In-situ Hybridization, Chromatin Immunoprecipitation (ChIP) Recommended Dilutions Western Blot 1:1000, ELISA, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation, In-situ Hybridization, Chromatin Immunoprecipitation (ChIP) Application Notes This secondary antibody may also be used for detection in immunoblotting assays that do not employ immunoprecipitation. It is provided as 1000X solutor To conserve reagent, we recommend incubating the blots from minigels in sealed bags (removing as much air as possible) with minimal volume (2-3 mused conservatively at 2.5mls/blot will yield enough reagent for 80 blots. Note that there are three key procedural considerations: 1. Protein A or G should not be used for the immunoprecipitation. Use of processors and the processor of the immunoprecipitation. Use of processor of the immunoprecipitation.	Isotype	IgG
Buffer 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 0.1 mg/ml Bovine Serum Albumin (BSA, IgG and Protease free), 50% (v/v) Glycerol Product Description Prepared from tissue culture supernatant by Protein G affinity chromatograp Assay by Immunoelectrophoresis resulted in a single precipitin arc against A Rabbit Serum. Host Mouse Reactivity Reactivity Reactivity is observed against native Rabbit IgG by both Western blot and ELISA. Immunogen Rabbit IgG Product Application Details Applications Western Blot, ELISA, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation, In-situ Hybridization, Chromatin Immunoprecipitation (ChIP) Recommended Dilutions Western Blot 1:1000, ELISA, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation, In-situ Hybridization, Chromatin Immunoprecipitation (ChIP) Application Notes This secondary antibody may also be used for detection in immunoblotting assays that do not employ immunoprecipitation. It is provided as 1000X soluto To conserve reagent, we recommend incubating the blots from minigels in sealed bags (removing as much air as possible) with minimal volume (2-3 mused conservatively at 2.5mls/blot will yield enough reagent for 80 blots. Note that there are three key procedural considerations: 1. Protein A or G should not be used for the immunoprecipitation. Use of procedural considerations.	Conjugate	HRP
Product Description Prepared from tissue culture supernatant by Protein G affinity chromatograp Assay by Immunoelectrophoresis resulted in a single precipitin arc against A Rabbit Serum. Host Mouse Species Rabbit Specificity/Sensitivity Reactivity is observed against native Rabbit IgG by both Western blot and ELISA. Immunogen Rabbit IgG Product Application Details Applications Western Blot, ELISA, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation, In-situ Hybridization, Chromatin Immunoprecipitation (ChIP) Recommended Dilutions Western Blot 1:1000, ELISA, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation, In-situ Hybridization, Chromatin Immunoprecipitation (ChIP) Application Notes This secondary antibody may also be used for detection in immunoblotting assays that do not employ immunoprecipitation. It is provided as 1000X soluto conserve reagent, we recommend incubating the blots from minigels in sealed bags (removing as much air as possible) with minimal volume (2-3 mused conservatively at 2.5mls/blot will yield enough reagent for 80 blots. Note that there are three key procedural considerations: 1. Protein A or G should not be used for the immunoprecipitation. Use of procedural considerations.	Purity	Protein G purified
Prepared from tissue culture supernatant by Protein G affinity chromatograp Assay by Immunoelectrophoresis resulted in a single precipitin arc against A Rabbit Serum. Host Mouse Species Rabbit Specificity/Sensitivity Reactivity is observed against native Rabbit IgG by both Western blot and ELISA. Immunogen Rabbit IgG Product Application Details Applications Western Blot, ELISA, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation, In-situ Hybridization, Chromatin Immunoprecipitation (ChIP) Recommended Dilutions Western Blot 1:1000, ELISA, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation, In-situ Hybridization, Chromatin Immunoprecipitation (ChIP) Application Notes This secondary antibody may also be used for detection in immunoblotting assays that do not employ immunoprecipitation. It is provided as 1000X soluto conserve reagent, we recommend incubating the blots from minigels in sealed bags (removing as much air as possible) with minimal volume (2-3 mused conservatively at 2.5mls/blot will yield enough reagent for 80 blots. Note that there are three key procedural considerations: 1. Protein A or G should not be used for the immunoprecipitation. Use of procedural considerations.	Buffer	
Assay by Immunoelectrophoresis resulted in a single precipitin arc against A Rabbit Serum. Mouse Species Rabbit Reactivity is observed against native Rabbit IgG by both Western blot and ELISA. Immunogen Rabbit IgG Product Application Details Applications Western Blot, ELISA, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation, In-situ Hybridization, Chromatin Immunoprecipitation (ChIP) Recommended Dilutions Western Blot 1:1000, ELISA, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation, In-situ Hybridization, Chromatin Immunoprecipitation (ChIP) Application Notes This secondary antibody may also be used for detection in immunoblotting assays that do not employ immunoprecipitation. It is provided as 1000X solu To conserve reagent, we recommend incubating the blots from minigels in sealed bags (removing as much air as possible) with minimal volume (2-3 m used conservatively at 2.5mls/blot will yield enough reagent for 80 blots. Note that there are three key procedural considerations: 1. Protein A or G should not be used for the immunoprecipitation. Use of procedural considerations.	Product Description	
Specificity/Sensitivity Reactivity is observed against native Rabbit IgG by both Western blot and ELISA. Immunogen Rabbit IgG Product Application Details Applications Western Blot, ELISA, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation, In-situ Hybridization, Chromatin Immunoprecipitation (ChIP) Recommended Dilutions Western Blot 1:1000, ELISA, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation, In-situ Hybridization, Chromatin Immunoprecipitation (ChIP) Application Notes This secondary antibody may also be used for detection in immunoblotting assays that do not employ immunoprecipitation. It is provided as 1000X soluto conserve reagent, we recommend incubating the blots from minigels in sealed bags (removing as much air as possible) with minimal volume (2-3 mused conservatively at 2.5mls/blot will yield enough reagent for 80 blots. Note that there are three key procedural considerations: 1. Protein A or G should not be used for the immunoprecipitation. Use of procedural considerations.	Description	Prepared from tissue culture supernatant by Protein G affinity chromatography. Assay by Immunoelectrophoresis resulted in a single precipitin arc against Anti-Rabbit Serum.
Reactivity is observed against native Rabbit IgG by both Western blot and ELISA.	Host	Mouse
Immunogen	Species	Rabbit
Product Application Details Applications Western Blot, ELISA, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation, In-situ Hybridization, Chromatin Immunoprecipitation (ChIP) Recommended Dilutions Western Blot 1:1000, ELISA, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation, In-situ Hybridization, Chromatin Immunoprecipitation (ChIP) Application Notes This secondary antibody may also be used for detection in immunoblotting assays that do not employ immunoprecipitation. It is provided as 1000X solu To conserve reagent, we recommend incubating the blots from minigels in sealed bags (removing as much air as possible) with minimal volume (2-3 mused conservatively at 2.5mls/blot will yield enough reagent for 80 blots. Note that there are three key procedural considerations: 1. Protein A or G should not be used for the immunoprecipitation. Use of procedural considerations.	Specificity/Sensitivity	
Applications Western Blot, ELISA, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation, In-situ Hybridization, Chromatin Immunoprecipitation (ChIP) Western Blot 1:1000, ELISA, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation, In-situ Hybridization, Chromatin Immunoprecipitation (ChIP) Application Notes This secondary antibody may also be used for detection in immunoblotting assays that do not employ immunoprecipitation. It is provided as 1000X solu To conserve reagent, we recommend incubating the blots from minigels in sealed bags (removing as much air as possible) with minimal volume (2-3 m used conservatively at 2.5mls/blot will yield enough reagent for 80 blots. Note that there are three key procedural considerations: 1. Protein A or G should not be used for the immunoprecipitation. Use of procedural considerations.	Immunogen	Rabbit IgG
Immunoprecipitation, In-situ Hybridization, Chromatin Immunoprecipitation (ChIP) Western Blot 1:1000, ELISA, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation, In-situ Hybridization, Chromatin Immunoprecipitation (ChIP) Application Notes This secondary antibody may also be used for detection in immunoblotting assays that do not employ immunoprecipitation. It is provided as 1000X soluto To conserve reagent, we recommend incubating the blots from minigels in sealed bags (removing as much air as possible) with minimal volume (2-3 mused conservatively at 2.5mls/blot will yield enough reagent for 80 blots. Note that there are three key procedural considerations: 1. Protein A or G should not be used for the immunoprecipitation. Use of procedural considerations.	Product Application Details	
Immunoprecipitation, In-situ Hybridization, Chromatin Immunoprecipitation (ChIP) Application Notes This secondary antibody may also be used for detection in immunoblotting assays that do not employ immunoprecipitation. It is provided as 1000X soluto conserve reagent, we recommend incubating the blots from minigels in sealed bags (removing as much air as possible) with minimal volume (2-3 mused conservatively at 2.5mls/blot will yield enough reagent for 80 blots. Note that there are three key procedural considerations: 1. Protein A or G should not be used for the immunoprecipitation. Use of procedural considerations.	Applications	Immunoprecipitation, In-situ Hybridization, Chromatin Immunoprecipitation
assays that do not employ immunoprecipitation. It is provided as 1000X solute To conserve reagent, we recommend incubating the blots from minigels in sealed bags (removing as much air as possible) with minimal volume (2-3 mused conservatively at 2.5mls/blot will yield enough reagent for 80 blots. Note that there are three key procedural considerations: 1. Protein A or G should not be used for the immunoprecipitation. Use of procedural considerations.	Recommended Dilutions	Immunoprecipitation, In-situ Hybridization, Chromatin Immunoprecipitation
immunoprecipitation, Anti-Rat IgG beads, or Anti-Rabbit IgG beads should be used for rat or rabbit immunoprecipitating antibodies, respectively. 2. Immunoprecipitate should be completely reduced. 3. BLOTTO/Milk should be used as the blocking protein for the immunoblot. recommended dilutions for listed applications are intended as an initial	Application Notes	assays that do not employ immunoprecipitation. It is provided as 1000X solution. To conserve reagent, we recommend incubating the blots from minigels in sealed bags (removing as much air as possible) with minimal volume (2-3 mls). If used conservatively at 2.5mls/blot will yield enough reagent for 80 blots. Note that there are three key procedural considerations: 1. Protein A or G should not be used for the immunoprecipitation. Use of protein A or G beads with the rabbit Pure-Blot will result in contaminating bands. For immunoprecipitation, Anti-Rat IgG beads, or Anti-Rabbit IgG beads should be used for rat or rabbit immunoprecipitating antibodies, respectively. 2. Immunoprecipitate should be completely reduced. 3. BLOTTO/Milk should be used as the blocking protein for the immunoblot. All recommended dilutions for listed applications are intended as an initial recommendation, specific conditions for each protein and antibody combination



Images

Western Blot: Mouse Pure-Blot anti-Rabbit IgG (H+L) Secondary Antibody (eB182) [HRP] [NBP3-11668] - Jurkat cell lysate (0.5 ml of 1x10e7 cells/ml) was incubated with rabbit anti-human Stat1 and immunoprecipitated using Protein G, Protein A and Anti-Rabbit Ig IP Beads. Precipitate from 5x10e5 cells was subjected to electrophoresis, transferred to a PVDF membrane, and Western blotted with anti-Stat1 using Mouse Pure-Blot anti-Rabbit IgG Secondary antibody (eB182) [HRP].





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Secondary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-11668

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

