## **Product Datasheet**

### NPM1 Antibody (NPM1/7072R) [Alexa Fluor® 647] NBP3-14019AF647

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com

G

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-14019AF647

Updated 10/26/2023 v.20.1

# Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-14019AF647



#### NBP3-14019AF647

NPM1 Antibody (NPM1/7072R) [Alexa Fluor® 647]

Unit Size   0.1 ml     Concentration   Please see the vial label for concentration. If unlisted please contact technical services.     Storage   Store at 4C in the dark.     Clonality   Monoclonal     Clone   NPM1/7072R     Preservative   0.05% Sodium Azide     Isotype   IgG Kappa     Conjugate   Alexa Fluor 647     Purity   Protein A or G purified     Buffer   50mM Sodium Borate     Product Description   Ka69     Gene ID   4869     Gene Symbol   NPM1     Species   Human     Marker   Acute Myeloid Leukemia Marker     Specificity/Sensitivity   Recognizes a 33kDa glycoprotein, identified as NPM1 (NPM). It is predominantly localized in the nucleus of cells in most tissues. NPM is involved in ribosomal assembly and rRNA transport. It is an abundant protein that is highly phosphorylated by Cdc2 kinase during mitosis. This phosphoprolet in moves between the nucleus of cells in cost tissues. NPM is involved in ribosomal assembly and rRNA transport. It is an abundant protein that is highly phosphorylated by Cdc2 kinase during mitosis. This phosphoprotein moves between the nucleus of cells in cost tissues. NPM is involved in ribosomal associated with acute myeloid leukemia. The antibody may be a useful aid for classification of acute myeloid leukemia. The antibody may be a useful aid for cla	, , , , , , , , , , , , , , , , , , ,	•
ConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneNPM1/7072RPreservative0.05% Sodium AzideIsotypeIgG KappaConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionRabbitGene ID4869Gene ID4869Gene SymbolNPM1SpeciesHumanMarkerAcute Myeloid Leukemia MarkerSpecificity/SensitivityRecognizes a 33kDa glycoprotein, identified as NPM1 (NPM). It is predominantly localized in the nucleus of cells in most norther in robosphorylated by Cdc2 kinase during mitosis. This phosphorprotein moves between the nucleus and the cytoplasm. It is thought to be involved in several processes including regulation of the ARF/p53 pathway. A number of genes are fusion partners, in particular the anaplastic lymphoma kinase gene on chromosome 2. Mutations in exon 12 affecting the C-terminus of the protein are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with an aberrant cytoplasmic location. Mutations in this gene are assoc	Product Information	
services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneNPM1/7072RPreservative0.05% Sodium AzideIsotypeIgG KappaConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionRabbitHostRabbitGene ID4869Gene SymbolNPM1SpeciesHumanMarkerAcute Myeloid Leukemia MarkerSpecificity/SensitivityRecognizes a 33kDa glycoprotein, identified as NPM1 (NPM). It is predominantly localized in the nucleus of cells in most tissues. NPM is involved in ribosomal assembly and rRNA transport. It is an abundant protein that is highly phosphorylated by Cdc2 kinase during mitosis. This phosphoprotein moves between the nucleus and the cytplasm. It is though to be protein are associated with an aberrant cytoplasm. It is though to be protein are associated with acute myeloid leukemia. The antibody may be a useful aid for classification of acute myeloid leukemia. The antibody may be a useful aid for classification of acute myeloid leukemia.ImmunogenRecombinant fragment of human NPM1 protein (exact sequence is proprietary)	Unit Size	0.1 ml
ClonalityMonoclonalCloneNPM1/7072RPreservative0.05% Sodium AzideIsotypeIgG KappaConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionRabbitHostRabbitGene ID4869Gene SymbolNPM1SpeciesHumanMarkerAcute Myeloid Leukemia MarkerSpecificity/SensitivityRecognizes a 33kDa glycoprotein, identified as NPM1 (NPM). It is predominantly localized in the nucleus of cells in most tissues. NPM is involved in ribosomal assembly and rRNA transport. It is an abundant protein that is highly phosphorylated by Cd2 kinase during mitosis. This phosphoprotein moves between the nucleus and the cytoplasm. It is thought to be involved in several processes including regulation of the ARF/p53 pathway. A number of genes are fusion partners, in particular the anaplastic lymphoma kinase gene on chromosome 2. Mutations in exon 12 affecting the C-terminus of the protein are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with acute myeloid leukemia.ImmunogenRecombinant fragment o	Concentration	
CloneNPM1/7072RPreservative0.05% Sodium AzideIsotypeIgG KappaConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionKabbitHostRabbitGene ID4869Gene SymbolNPM1SpeciesHumanMarkerAcute Myeloid Leukemia MarkerSpecificity/SensitivityRecognizes a 33kDa glycoprotein, identified as NPM1 (NPM). It is predominantly localized in the nucleus of cells in most tissues. NPM is involved in ribosomal assembly and rRNA transport. It is an abundant protein that is highly phosphorylated by Cdc2 kinase during mitosis. This phosphoprotein are associated with an aberrant cytoplasm. It is thought to be involved in several processes including regulation of the ARF/pS3 pathway. A number of genes are fusion partners, in particular the anaplastic location. Mutations in this gene are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with an aberrant cytoplasmic location. Mu	Storage	Store at 4C in the dark.
Preservative 0.05% Sodium Azide   Isotype IgG Kappa   Conjugate Alexa Fluor 647   Purity Protein A or G purified   Buffer 50mM Sodium Borate   Product Description Host   Host Rabbit   Gene ID 4869   Gene Symbol NPM1   Species Human   Marker Acute Myeloid Leukemia Marker   Specificity/Sensitivity Recognizes a 33kDa glycoprotein, identified as NPM1 (NPM). It is predominantly localized in the nucleus of cells in most tissues. NPM is involved in ribosomal assembly and rRNA transport. It is an abundant protein that is highly phosphorylated by Cdc2 kinase during mitosis. This phosphoprotein moves between the nucleus and the cytoplasm. It is thought to be involved in several processes including regulation of the ARF/pS3 pathway. A number of genes are fusion partners, in particular the anaplastic lymphoma kinase gene on chromosome 2. Mutations in exon 12 affecting the C-terminus of the protein are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with acute myeloid leukemia. </th <th>Clonality</th> <th>Monoclonal</th>	Clonality	Monoclonal
IsotypeIgG KappaConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID4869Gene SymbolNPM1SpeciesHumanMarkerAcute Myeloid Leukemia MarkerSpecificity/SensitivityRecognizes a 33kDa glycoprotein, identified as NPM1 (NPM). It is predominantly localized in the nucleus of cells in most tissues. NPM is involved in ribosomal assembly and rRNA transport. It is an abundant protein that is highly phosphorylated by Cdc2 kinase during mitosis. This phosphorpter moves between the nucleus and the cytoplasm. It is thought to be involved in several processes in cluding regulation of the ARF/PS3 pathway. A number of genes are fusion partners, in particular the anaplastic lymphoma kinase gene on chromosome 2. Mutations in exon 12 affecting the C-terminus of the protein are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with a cute myeloid leukemia.ImmunogenRecombinant fragment of human NPM1 protein (exact sequence is proprietary)	Clone	NPM1/7072R
ConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID4869Gene SymbolNPM1SpeciesHumanMarkerAcute Myeloid Leukemia MarkerSpecificity/SensitivityRecognizes a 33kDa glycoprotein, identified as NPM1 (NPM). It is predominantly localized in the nucleus of cells in most tissues. NPM is involved in ribosomal assembly and rRNA transport. It is an abundant protein that is highly phosphorylated by Cdc2 kinase during mitosis. This phosphoprotein moves between the nucleus and the cytoplasm. It is thought to be involved in several processes including regulation of the ARF/p53 pathway. A number of genes are fusion partners, in particular the anaplastic lymphoma kinase gene on chromosome 2. Mutations in exon 12 affecting the C-terminus of the protein are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with acute myeloid leukemia. The antibody may be a useful aid for classification of acute myeloid leukemia.ImmunogenRecombinant fragment of human NPM1 protein (exact sequence is proprietary)	Preservative	0.05% Sodium Azide
PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID4869Gene SymbolNPM1SpeciesHumanMarkerAcute Myeloid Leukemia MarkerSpecificity/SensitivityRecognizes a 33kDa glycoprotein, identified as NPM1 (NPM). It is predominantly localized in the nucleus of cells in most tissues. NPM is involved in ribosomal assembly and rRNA transport. It is an abundant protein that is highly phosphorylated by Cdc2 kinase during mitosis. This phosphoprotein moves between the nucleus and the cytoplasm. It is thought to be involved in several processes including regulation of the ARF/p53 pathway. A number of genes are fusion partners, in particular the anaplastic lymphoma kinase gene on chromosome 2. Mutations in exon 12 affecting the C-terminus of the protein are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with acute myeloid leukemia. The antibody may be a useful aid for classification of acute myeloid leukemia.ImmunogenRecombinant fragment of human NPM1 protein (exact sequence is proprietary)	Isotype	IgG Kappa
Buffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID4869Gene SymbolNPM1SpeciesHumanMarkerAcute Myeloid Leukemia MarkerSpecificity/SensitivityRecognizes a 33kDa glycoprotein, identified as NPM1 (NPM). It is predominantly localized in the nucleus of cells in most tissues. NPM is involved in ribosomal assembly and rRNA transport. It is an abundant protein that is highly phosphorylated by Cdc2 kinase during mitosis. This phosphoprotein moves between the nucleus and the cytoplasm. It is thought to be involved in several processes including regulation of the ARF/p53 pathway. A number of genes are fusion partners, in particular the anaplastic lymphoma kinase gene on chromosome 2. Mutations in exon 12 affecting the C-terminus of the protein are associated with a cute myeloid leukemia. The antibody may be a useful aid for classification of acute myeloid leukemia.ImmunogenRecombinant fragment of human NPM1 protein (exact sequence is proprietary)	Conjugate	Alexa Fluor 647
Product DescriptionHostRabbitGene ID4869Gene SymbolNPM1SpeciesHumanMarkerAcute Myeloid Leukemia MarkerSpecificity/SensitivityRecognizes a 33kDa glycoprotein, identified as NPM1 (NPM). It is predominantly localized in the nucleus of cells in most tissues. NPM is involved in ribosomal assembly and rRNA transport. It is an abundant protein that is highly phosphorylated by Cdc2 kinase during mitosis. This phosphoprotein moves between the nucleus and the cytoplasm. It is thought to be involved in several processes including regulation of the ARF/p53 pathway. A number of genes are fusion partners, in particular the anaplastic lymphoma kinase gene on chromosome 2. Mutations in exon 12 affecting the C-terminus of the protein are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with acute myeloid leukemia. The antibody may be a useful aid for classification of acute myeloid leukemia.ImmunogenRecombinant fragment of human NPM1 protein (exact sequence is proprietary)	Purity	Protein A or G purified
HostRabbitGene ID4869Gene SymbolNPM1SpeciesHumanMarkerAcute Myeloid Leukemia MarkerSpecificity/SensitivityRecognizes a 33kDa glycoprotein, identified as NPM1 (NPM). It is predominantly localized in the nucleus of cells in most tissues. NPM is involved in ribosomal assembly and rRNA transport. It is an abundant protein that is highly phosphorylated by Cdc2 kinase during mitosis. This phosphoprotein moves between the nucleus and the cytoplasm. It is thought to be involved in several processes including regulation of the ARF/p53 pathway. A number of genes are fusion partners, in particular the anaplastic lymphoma kinase gene on chromosome 2. Mutations in exon 12 affecting the C-terminus of the protein are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with a cute myeloid leukemia. The antibody may be a useful aid for classification of acute myeloid leukemia.ImmunogenRecombinant fragment of human NPM1 protein (exact sequence is proprietary)	Buffer	50mM Sodium Borate
Gene ID4869Gene SymbolNPM1SpeciesHumanMarkerAcute Myeloid Leukemia MarkerSpecificity/SensitivityRecognizes a 33kDa glycoprotein, identified as NPM1 (NPM). It is predominantly localized in the nucleus of cells in most tissues. NPM is involved in ribosomal assembly and rRNA transport. It is an abundant protein that is highly phosphorylated by Cdc2 kinase during mitosis. This phosphoprotein moves between the nucleus and the cytoplasm. It is thought to be involved in several processes including regulation of the ARF/p53 pathway. A number of genes are fusion partners, in particular the anaplastic lymphoma kinase gene on chromosome 2. Mutations in exon 12 affecting the C-terminus of the protein are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with acute myeloid leukemia. The antibody may be a useful aid for classification of acute myeloid leukemia.ImmunogenRecombinant fragment of human NPM1 protein (exact sequence is proprietary)	Product Description	
Gene SymbolNPM1SpeciesHumanMarkerAcute Myeloid Leukemia MarkerSpecificity/SensitivityRecognizes a 33kDa glycoprotein, identified as NPM1 (NPM). It is predominantly localized in the nucleus of cells in most tissues. NPM is involved in ribosomal assembly and rRNA transport. It is an abundant protein that is highly phosphorylated by Cdc2 kinase during mitosis. This phosphoprotein moves between the nucleus and the cytoplasm. It is thought to be involved in several processes including regulation of the ARF/p53 pathway. A number of genes are fusion partners, in particular the anaplastic lymphoma kinase gene on chromosome 2. Mutations in exon 12 affecting the C-terminus of the protein are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with acute myeloid leukemia. The antibody may be a useful aid for classification of acute myeloid leukemia.ImmunogenRecombinant fragment of human NPM1 protein (exact sequence is proprietary)	Host	Rabbit
SpeciesHumanMarkerAcute Myeloid Leukemia MarkerSpecificity/SensitivityRecognizes a 33kDa glycoprotein, identified as NPM1 (NPM). It is predominantly localized in the nucleus of cells in most tissues. NPM is involved in ribosomal assembly and rRNA transport. It is an abundant protein that is highly phosphorylated by Cdc2 kinase during mitosis. This phosphoprotein moves between the nucleus and the cytoplasm. It is thought to be involved in several processes including regulation of the ARF/p53 pathway. A number of genes are fusion partners, in particular the anaplastic lymphoma kinase gene on chromosome 2. Mutations in exon 12 affecting the C-terminus of the protein are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with acute myeloid leukemia. The antibody may be a useful aid for classification of acute myeloid leukemia.ImmunogenRecombinant fragment of human NPM1 protein (exact sequence is proprietary)	Gene ID	4869
MarkerAcute Myeloid Leukemia MarkerSpecificity/SensitivityRecognizes a 33kDa glycoprotein, identified as NPM1 (NPM). It is predominantly localized in the nucleus of cells in most tissues. NPM is involved in ribosomal assembly and rRNA transport. It is an abundant protein that is highly phosphorylated by Cdc2 kinase during mitosis. This phosphoprotein moves between the nucleus and the cytoplasm. It is thought to be involved in several processes including regulation of the ARF/p53 pathway. A number of genes are fusion partners, in particular the anaplastic lymphoma kinase gene on chromosome 2. Mutations in exon 12 affecting the C-terminus of the protein are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with acute myeloid leukemia. The antibody may be a useful aid for classification of acute myeloid leukemia.ImmunogenRecombinant fragment of human NPM1 protein (exact sequence is proprietary)	Gene Symbol	NPM1
Specificity/SensitivityRecognizes a 33kDa glycoprotein, identified as NPM1 (NPM). It is predominantly localized in the nucleus of cells in most tissues. NPM is involved in ribosomal assembly and rRNA transport. It is an abundant protein that is highly phosphorylated by Cdc2 kinase during mitosis. This phosphoprotein moves between the nucleus and the cytoplasm. It is thought to be involved in several processes including regulation of the ARF/p53 pathway. A number of genes are fusion partners, in particular the anaplastic lymphoma kinase gene on chromosome 2. Mutations in exon 12 affecting the C-terminus of the protein are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with acute myeloid leukemia. The antibody may be a useful aid for classification of acute myeloid leukemia.ImmunogenRecombinant fragment of human NPM1 protein (exact sequence is proprietary)	Species	Human
Iocalized in the nucleus of cells in most tissues. NPM is involved in ribosomal assembly and rRNA transport. It is an abundant protein that is highly phosphorylated by Cdc2 kinase during mitosis. This phosphoprotein moves between the nucleus and the cytoplasm. It is thought to be involved in several processes including regulation of the ARF/p53 pathway. A number of genes are fusion partners, in particular the anaplastic lymphoma kinase gene on chromosome 2. Mutations in exon 12 affecting the C-terminus of the protein are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with acute myeloid leukemia. The antibody may be a useful aid for classification of acute myeloid leukemia.ImmunogenRecombinant fragment of human NPM1 protein (exact sequence is proprietary)	Marker	Acute Myeloid Leukemia Marker
	Specificity/Sensitivity	localized in the nucleus of cells in most tissues. NPM is involved in ribosomal assembly and rRNA transport. It is an abundant protein that is highly phosphorylated by Cdc2 kinase during mitosis. This phosphoprotein moves between the nucleus and the cytoplasm. It is thought to be involved in several processes including regulation of the ARF/p53 pathway. A number of genes are fusion partners, in particular the anaplastic lymphoma kinase gene on chromosome 2. Mutations in exon 12 affecting the C-terminus of the protein are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with acute myeloid leukemia. The antibody may be a useful aid for
	Immunogen	



	Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.
Product Application Details	
Applications	Immunohistochemistry-Paraffin
<b>Recommended Dilutions</b>	Immunohistochemistry-Paraffin
Application Notes	Optimal dilution of this antibody should be experimentally determined.

Notes





#### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 novus@novusbio.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com Technical Support: technical@novusbio.com Orders: orders@novusbio.com General: novus@novusbio.com

#### Products Related to NBP3-14019AF647

NB110-61646PEP	NPM1 Antibody Blocking Peptide
NBL1-13748	NPM1 Overexpression Lysate
NB200-106	p19ARF/CDKN2A Antibody - BSA Free
NB200-103	p53 Antibody (PAb 240) - BSA Free

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-14019AF647

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

