Product Datasheet

Histone H2AE [ac Lys9] Antibody (3H5) - BSA Free NBP3-15226

Unit Size: 100 ul

Store at -20 to -70C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-15226

Updated 7/30/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP3-15226



NBP3-15226

Histone H2AE [ac Lys9] Antibody (3H5) - BSA Free

(3H5) - BSA Free				
100 ul				
Please see the vial label for concentration. If unlisted please contact technical services.				
Store at -20 to -70C. Avoid freeze-thaw cycles.				
Monoclonal				
3H5				
0.02% Sodium Azide				
IgG				
Affinity purified				
PBS, pH 7.4, 150mM NaCl, and 50% glycerol				
Product Description				
Rabbit				
3012				
H2AC8				
Human				
A synthesized peptide. UniProt ID (P04908).				
Product Application Details				
ELISA, Immunocytochemistry/ Immunofluorescence				
ELISA, Immunocytochemistry/ Immunofluorescence 1:50-1:500				


1	7	
۲	J	P

Images

Immunocytochemistry/Immunofluorescence: Histone H2AE [ac Lys9] Antibody (3H5) [NBP3-15226] - Immunocytochemistry analysis of Histone H2AE antibody diluted at 1:100 and staining in Hela cells. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Immunocytochemistry/Immunofluorescence: Histone H2AE [ac Lys9] Antibody (3H5) [NBP3-15226] - Immunofluorescence staining of Hela cells (treated by 15mM sodium butyrate for 30 minutes) with at Histone H2AE antibody 1:56, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4C.The secondary antibody was Alexa Fluor 488-congugated Goat Anti-Rabbit IgG (H+L).





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP3-15226

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control
NBL1-11567 H4/j Overexpression Lysate

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-15226

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

