Product Datasheet

PAK4+5+6 [p Ser474, p Ser560, p Ser602] Antibody (2D1) BSA Free NBP3-15234

Unit Size: 100 ul

Store at -20 to -70C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-15234

Updated 7/30/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP3-15234



NBP3-15234


PAK4+5+6 [p Ser474, p Ser560, p Ser602] Antibody (2D1) - BSA Free

| PAR4+5+6 [p Sei474, p Sei560, p Sei602] Allibody (2D1) - BSA Fiee | |
|---|---|
| Product Information | |
| Unit Size | 100 ul |
| Concentration | Please see the vial label for concentration. If unlisted please contact technical services. |
| Storage | Store at -20 to -70C. Avoid freeze-thaw cycles. |
| Clonality | Monoclonal |
| Clone | 2D1 |
| Preservative | 0.02% Sodium Azide |
| Isotype | IgG |
| Purity | Affinity purified |
| Buffer | PBS, pH 7.4, 150mM NaCl, and 50% glycerol |
| Product Description | |
| Host | Rabbit |
| Gene ID | 10298 |
| Gene Symbol | PAK4 |
| Species | Human |
| Immunogen | A synthesized peptide derived from human Phospho-PAK4+5+6 (S474/S560/S602). UniProt (O96013/Q9P286/Q9NQU5). |
| Product Application Details | |
| Applications | Western Blot, ELISA, Immunocytochemistry/ Immunofluorescence |
| Recommended Dilutions | Western Blot 1:500-1:5000, ELISA, Immunocytochemistry/ Immunofluorescence 1:20-1:200 |




Images

Western Blot: PAK4+5+6 [p Ser474, p Ser560, p Ser602] Antibody (2D1) [NBP3-15234] - Positive WB detected in 293 whole cell lysate (treated with Calyculin A or not). All lanes PAK4+5+6 antibody at 2.15ug/ml. Secondary: Goat polyclonal to rabbit IgG at 1/50000 dilution. Predicted band size: 64,75,80 KDa. Observed band size: 75,80 Kda.



Immunocytochemistry/Immunofluorescence: PAK4+5+6 [p Ser474, p Ser560, p Ser602] Antibody (2D1) [NBP3-15234] - Immunofluorescence staining of HepG2 cells (treated with 50mM Calyculin A for 30 minutes) with at PAK4+5+6 antibody 1:100, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4C. The secondary antibody was Alexa Fluor 488-congugated Goat Anti-Rabbit IgG (H+L).





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP3-15234

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

294-HG-005 HGF [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-15234

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

