

Product Datasheet

MED4 Antibody (8T5C9)

NBP3-15287-100ul

Unit Size: 100 ul

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP3-15287

Updated 7/31/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP3-15287



NBP3-15287-100ul

MED4 Antibody (8T5C9)

Product Information

Unit Size	100 ul
Concentration	Please see the vial label for concentration. If unlisted please contact technical services.
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	8T5C9
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Affinity purified
Buffer	PBS (pH 7.3), 50% glycerol, 0.05% BSA

Product Description


Host	Rabbit
Gene ID	29079
Gene Symbol	MED4
Species	Human, Mouse, Rat
Immunogen	Recombinant fusion protein containing a sequence corresponding to amino acids 150-250 of human MED4 (Q9NPJ6). KYAHRISASNAVCAPLTWVPGDPRRPYPTDLEMRSGLLGQMNNPSTNGVNGH LPGDALAAGRLPDVLAPQYPWQSNDSMNMLPPNHSSDFLLEPPGHNKE

Product Application Details


Applications	Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry
Recommended Dilutions	Western Blot 1:500 - 1:1000, Immunohistochemistry 1:50 - 1:200, Immunohistochemistry-Paraffin

Images

Western Blot: MED4 Antibody (8T5C9) [NBP3-15287] - Western blot analysis of extracts of various cell lines, using MED4 antibody (NBP3-15287) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 5s.




Immunohistochemistry-Paraffin: MED4 Antibody (8T5C9) [NBP3-15287]
 - Immunohistochemistry of paraffin-embedded rat liver using MED4 Rabbit mAb (NBP3-15287) at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.




Immunohistochemistry-Paraffin: MED4 Antibody (8T5C9) [NBP3-15287]
 - Immunohistochemistry of paraffin-embedded human colon carcinoma using MED4 Rabbit mAb (NBP3-15287) at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.




Immunohistochemistry-Paraffin: MED4 Antibody (8T5C9) [NBP3-15287]
 - Immunohistochemistry of paraffin-embedded mouse lung using MED4 Rabbit mAb (NBP3-15287) at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.




Immunohistochemistry: MED4 Antibody (8T5C9) [NBP3-15287] -
 Immunohistochemistry analysis of MED4 in paraffin-embedded rat colon tissue using MED4 Rabbit mAb at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry: MED4 Antibody (8T5C9) [NBP3-15287] - Immunohistochemistry analysis of MED4 in paraffin-embedded human breast tissue using MED4 Rabbit mAb at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry: MED4 Antibody (8T5C9) [NBP3-15287] - Immunohistochemistry analysis of MED4 in paraffin-embedded human colon tissue using MED4 Rabbit mAb at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry: MED4 Antibody (8T5C9) [NBP3-15287] - Immunohistochemistry analysis of MED4 in paraffin-embedded mouse colon tissue using MED4 Rabbit mAb at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry: MED4 Antibody (8T5C9) [NBP3-15287] - Immunohistochemistry analysis of MED4 in paraffin-embedded human esophagus tissue using MED4 Rabbit mAb at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry: MED4 Antibody (8T5C9) [NBP3-15287] - Immunohistochemistry analysis of MED4 in paraffin-embedded human breast cancer tissue using MED4 Rabbit mAb at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP3-15287-100ul

HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control
NBP1-44479	Recombinant Human MED4 His Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-15287

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

