# **Product Datasheet**

# SUMO Activating Enzyme E1 (SAE1) Antibody (0L2N2) NBP3-15317-100ul

Unit Size: 100 ul

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-15317

Updated 7/31/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP3-15317



# NBP3-15317-100ul

SUMO Activating Enzyme E1 (SAE1) Antibody (0L2N2)


SUMO Activating Enzyme E1 (SAE1) Antibody (0L2N2)	
100 ul	
Please see the vial label for concentration. If unlisted please contact technical services.	
Store at -20C. Avoid freeze-thaw cycles.	
Monoclonal	
0L2N2	
0.02% Sodium Azide	
IgG	
Affinity purified	
PBS (pH 7.3), 50% glycerol, 0.05% BSA	
Product Description	
Rabbit	
10055	
SAE1	
Human, Mouse, Rat	
A synthetic peptide corresponding to a sequence within amino acids 200-300 of human SUMO Activating Enzyme E1 (SAE1) (Q9UBE0).  DSSETTMVKKKVVFCPVKEALEVDWSSEKAKAALKRTTSDYFLLQVLLKFRTD KGRDPSSDTYEEDSELLLQIRNDVLDSLGISPDLLPEDFVRYCFSEMA	
Western Blot	

	1.10.12.002.1.22022223
<b>Product Application Details</b>	
Applications	Western Blot
Recommended Dilutions	Western Blot 1:500 - 1:2000




#### **Images**

Western Blot: SUMO Activating Enzyme E1 (SAE1) Antibody (0L2N2) [NBP3-15317] - Western blot analysis of extracts of various cell lines, using SUMO Activating Enzyme E1 (SAE1) Rabbit mAb (NBP3-15317) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 1s.



Western Blot: SUMO Activating Enzyme E1 (SAE1) Antibody (0L2N2) [NBP3-15317] - Western blot analysis of extracts of various cell lines, using SUMO Activating Enzyme E1 (SAE1) Rabbit mAb (NBP3-15317) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 5s.





## **Novus Biologicals USA**

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

## Products Related to NBP3-15317-100ul

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

NBP2-13272PEP SUMO Activating Enzyme E1 (SAE1) Recombinant Protein Antigen

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-15317

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

