

Product Datasheet

HO-1/HMOX1/HSP32 Antibody (7H7W10) NBP3-15640-100ul

Unit Size: 100 ul

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP3-15640

Updated 7/31/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP3-15640



NBP3-15640-100ul

HO-1/HMOX1/HSP32 Antibody (7H7W10)


Product Information	
Unit Size	100 ul
Concentration	Please see the vial label for concentration. If unlisted please contact technical services.
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	7H7W10
Preservative	0.05% Proclin 300
Isotype	IgG
Purity	Affinity purified
Buffer	PBS (pH 7.3), 50% glycerol, 0.05% BSA
Target Molecular Weight	33 kDa

Product Description	
Host	Rabbit
Gene ID	3162
Gene Symbol	HMOX1
Species	Human, Mouse, Rat
Immunogen	Recombinant fusion protein containing a sequence corresponding to amino acids 1-265 of human Heme Oxygenase 1 (HO-1/HMOX1) (NP_002124.1). MERPQPDSMPQDLSEALKEATKEVHTQAENAEFMRNFQKGQVTRDGFKLVM ASLYHIYVALEEEIERNKESPVFAPVYFPEELHRKAALEQDLAFWYGPRWQEV PYTPAMQRYVKRLHEVGRTEPELLVAHAYTRYLGDLGGQVLKKIAQKALDLP SSGEGLAFFTFPNIASATKFKQLYRSRMNSLEMPAVRQRVIEEAKTAFLLNIQL FEELQELLTHDTKDQSPSRAPGLRQRASNKVQDSAPVETPRGKPLNTRSQA


Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, ELISA, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Knockout Validated
Recommended Dilutions	Western Blot 1:2000 - 1:10000, ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements., Immunohistochemistry 1:200 - 1:800, Immunocytochemistry/Immunofluorescence 1:50 - 1:200, Immunohistochemistry-Paraffin 1:200 - 1:800, Knockout Validated

Images


analysis of extracts of Raw264.7 cells, using Heme Oxygenase 1 (HO-1/HMOX1) antibody (A19062) at 1:10000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 3s.




HO-1/HMOX1/HSP32 Antibody (3L2U4) [NBP3-15640] - Analysis of extracts of HepG2 cells, using Heme Oxygenase 1 (HO-1/HMOX1) antibody at 1:10000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 180s.




Western Blot: Rabbit HO-1/HMOX1/HSP32 mAb (7H7W10) [NBP3-15640] - Analysis of extracts from wild type(WT) and Heme Oxygenase 1 (HO-1/HMOX1) knockout (KO) HeLa cells, using Heme Oxygenase 1 (HO-1/HMOX1) antibody at 1:10000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 180s.




Immunohistochemistry: HO-1/HMOX1/HSP32 Antibody (7H7W10) [NBP3-15640] - Immunohistochemistry analysis of paraffin-embedded Human tonsil tissue using [KD Validated] HO-1/HMOX1/HSP32 Rabbit mAb at a dilution of 1:500 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.




Immunohistochemistry: HO-1/HMOX1/HSP32 Antibody (7H7W10) [NBP3-15640] - Immunohistochemistry analysis of paraffin-embedded Mouse spleen tissue using [KD Validated] HO-1/HMOX1/HSP32 Rabbit mAb at a dilution of 1:500 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry: HO-1/HMOX1/HSP32 Antibody (7H7W10) [NBP3-15640] - Immunohistochemistry analysis of paraffin-embedded Human liver tissue using [KD Validated] HO-1/HMOX1/HSP32 Rabbit mAb at a dilution of 1:500 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunocytochemistry/ Immunofluorescence: HO-1/HMOX1/HSP32 Antibody (7H7W10) [NBP3-15640] - Immunofluorescence analysis of paraffin-embedded human spleen using [KD Validated] HO-1/HMOX1/HSP32 Rabbit mAb at dilution of 1:50 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP3-15640-100ul

HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control
NBP1-77459PEP	HO-1/HMOX1/HSP32 Antibody Blocking Peptide

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-15640

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

