Product Datasheet

PRP19 Antibody (4M3Z1) NBP3-16853-100ul

Unit Size: 100 ul

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-16853

Updated 7/31/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-16853



NBP3-16853-100ul

PRP19 Antibody (4M3Z1)

,	
Product Information	
Unit Size	100 ul
Concentration	Please see the vial label for concentration. If unlisted please contact technical services.
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	4M3Z1
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Affinity purified
Buffer	PBS (pH 7.3), 50% glycerol, 0.05% BSA
Product Description	
Host	Rabbit
Gene ID	27339
Gene Symbol	PRPF19
Species	Human, Mouse, Rat
Immunogen	Recombinant fusion protein containing a sequence corresponding to amino acids 400-504 of human PRPF19(Q9UMS4). FSENGYYLATAADDSSVKLWDLRKLKNFKTLQLDNNFEVKSLIFDQSGTYLALG GTDVQIYICKQWTEILHFTEHSGLTTGVAFGHHAKFIASTGMDRSLKFYSL
Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry
il	

Western Blot 1:500 - 1:2000, Immunohistochemistry 1:50 - 1:200,


Immunocytochemistry/ Immunofluorescence 1:50 - 1:200,

Immunohistochemistry-Paraffin


Images

Recommended Dilutions


Western Blot: PRP19 Antibody (4M3Z1) [NBP3-16853] - Western blot analysis of extracts of various cell lines, using PRP19 Rabbit mAb (NBP3 -16853) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 10s.




Western Blot: PRP19 Antibody (4M3Z1) [NBP3-16853] - Western blot analysis of extracts of Rat spleen, using PRP19 Rabbit mAb (NBP3-16853) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 3min.




Immunocytochemistry/Immunofluorescence: PRP19 Antibody (4M3Z1) [NBP3-16853] - Immunofluorescence analysis of U-2 OS cells using PRP19 Rabbit mAb (NBP3-16853) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.




Immunohistochemistry: PRP19 Antibody (4M3Z1) [NBP3-16853] - Immunohistochemistry analysis of PRP19 in paraffin-embedded human colon carcinoma tissue using PRP19 Rabbit mAb at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry: PRP19 Antibody (4M3Z1) [NBP3-16853] - Immunohistochemistry analysis of PRP19 in paraffin-embedded mouse testis tissue using PRP19 Rabbit mAb at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry: PRP19 Antibody (4M3Z1) [NBP3-16853] - Immunohistochemistry analysis of PRP19 in paraffin-embedded mouse intestin tissue using PRP19 Rabbit mAb at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry: PRP19 Antibody (4M3Z1) [NBP3-16853] - Immunohistochemistry analysis of PRP19 in paraffin-embedded rat colon tissue using PRP19 Rabbit mAb at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry: PRP19 Antibody (4M3Z1) [NBP3-16853] - Immunohistochemistry analysis of PRP19 in paraffin-embedded mouse brain tissue using PRP19 Rabbit mAb at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP3-16853-100ul

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

H00027339-P01-10ug Recombinant Human PRP19 GST (N-Term) Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-16853

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

