# **Product Datasheet**

# Chorionic Gonadotropin alpha Chain (hCG alpha) Antibody NBP3-18630-100ug

Unit Size: 100 ug

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-18630

Updated 6/15/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-18630



## NBP3-18630-100ug


Chorionic Gonadotropin alpha Chain (hCG alpha) Antibody

Chorionic Gonadotropin alpha	a Chain (hCG alpha) Antibody	
Product Information		
Unit Size	100 ug	
Concentration	Please see the vial label for concentration. If unlisted please contact technical services.	
Storage	Store at -20C. Avoid freeze-thaw cycles.	
Clonality	Polyclonal	
Preservative	0.01% Sodium Azide	
Isotype	IgG	
Purity	Affinity purified	
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2	
<b>Product Description</b>		
Description	Store vial at -20C prior to opening. Aliquot contents and freeze at -20C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4 degrees C as an undiluted liquid. Dilute only prior to immediate use.  This product was affinity purified from monospecific antiserum by immunoaffinity purification.	
Host	Goat	
Gene ID	1081	
Gene Symbol	CGA	
Species	Human	
Specificity/Sensitivity	This product reacts with Full Human Chorionic Gonadotropin (hCG) and Human Chorionic Gonadotropin Alpha chain (alpha-hCG).	
Immunogen	Prepared from whole goat serum produced by repeated immunizations with Human Chorionic Gonadotropin alpha Chain (hCG alpha) native protein.	
Notes	Store vial at -20C prior to opening. Aliquot contents and freeze at -20C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4 degrees C as an undiluted liquid. Dilute only prior to immediate use.  This product was affinity purified from monospecific antiserum by immunoaffinity purification.	
<b>Product Application Details</b>		
Applications	Western Blot, Dot Blot, ELISA, Immunohistochemistry, Immunohistochemistry-Paraffin	
Recommended Dilutions	Western Blot, ELISA, Immunohistochemistry 1:500 - 1:1000, Immunohistochemistry-Paraffin 1:500-1:1000, Dot Blot	
Application Notes	This product has been tested in ELISA and WB. Expect a band at ~14.9kDa for the alpha chain in western blot using appropriate tissues and lysates. ELISA reactivity expected against alpha hCG with an expected titer of ~1:100,000 and full hCG at ~1:50,000 dilution.	




#### **Images**

Western Blot: Chorionic Gonadotropin alpha Chain (hCG alpha) Antibody [NBP3-18630] - Lane 1: Opal Prestained Molecular Weight Marker. Lane 2: hCG alpha (0.05ug) [+]. Lane 3: hCG alpha/U-20S WCL (0.05ug/20ug) [+]. Lane 4: U-20S WCL (20ug) [-]. Lane 5: HeLa WCL (35ug) [+]. Primary Antibody: Anti-alpha-hCG at 1:1000 overnight at 2-8C. Secondary Antibody: Donkey Anti-Goat IgG mx8 HRP Conjugated at 1:40,000 for 30 mins at RT. Block: BlockOut Buffer. Exposure: 30 sec. Predicted MW: alpha subunit ~14.9kDa, full hCG ~37kDa, luteinizing hormone (LH) ~29kDa, thyroid stimulating hormone (TSH) ~28kDa, follicle stimulating hormone (FSH) ~30kDa. Observed MW: alpha subunit ~15-20kDa, full hCG ~35kDa.



ELISA: Chorionic Gonadotropin alpha Chain (hCG alpha) Antibody [NBP3-18630] - Each well was coated in duplicate at 10ug/mL of hCG alpha (RED) and full hCG (BLUE) protein. The starting dilution of antibody was 5ug/mL and the X-axis represents the Log10 of a 3-fold dilution. This titration is a 4-parameter curve fit where the IC50 is defined as the titer of the antibody. Assay performed using 3% Fish Gel/PBS Blocking Buffer, Donkey Anti-Goat IgG MX8 HRP Conjugated Secondary, and TMB ELISA Peroxidase substrate.





### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966 novus@novusbio.com

# Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom

Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

### **General Contact Information**

www.novusbio.com

Technical Support: technical@novusbio.com

Orders: orders@novusbio.com General: novus@novusbio.com

### Products Related to NBP3-18630-100ug

HAF017	Rabbit anti-Goat IgG Secondary	Antibody [HRP (Horseradish
11/11 01/	Trabbit anti Obat Ido Occondary	

Peroxidase)1

HAF109 Donkey anti-Goat IgG Secondary Antibody [HRP (Horseradish

Peroxidase)]

NB410-28088-1mg Goat IgG Isotype Control

H00001081-P01-2ug Recombinant Human Chorionic Gonadotropin alpha Chain (hCG alpha)

GST (N-Term) Protein

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-18630

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications



