Product Datasheet

BAP1 Antibody (C-4) NBP3-21456

Unit Size: 0.2 ml Store at 4C.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-21456

Updated 6/4/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-21456



NBP3-21456

BAP1 Antibody (C-4)

Product Information Unit Size 0.2 ml Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C. Clonality Monoclonal Clone C-4 Preservative 0.1% Sodium Azide Isotype IgG1 Purity Protein A or G purified Buffer PBS, 0.1% gelatin Product Description Host Mouse Gene ID 8314 Gene Symbol BAP1 Species Human Immunogen Amino acids 430-729 of human BAP1 Product Application Details Applications Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry, Immunohistochemistry-Paraffin 1:50-1:500 Application Notes Pre-treatment: Heat induced epitope retrieval in 10 mM citrate buffer, pH6.0, or in 50 mM Tris buffer pH9.5, for 20 minutes is required for IHC staining on formalin-fixed, paraffin embedded tissue sections.	Brit 1 Filiabody (O +)	
Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C. Clonality Monoclonal Clone C-4 Preservative 0.1% Sodium Azide Isotype IgG1 Purity Protein A or G purified Buffer PBS, 0.1% gelatin Product Description Host Mouse Gene ID 8314 Gene Symbol BAP1 Species Human Immunogen Amino acids 430-729 of human BAP1 Product Application Details Applications Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry, Immunohistochemistry-Paraffin 1:50-1:500 Application Notes Pre-treatment: Heat induced epitope retrieval in 10 mM citrate buffer, pH6.0, or in 50 mM Tris buffer pH9.5, for 20 minutes is required for IHC staining on	Product Information	
Storage Store at 4C. Clonality Monoclonal Clone C-4 Preservative 0.1% Sodium Azide Isotype IgG1 Purity Protein A or G purified Buffer PBS, 0.1% gelatin Product Description Host Mouse Gene ID 8314 Gene Symbol BAP1 Species Human Immunogen Amino acids 430-729 of human BAP1 Product Application Details Application Notes Inmunohistochemistry, Immunohistochemistry-Paraffin 1:50-1:500 Application Notes Pre-treatment: Heat induced epitope retrieval in 10 mM citrate buffer, pH6.0, or in 50 mM Tris buffer pH9.5, for 20 minutes is required for IHC staining on	Unit Size	0.2 ml
Clone C-4 Preservative 0.1% Sodium Azide Isotype IgG1 Purity Protein A or G purified Buffer PBS, 0.1% gelatin Product Description Host Mouse Gene ID 8314 Gene Symbol BAP1 Species Human Immunogen Amino acids 430-729 of human BAP1 Product Application Details Applications Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry, Immunohistochemistry-Paraffin 1:50-1:500 Application Notes Pre-treatment: Heat induced epitope retrieval in 10 mM citrate buffer, pH6.0, or in 50 mM Tris buffer pH9.5, for 20 minutes is required for IHC staining on	Concentration	· ·
Clone C-4 Preservative 0.1% Sodium Azide Isotype IgG1 Purity Protein A or G purified Buffer PBS, 0.1% gelatin Product Description Host Mouse Gene ID 8314 Gene Symbol BAP1 Species Human Immunogen Amino acids 430-729 of human BAP1 Product Application Details Applications Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry, Immunohistochemistry-Paraffin 1:50-1:500 Application Notes Pre-treatment: Heat induced epitope retrieval in 10 mM citrate buffer, pH6.0, or in 50 mM Tris buffer pH9.5, for 20 minutes is required for IHC staining on	Storage	Store at 4C.
Preservative 0.1% Sodium Azide Isotype IgG1 Purity Protein A or G purified Buffer PBS, 0.1% gelatin Product Description Host Mouse Gene ID 8314 Gene Symbol BAP1 Species Human Immunogen Amino acids 430-729 of human BAP1 Product Application Details Applications Immunohistochemistry, Immunohistochemistry-Paraffin 1:50-1:500 Application Notes Pre-treatment: Heat induced epitope retrieval in 10 mM citrate buffer, pH6.0, or in 50 mM Tris buffer pH9.5, for 20 minutes is required for IHC staining on	Clonality	Monoclonal
Isotype IgG1 Purity Protein A or G purified Buffer PBS, 0.1% gelatin Product Description Host Mouse Gene ID 8314 Gene Symbol BAP1 Species Human Immunogen Amino acids 430-729 of human BAP1 Product Application Details Applications Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry, Immunohistochemistry-Paraffin 1:50-1:500 Application Notes Pre-treatment: Heat induced epitope retrieval in 10 mM citrate buffer, pH6.0, or in 50 mM Tris buffer pH9.5, for 20 minutes is required for IHC staining on	Clone	C-4
Purity Protein A or G purified Buffer PBS, 0.1% gelatin Product Description Host Mouse Gene ID 8314 Gene Symbol BAP1 Species Human Immunogen Amino acids 430-729 of human BAP1 Product Application Details Applications Immunohistochemistry, Immunohistochemistry-Paraffin 1:50-1:500 Application Notes Pre-treatment: Heat induced epitope retrieval in 10 mM citrate buffer, pH6.0, or in 50 mM Tris buffer pH9.5, for 20 minutes is required for IHC staining on	Preservative	0.1% Sodium Azide
Buffer PBS, 0.1% gelatin Product Description Host Mouse Gene ID 8314 Gene Symbol BAP1 Species Human Immunogen Amino acids 430-729 of human BAP1 Product Application Details Applications Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry, Immunohistochemistry-Paraffin 1:50-1:500 Application Notes Pre-treatment: Heat induced epitope retrieval in 10 mM citrate buffer, pH6.0, or in 50 mM Tris buffer pH9.5, for 20 minutes is required for IHC staining on	Isotype	IgG1
Product Description Host Mouse Gene ID 8314 Gene Symbol BAP1 Species Human Immunogen Amino acids 430-729 of human BAP1 Product Application Details Applications Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry, Immunohistochemistry-Paraffin 1:50-1:500 Application Notes Pre-treatment: Heat induced epitope retrieval in 10 mM citrate buffer, pH6.0, or in 50 mM Tris buffer pH9.5, for 20 minutes is required for IHC staining on	Purity	Protein A or G purified
Host Mouse Gene ID 8314 Gene Symbol BAP1 Species Human Immunogen Amino acids 430-729 of human BAP1 Product Application Details Applications Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry, Immunohistochemistry-Paraffin 1:50-1:500 Application Notes Pre-treatment: Heat induced epitope retrieval in 10 mM citrate buffer, pH6.0, or in 50 mM Tris buffer pH9.5, for 20 minutes is required for IHC staining on	Buffer	PBS, 0.1% gelatin
Gene Symbol BAP1 Species Human Immunogen Amino acids 430-729 of human BAP1 Product Application Details Applications Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry, Immunohistochemistry-Paraffin 1:50-1:500 Application Notes Pre-treatment: Heat induced epitope retrieval in 10 mM citrate buffer, pH6.0, or in 50 mM Tris buffer pH9.5, for 20 minutes is required for IHC staining on	Product Description	
Gene Symbol Species Human Immunogen Amino acids 430-729 of human BAP1 Product Application Details Applications Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry, Immunohistochemistry-Paraffin 1:50-1:500 Application Notes Pre-treatment: Heat induced epitope retrieval in 10 mM citrate buffer, pH6.0, or in 50 mM Tris buffer pH9.5, for 20 minutes is required for IHC staining on	Host	Mouse
Species Human Amino acids 430-729 of human BAP1 Product Application Details Applications Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry, Immunohistochemistry-Paraffin 1:50-1:500 Application Notes Pre-treatment: Heat induced epitope retrieval in 10 mM citrate buffer, pH6.0, or in 50 mM Tris buffer pH9.5, for 20 minutes is required for IHC staining on	Gene ID	8314
Immunogen Amino acids 430-729 of human BAP1 Product Application Details Applications Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry, Immunohistochemistry-Paraffin 1:50-1:500 Application Notes Pre-treatment: Heat induced epitope retrieval in 10 mM citrate buffer, pH6.0, or in 50 mM Tris buffer pH9.5, for 20 minutes is required for IHC staining on	Gene Symbol	BAP1
Product Application Details Applications Immunohistochemistry, Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry, Immunohistochemistry-Paraffin 1:50-1:500 Application Notes Pre-treatment: Heat induced epitope retrieval in 10 mM citrate buffer, pH6.0, or in 50 mM Tris buffer pH9.5, for 20 minutes is required for IHC staining on	Species	Human
ApplicationsImmunohistochemistry, Immunohistochemistry-ParaffinRecommended DilutionsImmunohistochemistry, Immunohistochemistry-Paraffin 1:50-1:500Application NotesPre-treatment: Heat induced epitope retrieval in 10 mM citrate buffer, pH6.0, or in 50 mM Tris buffer pH9.5, for 20 minutes is required for IHC staining on	Immunogen	Amino acids 430-729 of human BAP1
Recommended Dilutions Immunohistochemistry, Immunohistochemistry-Paraffin 1:50-1:500 Application Notes Pre-treatment: Heat induced epitope retrieval in 10 mM citrate buffer, pH6.0, or in 50 mM Tris buffer pH9.5, for 20 minutes is required for IHC staining on	Product Application Details	
Application Notes Pre-treatment: Heat induced epitope retrieval in 10 mM citrate buffer, pH6.0, or in 50 mM Tris buffer pH9.5, for 20 minutes is required for IHC staining on	Applications	Immunohistochemistry, Immunohistochemistry-Paraffin
in 50 mM Tris buffer pH9.5, for 20 minutes is required for IHC staining on	Recommended Dilutions	Immunohistochemistry, Immunohistochemistry-Paraffin 1:50-1:500
	Application Notes	in 50 mM Tris buffer pH9.5, for 20 minutes is required for IHC staining on





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966 novus@novusbio.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: technical@novusbio.com

Orders: orders@novusbio.com General: novus@novusbio.com

Products Related to NBP3-21456

HAF007 Goat anti-Mouse IgG Secondary Antibody [HRP]

NB720-B Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]

NBP1-97005-0.5mg Mouse IgG1 Isotype Control (MG1)
NBP2-57089PEP BAP1 Recombinant Protein Antigen

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-21456

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

