# **Product Datasheet**

# Myosin heavy chain 11 Antibody (MYH11/8185R) - Azide and BSA Free NBP3-24304

Unit Size: 100 ug

Store at -20 to -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-24304

Updated 7/16/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP3-24304



#### NBP3-24304

| Myosin heavy chain 11 Antibody (MYH11/8185R) - Azide and BSA Free |   |
|---|---|
| Product Information   |   |
| Unit Size   | 100 ug  |
| Concentration   | 1 mg/ml   |
| Storage   | Store at -20 to -80C. Avoid freeze-thaw cycles.   |
| Clonality   | Monoclonal  |
| Clone   | MYH11/8185R   |
| Preservative  | No Preservative   |
| Isotype   | IgG Kappa   |
| Purity  | Protein A or G purified   |
| Buffer  | 10mM PBS  |
| Product Description   |   |
| Description   | Antibody without azide - store at -20 to -80C.  |
| Host  | Rabbit  |
| Gene ID   | 4629  |
| Gene Symbol   | MYH11   |
| Species   | Human   |
| Marker  | SM-MHCLeiomyosarcoma & Myoepithelial Cell Marker  |
| Specificity/Sensitivity   | The antibody reacts with smooth muscle cells and myoepithelial cells, but not with myofibroblasts. It is very helpful in distinguishing between benign sclerosing breast lesions and infiltrating carcinomas in difficult cases since it strongly stains the myoepithelial layer in the benign lesions while it is negative in the infiltrating carcinomas. |
| Immunogen   | Recombinant fragment (around aa1-200) of human Myosin heavy chain 11 protein (exact sequence is proprietary)  |
| Product Application Details                                       |   |
| Applications  | Immunohistochemistry-Paraffin   |
| Recommended Dilutions   | Immunohistochemistry-Paraffin 1-2ug/ml  |
| Application Notes   | Immunohistochemistry-Paraffin: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes. Optimal dilution for a specific application should be determined.   |





### **Novus Biologicals USA**

10730 E. Briarwood Avenue Centennial, CO 80112 USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com
Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

## **Products Related to NBP3-24304**

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP1-87026PEP Myosin heavy chain 11 Recombinant Protein Antigen
H00027130-P02-10ug Recombinant Human Inversin GST (N-Term) Protein

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-24304

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

