# **Product Datasheet**

# TTF-1 / NKX2-1 Antibody (6C11) - BSA Free NBP3-26111-100ul

Unit Size: 100 ul

Store at -20 to -70C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-26111

Updated 7/30/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP3-26111



# NBP3-26111-100ul


TTF-1 / NKX2-1 Antibody (6C11) - BSA Free

TTF-T/ NKAZ-T AHIIDOUY (OCTT) - DOA FTEE	
Product Information	
Unit Size	100 ul
Concentration	Please see the vial label for concentration. If unlisted please contact technical services.
Storage	Store at -20 to -70C. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	6C11
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Affinity purified
Buffer	PBS, pH 7.4, 150mM NaCl, and 50% glycerol
Product Description	
Host	Rabbit
Gene ID	7080
Gene Symbol	NKX2-1
Species	Human
Immunogen	A synthesized peptide derived from Human TTF-1/NKX2-1 [UniProt P43699]
Product Application Details	
Applications	Western Blot, ELISA, Flow Cytometry, Immunohistochemistry
Recommended Dilutions	Western Blot 1:500-1:5000, Flow Cytometry 1:20-1:200, ELISA, Immunohistochemistry 1:50-1:200




#### **Images**

Immunohistochemistry: TTF-1 / NKX2-1 Antibody (6C11) [NBP3-26111] - Image of TTF-1 / NKX2-1 Antibody (6C11) diluted at 1:100 and staining in paraffin-embedded human lung tissue performed. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.



Immunohistochemistry: TTF-1 / NKX2-1 Antibody (6C11) [NBP3-26111] - Image of TTF-1 / NKX2-1 Antibody (6C11) diluted at 1:100 and staining in paraffin-embedded human lung tissue performed. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.




Western Blot: TTF-1 / NKX2-1 Antibody (6C11) [NBP3-26111] - Positive Western Blot detected in: HepG2 whole cell lysate, U251 whole cell

lysate, U87 whole cell lysate.


All lanes: TTF-1 / NKX2-1 Antibody at 1: 2000

Secondary: Goat polyclonal to rabbit IgG at 1/50000 dilution.

Predicted band size: 39 kDa Observed band size: 39 kDa



Flow Cytometry: TTF-1 / NKX2-1 Antibody (6C11) [NBP3-26111] - Overlay histogram showing A549 cells stained with TTF-1 / NKX2-1 Antibody (6C11) (red line) at 1:50. The cells were fixed with 70% Ethylalcohol (18h) and then incubated in 10% normal goat serum to block non-specific protein-protein interactions followed by the antibody (1ug/1\*10^6 cells) for 1 h at 4C. The secondary antibody used was FITC-conjugated goat anti-rabbit IgG (H+L) at 1/200 dilution for 30min at 4C. Control antibody (green line) was Rabbit IgG (1ug/1\*10^6 cells) used under the same conditions. Acquisition of >10,000 events was performed.





## Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

### Products Related to NBP3-26111-100ul

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

H00007080-Q01-10ug Recombinant Human TTF-1 / NKX2-1 GST (N-Term) Protein

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-26111

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

