Product Datasheet

PBR Antibody (23G2) - BSA Free NBP3-26270-100ul

Unit Size: 100 ul

Store at -20 to -70C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-26270

Updated 7/30/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP3-26270



NBP3-26270-100ul

PBR Antibody (23G2) - BSA Free

,	
Product Information	
Unit Size	100 ul
Concentration	Please see the vial label for concentration. If unlisted please contact technical services.
Storage	Store at -20 to -70C. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	23G2
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Affinity purified
Buffer	PBS, pH 7.4, 150mM NaCl, and 50% glycerol
Draduat Decariation	

Product Description	
Host	Rabbit
Gene ID	706
Gene Symbol	TSPO
Species	Human, Mouse
Immunogen	A synthesized peptide derived from Human PBR [UniProt P30536]

Product Application Details	
Applications	Western Blot, ELISA, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunoprecipitation
Recommended Dilutions	Western Blot 1:500-1:5000, Flow Cytometry, ELISA, Immunohistochemistry 1:50 -1:200, Immunocytochemistry/ Immunofluorescence 1:20-1:200, Immunoprecipitation 1:200-1:1000

Images


Immunoprecipitation: PBR Antibody (23G2) [NBP3-26270] -

Immunoprecipitating PBR in Hela whole cell lysate.


Lane 1: Rabbit control IgG instead of NBP3-26270 in Hela whole cell lysate. For western blotting, a HRP-conjugated Protein G antibody was used as the secondary antibody (1/2000).

Lane 2: NBP3-26270 (3ug) + Hela whole cell lysate (500ug).


Lane 3: Hela whole cell lysate (20ug).



Immunohistochemistry: PBR Antibody (23G2) [NBP3-26270] - Image of PBR Antibody (23G2) diluted at 1:117 and staining in paraffin-embedded human pancreatic cancer performed. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Immunohistochemistry: PBR Antibody (23G2) [NBP3-26270] - Image of PBR Antibody (23G2) diluted at 1:117 and staining in paraffin-embedded human pancreatic cancer performed. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.




Western Blot: PBR Antibody (23G2) [NBP3-26270] - Positive Western Blot detected in: Hela whole cell lysate, MCF-7 whole cell lysate, HepG2 whole cell lysate, A549 whole cell lysate, Mouse kidney tissue.


All lanes: PBR Antibody at 1.2ug/ml.

Secondary: Goat polyclonal to rabbit IgG at 1/50000 dilution.


Predicted band size: 19, 11 KDa Observed band size: 19 KDa



Immunocytochemistry/Immunofluorescence: PBR Antibody (23G2) [NBP3-26270] - Staining of PC3 cells with PBR Antibody (23G2) at 1:39, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4C. The secondary antibody was Alexa Fluor 488-conjugated Goat Anti-Rabbit IgG (H+L).



Flow Cytometry: PBR Antibody (23G2) [NBP3-26270+156:156D172158:192153:192D172158:192149:192D172158:192 145:192D1168:192] - Overlay histogram showing HepG2 cells stained with PBR Antibody (23G2) (red line) at 1:50. The cells were fixed with 70% Ethylalcohol (18h) and then permeabilized with 0.3% Triton X-100 for 2 min. The cells were then incubated in 1x PBS /10% normal goat serum to block non-specific protein-protein interactions followed by primary antibody for 1 h at 4C. The secondary antibody used was FITC goat anti-rabbit IgG (H+L) at 1/200 dilution for 1 h at 4C. Control antibody (green line) was used under the same conditions. Acquisition of >10,000 events was performed.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP3-26270-100ul

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

H00000706-P01-10ug Recombinant Human PBR GST (N-Term) Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-26270

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

