Product Datasheet

CCL23/Ck beta 8-1/MIP3 Antibody (CCL23/4036) NBP3-26728-100ug

Unit Size: 100 ug
Store at 4C.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-26728

Updated 7/16/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-26728



NBP3-26728-100ug

| CCL23/Ck beta 8-1/MIP3 Antibod | ly (CCL23/4036) | |
|--------------------------------|--|--|
| Product Information | | |
| Unit Size | 100 ug | |
| Concentration | 0.2 mg/ml | |
| Storage | Store at 4C. | |
| Clonality | Monoclonal | |
| Clone | CCL23/4036 | |
| Preservative | 0.05% Sodium Azide | |
| Isotype | IgG2c Kappa | |
| Purity | Protein A or G purified | |
| Buffer | 10mM PBS with 0.05% BSA | |
| Product Description | | |
| Description | Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Non-hazardous. No MSDS required. | |
| Host | Mouse | |
| Gene ID | 6368 | |
| Gene Symbol | CCL23 | |
| Species | Human | |
| Immunogen | Recombinant fragment (around aa1-120) of human CCL23/Ck beta 8-1/MIP3 protein (exact sequence is proprietary) | |
| Product Application Details | | |
| Applications | Immunohistochemistry-Paraffin | |
| Recommended Dilutions | Immunohistochemistry-Paraffin 1-2ug/ml for 30 minutes at RT | |
| Application Notes | Immunohistochemistry-Paraffin 1-2ug/ml for 30 minutes at RT. Optimal dilution for a specific application should be determined Staining of formalin-fixed tissues | |

| Product Application Details | | |
|-----------------------------|--|--|
| Applications | Immunohistochemistry-Paraffin | |
| Recommended Dilutions | Immunohistochemistry-Paraffin 1-2ug/ml for 30 minutes at RT | |
| Application Notes | Immunohistochemistry-Paraffin 1-2ug/ml for 30 minutes at RT. Optimal dilution for a specific application should be determined Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes | |



| Images | |
|---|---|
| Formalin-fixed, paraffin-embedded human adrenal gland stained with CCL23/Ck beta 8-1/MIP3 Antibody (CCL23/4036). HIER: Tris/EDTA, pH9.0, 45min. 2°C: HRP-polymer, 30min. DAB, 5min. | X |
| Analysis of Protein Array containing more than 19,000 full-length human proteins using CCL23-Monospecific Mouse Monoclonal Antibody (CCL23/4036). | X |





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom

Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP3-26728-100ug

HAF007 Goat anti-Mouse IgG Secondary Antibody [HRP]

NB720-B Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
NBP2-35021-5ug Recombinant Human CCL23/Ck beta 8-1/MIP3 Protein

210-TA-005 TNF-alpha [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-26728

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

