

# Product Datasheet

## **SARS-CoV-2 Spike S1 Antibody - BSA Free NBP3-26918-0.1mg**

Unit Size: 0.1 mg

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

[www.novusbio.com](http://www.novusbio.com)



[technical@novusbio.com](mailto:technical@novusbio.com)

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:  
[www.novusbio.com/NBP3-26918](http://www.novusbio.com/NBP3-26918)

Updated 4/1/2024 v.20.1

Earn rewards for product  
reviews and publications.

Submit a publication at [www.novusbio.com/publications](http://www.novusbio.com/publications)

Submit a review at [www.novusbio.com/reviews/destination/NBP3-26918](http://www.novusbio.com/reviews/destination/NBP3-26918)



**NBP3-26918-0.1mg**

SARS-CoV-2 Spike S1 Antibody - BSA Free


Product Information	
Unit Size	0.1 mg
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Peptide affinity purified
Buffer	PBS

Product Description	
Host	Rabbit
Gene ID	43740568
Species	SARS-CoV-2
Immunogen	SARS-CoV-2 Spike S1 Antibody was raised against a peptide corresponding to 16 amino acids near the amino terminus of SARS-CoV-2 (COVID-19) Spike S1 glycoprotein. The immunogen is located within the first 50 amino acids of SARS-CoV-2 (COVID-19) Spike S1 protein.


Product Application Details	
Applications	Western Blot, ELISA, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry
Recommended Dilutions	Western Blot 1-2 ug/mL, ELISA, Immunohistochemistry 0.5 ug/mL, Immunocytochemistry/ Immunofluorescence
Application Notes	SARS-CoV-2 Spike S1 Antibody can be used for the detection of SARS-CoV-2 (COVID-19) Spike protein in ELISA. It will detect 4 ng of free peptide at 1 ug/mL.

**Images**


ELISA: SARS-CoV-2 Spike S1 Antibody - BSA Free [NBP3-26918] - Validation with SARS-CoV-2 Spike Recombinant Protein. Antibodies: SARS-CoV-2 Spike S1 antibody, NBP3-26918. A direct ELISA was performed using SARS-CoV-2 (COVID-19) Spike S1 recombinant protein as coating antigen and the anti-SARS-CoV-2 Spike S1 antibody as the capture antibody. Secondary: Goat anti-rabbit IgG HRP conjugate at 1:20000 dilution. Detection range is from 8 ng/mL to 1000 ng/mL.




**Immunohistochemistry: SARS-CoV-2 Spike S1 Antibody - BSA Free [NBP3-26918] - Validation of SARS-CoV-2 Spike S1 in COVID-19 Patient Lung.** Analysis of paraffin-embedded COVID-19 patient lung tissue using anti- SARS-CoV-2 Spike S1 antibody (0.5 ug/mL). Tissue was fixed with formaldehyde and blocked with 10% serum for 1 h at RT; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody overnight at 4C. A goat anti-rabbit IgG H&L (HRP) at 1/250 was used as secondary. Counter stained with Hematoxylin. Strong signal of SARS-COV-2 spike protein was observed in macrophage of COVID-19 patient lung, but not in non-COVID-19 patient lung.




**Western Blot: SARS-CoV-2 Spike S1 Antibody - BSA Free [NBP3-26918] - Validation with SARS-CoV-2 Spike Recombinant Protein.** Loading: 50 ng per lane of SARS-CoV-2 Spike S1 recombinant protein. Antibodies: SARS-CoV-2 Spike S1, NBP3-26918, 1h incubation at RT in 5% NFDm/TBST. Secondary: Goat anti-rabbit IgG HRP conjugate at 1:10000 dilution. Lane 1: 1 ug/mL and Lane 2: 2 ug/mL.



**Immunocytochemistry/Immunofluorescence: SARS-CoV-2 Spike S1 Antibody - BSA Free [NBP3-26918] - Validation of SARS-CoV-2 Spike S1 in 293 Transfected Cells.** Analysis of 4% paraformaldehyde-fixed Spike transfected 293 cells labeling SARS-CoV-2 Spike S1 with NBP3-26918 at 5 ug/mL, followed by goat anti-rabbit IgG secondary antibody at 1/500 dilution (green) and DAPI staining (blue).



**Western Blot: SARS-CoV-2 Spike S1 Antibody - BSA Free [NBP3-26918] - Overexpression Validation in Spike Transfected 293 Cells.** Loading: 10 ug per lane of 293 cell lysate. Antibodies: SARS-CoV-2 Spike S1, NBP3-26918 (1 ug/mL), 1h incubation at RT in 5% NFDm/TBST. Secondary: Goat anti-rabbit IgG HRP conjugate at 1:10000 dilution. Lane 1: WT 293 cells and Lane 2: SARS-CoV-2 Spike overexpressed 293 cells.





### **Novus Biologicals USA**

10730 E. Briarwood Avenue  
Centennial, CO 80112  
USA  
Phone: 303.730.1950  
Toll Free: 1.888.506.6887  
Fax: 303.730.1966  
nb-customerservice@bio-techne.com

### **Bio-Techne Canada**

21 Canmotor Ave  
Toronto, ON M8Z 4E6  
Canada  
Phone: 905.827.6400  
Toll Free: 855.668.8722  
Fax: 905.827.6402  
canada.inquires@bio-techne.com

### **Bio-Techne Ltd**

19 Barton Lane  
Abingdon Science Park  
Abingdon, OX14 3NB, United Kingdom  
Phone: (44) (0) 1235 529449  
Free Phone: 0800 37 34 15  
Fax: (44) (0) 1235 533420  
info.EMEA@bio-techne.com

### **General Contact Information**

www.novusbio.com  
Technical Support: nb-technical@bio-techne.com  
Orders: nb-customerservice@bio-techne.com  
General: novus@novusbio.com

### **Products Related to NBP3-26918-0.1mg**

---

HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control
NBP2-90985	Recombinant SARS-CoV-2 Spike S1 His (C-Term) Protein

---

### **Limitations**

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit [www.novusbio.com/guarantee](http://www.novusbio.com/guarantee)

Earn gift cards/discounts by submitting a review: [www.novusbio.com/reviews/submit/NBP3-26918](http://www.novusbio.com/reviews/submit/NBP3-26918)

Earn gift cards/discounts by submitting a publication using this product:  
[www.novusbio.com/publications](http://www.novusbio.com/publications)

