

Product Datasheet

3-methylcytosine (3-mC) Antibody (PSH03-21) NBP3-31984

Unit Size: 100 ul

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP3-31984

Updated 8/7/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP3-31984



NBP3-31984**3-methylcytosine (3-mC) Antibody (PSH03-21)**

Product Information	
Unit Size	100 ul
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	PSH03-21
Preservative	0.05% Sodium Azide
Isotype	IgG
Purity	Protein A purified
Buffer	PBS (pH7.4), 0.1% BSA and 40% Glycerol
Product Description	
Host	Rabbit
Species	Non-species specific
Immunogen	3-mc-OVA
Product Application Details	
Applications	Dot Blot, Immunocytochemistry/ Immunofluorescence
Recommended Dilutions	Immunocytochemistry/ Immunofluorescence 1:100, Dot Blot 1:2000



Images

Dot Blot: 3-methylcytosine (3-mC) Antibody (PSH03-21) [NBP3-31984] - Dot blot analysis of 3-methylcytosine (3-mC) on different proteins with Rabbit anti-3mC antibody (NBP3-31984) at 1/2,000 dilution. Goat Anti-Rabbit IgG - HRP Secondary Antibody at 1/50,000 dilution for 1 hour at room temperature.

Lane 1: Uridine-BSA (negative)
 Lane 2: Cytidine-BSA (negative)
 Lane 3: 5-Methylcytosine-BSA (negative)
 Lane 4: 3-Methylcytosine-BSA (positive)
 Lane 5: 5-Hydroxymethylcytidine-BSA (negative)


Proteins loading: 50ng, 25ng, 12.5ng, 6.25ng;

Blocking and dilution buffer: 5% NFDM/TBST;

Exposure time: 1 minute.

Immunocytochemistry/ Immunofluorescence: 3-methylcytosine (3-mC) Antibody (PSH03-21) [NBP3-31984] - Immunocytochemistry analysis of U-2 OS cells treated with 4mM Methyl Methanesulfonate for 1 hour labeling 3-methylcytosine (3-mC) with Rabbit anti-3-methylcytosine (3-mC) antibody (NBP3-31984) at 1/100 dilution.

Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-3-methylcytosine (3-mC) antibody (NBP3-31984) at 1/100 dilution in 1% BSA in PBST overnight at 4 °C. Goat Anti-Rabbit IgG H&L (iFluor™ 488) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-
techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP3-31984

HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-31984

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

