

Product Datasheet

Aly Antibody (JE36-45) NBP3-32033

Unit Size: 100 ul

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP3-32033

Updated 8/7/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP3-32033



NBP3-32033

Aly Antibody (JE36-45)

Product Information	
Unit Size	100 ul
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	JE36-45
Preservative	0.05% Sodium Azide
Isotype	IgG
Purity	Protein A purified
Buffer	1*TBS (pH7.4), 0.05% BSA and 40% Glycerol
Target Molecular Weight	27 kDa
Product Description	
Host	Rabbit
Gene ID	10189
Gene Symbol	ALYREF
Species	Human, Mouse, Rat
Immunogen	Synthetic peptide within Human Aly aa 208-257 / 257. (Uniprot: Q86V81)
Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry
Recommended Dilutions	Western Blot 1:1000, Flow Cytometry 1:1000, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1:250, Immunohistochemistry-Paraffin 1:2000



Images

Western Blot: Aly Antibody (JE36-45) [NBP3-32033] - Western blot analysis of Aly on different lysates with Rabbit anti-Aly antibody (NBP3-32033) at 1/1,000 dilution.

Lane 1: HepG2 cell lysate (20 ug/Lane)
 Lane 2: K-562 cell lysate (20 ug/Lane)
 Lane 3: 293T cell lysate (20 ug/Lane)
 Lane 4: A549 cell lysate (20 ug/Lane)
 Lane 5: SK-OV-3 cell lysate (20 ug/Lane)
 Lane 6: HL-60 cell lysate (20 ug/Lane)
 Lane 7: NIH/3T3 cell lysate (20 ug/Lane)
 Lane 8: Neuro-2a cell lysate (20 ug/Lane)
 Lane 9: C6 cell lysate (20 ug/Lane)
 Lane 10: Mouse testis tissue lysate (40 ug/Lane)
 Lane 11: Mouse spleen tissue lysate (40 ug/Lane)
 Lane 12: Rat testis tissue lysate (40 ug/Lane)

Predicted band size: 27 kDa
 Observed band size: 27 kDa

Exposure time: 2 minutes;

4-20% SDS-PAGE gel.




Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (NBP3-32033) at 1/1,000 dilution was used in 5% NFDM/TBST at 4C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody at 1/50,000 dilution was used for 1 hour at room temperature.

Immunohistochemistry: Aly Antibody (JE36-45) [NBP3-32033] - Immunohistochemical analysis of paraffin-embedded human brain tissue with Rabbit anti-Aly antibody (NBP3-32033) at 1/2,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) for 2 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (NBP3-32033) at 1/2,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.


Immunocytochemistry/ Immunofluorescence: Aly Antibody (JE36-45) [NBP3-32033] - Immunocytochemistry analysis of HepG2 cells labeling Aly with Rabbit anti-Aly antibody (NBP3-32033) at 1/250 dilution.

Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-Aly antibody (NBP3-32033) at 1/250 dilution in 1% BSA in PBST overnight at 4 °C. Goat Anti-Rabbit IgG H&L (iFluor™ 488) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.



Flow Cytometry: Aly Antibody (JE36-45) [NBP3-32033] - Flow cytometric analysis of HepG2 cells labeling Aly.

Cells were fixed and permeabilized. Then stained with the primary antibody (NBP3-32033, 1 μ g/mL) (red) compared with Rabbit IgG Isotype Control (green). After incubation of the primary antibody at +4 $^{\circ}$ C for an hour, the cells were stained with a iFluorTM 488 conjugate-Goat anti-Rabbit IgG Secondary antibody at 1/1,000 dilution for 30 minutes at +4 $^{\circ}$ C. Unlabelled sample was used as a control (cells without incubation with primary antibody; black).





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP3-32033

HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control
H00010189-Q01-10ug	Recombinant Human Aly GST (N-Term) Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-32033

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

