Product Datasheet

Atlastin-3 Antibody (PSH02-47) NBP3-32060

Unit Size: 100 ul

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-32060

Updated 8/7/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP3-32060



NBP3-32060

Atlastin-3 Antibody (PSH02-47)

Aliasiii-5 Aliibouy (1 51 102-41)	
Product Information	
Unit Size	100 ul
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	PSH02-47
Preservative	0.05% Sodium Azide
Isotype	IgG
Purity	Protein A purified
Buffer	PBS (pH7.4), 0.1% BSA and 40% Glycerol
Target Molecular Weight	61 kDa
Product Description	
Host	Rabbit
Gene ID	25923
Gene Symbol	ATL3
Species	Human, Mouse, Rat, Monkey
Immunogen	Recombinant protein within human Atlastin-3 aa 1-488 / 541. (Uniprot: Q6DD88)
Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry
Recommended Dilutions	Western Blot 1:1000, Flow Cytometry 1:1000, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1:100, Immunohistochemistry-



Paraffin 1:1000

Images

Western Blot: Atlastin-3 Antibody (PSH02-47) [NBP3-32060] - Western blot analysis of Atlastin-3 on different lysates with Rabbit anti-Atlastin-3 antibody (NBP3-32060) at 1/1,000 dilution.

Lane 1: HeLa cell lysate

Lane 2: Jurkat cell lysate

Lane 3: A549 cell lysate

Lane 4: HepG2 cell lysate

Lane 5: COS-1 cell lysate

Lane 6: NIH/3T3 cell lysate

Lane 7: RAW264.7 cell lysate

Lane 8: Neuro-2a cell lysate

Lane 9: PC-12 cell lysate

Lane 10: L6 cell lysate

Lane 11: Mouse liver tissue lysate

Lane 12: Rat liver tissue lysate

Lysates/proteins at 30 ug/Lane.

Predicted band size: 61 kDa Observed band size: 61 kDa


Exposure time: 30 seconds;


4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (NBP3-32060) at 1/1,000 dilution was used in 5% NFDM/TBST at room temperature for 2 hours. Goat Anti-Rabbit IgG - HRP Secondary Antibody at 1/50,000 dilution was used for 1 hour at room temperature.

Immunohistochemistry: Atlastin-3 Antibody (PSH02-47) [NBP3-32060] - Immunohistochemical analysis of paraffin-embedded mouse liver tissue with Rabbit anti-Atlastin-3 antibody (NBP3-32060) at 1/1,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH2O and PBS, and then probed with the primary antibody (NBP3-32060) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.








Immunocytochemistry/ Immunofluorescence: Atlastin-3 Antibody (PSH02 -47) [NBP3-32060] - Immunocytochemistry analysis of HeLa cells labeling Atlastin-3 with Rabbit anti-Atlastin-3 antibody (NBP3-32060) at 1/100 dilution.


Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-Atlastin-3 antibody (NBP3-32060) at 1/100 dilution in 1% BSA in PBST overnight at 4 □. Goat Anti-Rabbit IgG H&L (iFluor™ 488) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.

Beta tubulin (red) was stained at 1/100 dilution overnight at +4□. Goat Anti-Mouse IgG H&L (iFluor™ 594) was used as the secondary antibody at 1/1,000 dilution.

Flow Cytometry: Atlastin-3 Antibody (PSH02-47) [NBP3-32060] - Flow cytometric analysis of HeLa cells labeling Atlastin-3.

Cells were fixed and permeabilized. Then stained with the primary antibody (NBP3-32060, 1µg/mL) (red) compared with Rabbit IgG Isotype Control (green). After incubation of the primary antibody at +4□ for an hour, the cells were stained with a iFluor™ 488 conjugate-Goat anti-Rabbit IgG Secondary antibody at 1/1,000 dilution for 30 minutes at +4□. Unlabelled sample was used as a control (cells without incubation with primary antibody; black).









Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP3-32060

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

H00025923-P01-10ug Recombinant Human Atlastin-3 GST (N-Term) Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-32060

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

