# **Product Datasheet**

# c-Myc Antibody (PS00-11) NBP3-32201

Unit Size: 100 ul

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-32201

Updated 8/7/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-32201



## NBP3-32201

c-Myc Antibody (PS00-11)

o myo rumbody (1 000 11)	
Product Information	
100 ul	
1 mg/ml	
Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.	
Monoclonal	
PS00-11	
0.05% Sodium Azide	
IgG	
Protein A purified	
PBS (pH7.4), 0.1% BSA and 40% Glycerol	
49 kDa	
Product Description	
Rabbit	
4609	
MYC	
Human	
Synthetic peptide. (Uniprot: P01106)	
Product Application Details	
Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry	
Western Blot 1:1000, Immunohistochemistry, Immunohistochemistry-Paraffin 1:200	



#### **Images**

Western Blot: c-Myc Antibody (PS00-11) [NBP3-32201] - Western blot analysis of c-Myc on different lysates with Rabbit anti-c-Myc antibody (NBP3-32201) at 1/1,000 dilution.

Lane 1: Jurkat cell lysate Lane 2: Daudi cell lysate Lane 3: K562 cell lysate Lane 4: HL-60 cell lysate

Lysates/proteins at 10 ug/Lane.

Predicted band size: 49 kDa Observed band size: 55 kDa


Exposure time: 2 minutes;


10% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (NBP3-32201) at 1/1,000 dilution was used in 5% NFDM/TBST at room temperature for 2 hours. Goat Anti-Rabbit IgG - HRP Secondary Antibody at 1:300,000 dilution was used for 1 hour at room temperature.

Immunohistochemistry: c-Myc Antibody (PS00-11) [NBP3-32201] - Immunohistochemical analysis of paraffin-embedded human B lymphocytoma tissue with Mouse anti-c-Myc antibody (NBP3-32201) at 1/200 dilution.

The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) for 2 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH2O and PBS, and then probed with the primary antibody (NBP3-32201) at 1/200 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.









### **Novus Biologicals USA**

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

### **Products Related to NBP3-32201**

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

H00004609-P01-10ug Recombinant Human c-Myc GST (N-Term) Protein

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-32201

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

