

Product Datasheet

Histone H3 [ac Lys4] Antibody (PSH03-71) NBP3-32429

Unit Size: 100 ul

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP3-32429

Updated 8/7/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP3-32429



NBP3-32429

Histone H3 [ac Lys4] Antibody (PSH03-71)


Product Information	
Unit Size	100 ul
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	PSH03-71
Preservative	0.05% Sodium Azide
Isotype	IgG
Purity	Protein A purified
Buffer	PBS (pH7.4), 0.1% BSA and 40% Glycerol
Target Molecular Weight	15 kDa

Product Description	
Host	Rabbit
Gene ID	126961
Gene Symbol	H3C14
Species	Human, Mouse, Rat
Immunogen	Synthetic peptide within human Histone H3 (ac Lys 4) aa 1-50. (Uniprot: P68431)

Product Application Details	
Applications	Western Blot, Chromatin Immunoprecipitation, Dot Blot, Immunocytochemistry/Immunofluorescence
Recommended Dilutions	Western Blot 1:1000, Chromatin Immunoprecipitation Use 0.5-2 ug for 25 ug of chromatin, Immunocytochemistry/ Immunofluorescence 1:500-1:1000, Dot Blot 1:1000

Images

Chromatin Immunoprecipitation: Histone H3 [ac Lys4] Antibody (PSH03-71) [NBP3-32429] - Chromatin immunoprecipitations were performed with cross-linked chromatin from HeLa cells treated 500ng/mL TSA for 4 hours and either Histone H3 (NBP3-32429) or Normal Rabbit IgG according to the ChIP protocol. The enriched DNA was quantified by real-time PCR using indicated primers. The amount of immunoprecipitated DNA in each sample is represented as signal relative to the total amount of input chromatin, which is equivalent to one.

ChIP: Acetyl-Histone H3 (Lys4)

Dot Blot: Histone H3 [ac Lys4] Antibody (PSH03-71) [NBP3-32429] - Dot blot analysis of Histone H3 on different proteins with Rabbit anti-Histone H3 antibody (NBP3-32429) at 1/1,000 dilution. Goat Anti-Rabbit IgG - HRP Secondary Antibody at 1/50,000 dilution for 1 hour at room temperature.

Lane 1: Unmodified Histone H3 (negative)
 Lane 2: Mono-Methyl-Histone H3 (Lys4) (negative)
 Lane 3: Di-Methyl-Histone H3 (Lys4) (negative)
 Lane 4: Tri-Methyl-Histone H3 (Lys4) (negative)
 Lane 5: Acetyl-Histone H3 (Lys4) (positive)
 Lane 6: Acetyl-Histone H3 (Lys9) (negative)
 Lane 7: Acetyl-Histone H3 (Lys27) (negative)

Proteins loading: 100ng, 25ng, 5ng;



Blocking and dilution buffer: 5% NFDm/TBST;

Exposure time: 50 seconds.

Immunocytochemistry/ Immunofluorescence: Histone H3 [ac Lys4] Antibody (PSH03-71) [NBP3-32429] - Immunocytochemistry analysis of HeLa cells treated with or without 500ng/mL TSA for 4 hours labeling Histone H3 with Rabbit anti-Histone H3 antibody (NBP3-32429) at 1/500 dilution.

Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-Histone H3 antibody (NBP3-32429) at 1/500 dilution in 1% BSA in PBST overnight at 4 °C. Goat Anti-Rabbit IgG H&L (iFluor™ 488) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.

Beta tubulin (red) was stained at 1/100 dilution overnight at +4°C. Goat Anti-Mouse IgG H&L (iFluor™ 594) was used as the secondary antibody at 1/1,000 dilution.



Western Blot: Histone H3 [ac Lys4] Antibody (PSH03-71) [NBP3-32429]

- Western blot analysis of Histone H3 on different lysates with Rabbit anti-Histone H3 antibody (NBP3-32429) at 1/1,000 dilution.

Lane 1: HeLa cell lysate

Lane 2: HeLa treated with 500ng/mL TSA for 4 hours cell lysate

Lane 3: NIH/3T3 cell lysate

Lane 4: NIH/3T3 treated with 400nM TSA for 18 hours cell lysate

Lysates/proteins at 20 µg/Lane.


Predicted band size: 15 kDa

Observed band size: 15 kDa

Exposure time: 1 minute 22 seconds;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody at 1/1,000 dilution was used in 5% NFDM/TBST at 4°C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody at 1/50,000 dilution was used for 1 hour at room temperature.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP3-32429

HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control
NB21-1101PEP	Histone H3 [p Thr11] Antibody Blocking Peptide

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-32429

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

