Product Datasheet

Histone H4 [ac Lys8] Antibody (PS01-25) NBP3-32432

Unit Size: 100 ul

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-32432

Updated 8/7/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP3-32432



NBP3-32432

Histone H4 [ac Lys8] Antibody (PS01-25)

misione n4 [ac Lyso] Antibody (P501-25)	
Product Information	
Unit Size	100 ul
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	PS01-25
Preservative	0.05% Sodium Azide
Isotype	IgG
Purity	Protein A purified
Buffer	PBS (pH7.4), 0.1% BSA and 40% Glycerol
Target Molecular Weight	11 kDa
Product Description	
Host	Rabbit
Gene ID	121504
Gene Symbol	H4C16
Species	Human, Mouse, Rat
Immunogen	Synthetic peptide within Human Histone H4 aa 1-20 (ac Lys 8). (Uniprot: P62805)
Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry
Recommended Dilutions	Western Blot 1:1000, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1:50, Immunohistochemistry-Paraffin 1:1000-1:5000



Images

Western Blot: Histone H4 [ac Lys8] Antibody (PS01-25) [NBP3-32432] - Western blot analysis of Histone H4 on different lysates with Rabbit anti-Histone H4 antibody (NBP3-32432) at 1/500 dilution.

Lane 1: Hela cell lysate, 10 ug/Lane Lane 2: NIH/3T3 cell lysate, 10 ug/Lane Lane 3: PC-12 cell lysate, 10 ug/Lane

Predicted band size: 11 kDa Observed band size: 14/16 kDa

Exposure time: 2 minutes;

15% SDS-PAGE gel.


Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (NBP3-32432) at 1/500 dilution was used in 5% NFDM/TBST at room temperature for 2 hours. Goat Anti-Rabbit IgG - HRP Secondary Antibody at 1:300,000 dilution was used for 1 hour at room temperature.


Immunohistochemistry: Histone H4 [ac Lys8] Antibody (PS01-25) [NBP3 -32432] - Immunohistochemical analysis of paraffin-embedded human skin tissue with Rabbit anti-Histone H4 antibody (NBP3-32432) at 1/1,000 dilution.


The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) for 2 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH2O and PBS, and then probed with the primary antibody (NBP3-32432) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.

Immunocytochemistry/ Immunofluorescence: Histone H4 [ac Lys8] Antibody (PS01-25) [NBP3-32432] - Immunocytochemistry analysis of PC-12 cells labeling Histone H4 with Rabbit anti-Histone H4 antibody (NBP3-32432) at 1/50 dilution.

Cells were fixed in 4% paraformaldehyde for 10 minutes at 37 \square , permeabilized with 0.05% Triton X-100 in PBS for 20 minutes, and then blocked with 2% negative goat serum for 30 minutes at room temperature. Cells were then incubated with Rabbit anti-Histone H4 antibody (NBP3-32432) at 1/50 dilution in 2% negative goat serum overnight at 4 \square . Goat Anti-Rabbit IgG H&L (iFluorTM 488) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.











Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP3-32432

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

NB21-2090PEP Histone H4 [Trimethyl Lys20] Antibody Blocking Peptide

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-32432

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

