Product Datasheet

Lunatic Fringe Antibody (JE66-62) NBP3-32530

Unit Size: 100 ul

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-32530

Updated 8/7/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP3-32530



NBP3-32530

Lunatic Fringe Antibody (JE66-62)

Lunatic Filinge Antibody (JE66-62)	
Product Information	
Unit Size	100 ul
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	JE66-62
Preservative	0.05% Sodium Azide
Isotype	IgG
Purity	Protein A purified
Buffer	1*TBS (pH7.4), 0.05% BSA and 40% Glycerol
Target Molecular Weight	42 kDa
Product Description	
Host	Rabbit
Gene ID	3955
Gene Symbol	LFNG
Species	Human, Mouse
Immunogen	Synthetic peptide within Human Lunatic Fringe aa 330-379 / 379. (Uniprot: Q8NES3)
Product Application Details	
Applications	Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence
Recommended Dilutions	Western Blot 1:1000, Flow Cytometry 1:1000, Immunocytochemistry/

Immunofluorescence 1:500



Images

Western Blot: Lunatic Fringe Antibody (JE66-62) [NBP3-32530] - Western blot analysis of Lunatic Fringe on different lysates with Rabbit anti-Lunatic Fringe antibody (NBP3-32530) at 1/1,000 dilution.

Lane 1: Hela cell lysate (15 ug/Lane)

Lane 2: K-562 cell lysate (15 ug/Lane)

Lane 3: HL-60 cell lysate (15 ug/Lane)

Lane 4: MCF7 cell lysate (15 ug/Lane)

Lane 5: HT-29 cell lysate (15 ug/Lane)

Lane 6: C2C12 cell lysate (15 ug/Lane)

Lane 7: PC-12 cell lysate (15 ug/Lane)

Lane 8: Mouse testis tissue lysate (30 ug/Lane)

Predicted band size: 42 kDa Observed band size: 42 kDa

Exposure time: 7 seconds;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (NBP3-32530) at 1/1,000 dilution was used in 5% NFDM/TBST at room temperature for 2 hours. Goat Anti-Rabbit IgG - HRP Secondary Antibody at 1:100,000 dilution was used for 1 hour at room temperature.


Immunocytochemistry/ Immunofluorescence: Lunatic Fringe Antibody (JE66-62) [NBP3-32530] - Immunocytochemistry analysis of HUVEC cells labeling Lunatic Fringe with Rabbit anti-Lunatic Fringe antibody (NBP3-32530) at 1/500 dilution.


Cells were fixed in 4% paraformaldehyde for 10 minutes at 37 □, permeabilized with 0.05% Triton X-100 in PBS for 20 minutes, and then blocked with 2% negative goat serum for 30 minutes at room temperature. Cells were then incubated with Rabbit anti-Lunatic Fringe antibody (NBP3-32530) at 1/200 dilution in 2% negative goat serum overnight at 4 □. Goat Anti-Rabbit IgG H&L (iFluor™ 488) was used as the secondary antibody at 1/1,000 dilution. Nuclear DNA was labelled in blue with DAPI.


Beta tubulin (red) was stained at 1/200 dilution overnight at +4□. Goat Anti-Mouse IgG H&L (iFluor™ 594) was used as the secondary antibody at 1/1,000 dilution.

Flow Cytometry: Lunatic Fringe Antibody (JE66-62) [NBP3-32530] - Flow cytometric analysis of K-562 cells labeling Lunatic Fringe.

Cells were fixed and permeabilized. Then stained with the primary antibody (NBP3-32530, 1ug/ml) (red) compared with Rabbit IgG Isotype Control (green). After incubation of the primary antibody at +4□ for an hour, the cells were stained with a iFluor™ 488 conjugate-Goat anti-Rabbit IgG Secondary antibody at 1/1,000 dilution for 30 minutes at +4□. Unlabelled sample was used as a control (cells without incubation with primary antibody; black).











Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP3-32530

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

H00003955-P01-10ug Recombinant Human Lunatic Fringe GST (N-Term) Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-32530

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

