Product Datasheet

PKA C (pan) [p Thr197] Antibody (JE45-17) - (Catalytic Subunit) NBP3-32788

Unit Size: 100 ul

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-32788

Updated 8/7/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-32788



NBP3-32788

PKA C (pan) [p Thr197] Antibody (JE45-17) - (Catalytic Subunit)

PKA C (pan) [p Thr197] Antibody (JE45-17) - (Catalytic Subunit)	
Product Information	
Unit Size	100 ul
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	JE45-17
Preservative	0.05% Sodium Azide
Isotype	IgG
Purity	Protein A purified
Buffer	1*TBS (pH7.4), 0.05% BSA and 40% Glycerol
Target Molecular Weight	40 kDa
Product Description	
Host	Rabbit
Gene ID	5566
Gene Symbol	PRKACA
Species	Human, Mouse, Rat
Immunogen	Synthetic phosphopeptide corresponding to residues surrounding Thr197 of PKA C (pan). (Uniprot: P17612)
Product Application Details	
Applications	Western Blot
Recommended Dilutions	Western Blot 1:1000



Images

Western Blot: PKA C (pan) [p Thr197] Antibody (JE45-17) - (Catalytic Subunit) [NBP3-32788] - Western blot analysis of PKA C (pan) on different lysates with Rabbit anti-PKA C (pan) antibody (NBP3-32788) at 1/1,000 dilution.

Lane 1: NIH/3T3 cell lysate Lane 2: C6 cell lysate Lane 3: HeLa cell lysate

Lane 4: HeLa serum starved overnight then treated with 100nM

Calyculin A for 1 hour cell lysate


Lysates/proteins at 20 ug/Lane.

Predicted band size: 40 kDa Observed band size: 40 kDa

Exposure time: 46 seconds;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (NBP3-32788) at 1/1,000 dilution was used in 5% NFDM/TBST at 4C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody at 1/50,000 dilution was used for 1 hour at room temperature.







Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP3-32788

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

236-EG-200 EGF [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-32788

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

