# **Product Datasheet**

# VASP [p Ser156] Antibody (JE45-91) NBP3-32815

Unit Size: 100 ul

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-32815

Updated 8/7/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-32815



### NBP3-32815

VASP [p Ser156] Antibody (JE45-91)

VASI [p Seriso] Antibody (3E45-91)	
Product Information	
Unit Size	100 ul
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	JE45-91
Preservative	0.05% Sodium Azide
Isotype	IgG
Purity	Protein A purified
Buffer	1*TBS (pH7.4), 0.05% BSA and 40% Glycerol
Target Molecular Weight	40 kDa
Product Description	
Host	Rabbit
Gene ID	7408
Gene Symbol	VASP
Species	Human, Mouse, Rat
Immunogen	Synthetic phosphopeptide corresponding to residues surrounding Ser156 of human VASP. (Uniprot: P50552)
Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry
Recommended Dilutions	Western Blot 1:1000, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1:100, Immunohistochemistry-Paraffin 1:1000



#### **Images**

Western Blot: VASP [p Ser156] Antibody (JE45-91) [NBP3-32815] - Western blot analysis of VASP on different lysates with Rabbit anti-VASP antibody (NBP3-32815) at 1/1,000 dilution.

Lane 1: HeLa cell lysate

Lane 2: HeLa treated with 100ng/mL Calyculin A for 30 minutes cell

iysate

Lane 3: HeLa treated with 100ng/mL Calyculin A for 30 minutes cell lysate, then the membrane treated with pp for 1 hour

Lysates/proteins at 20 ug/Lane.

Predicted band size: 40 kDa Observed band size: 50 kDa

Exposure time: 3 minutes; ECL;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (NBP3-32815) at 1/1,000 dilution was used in 5% NFDM/TBST at 4C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody at 1/50,000 dilution was used for 1 hour at room temperature.


Immunohistochemistry: VASP [p Ser156] Antibody (JE45-91) [NBP3-32815] - Immunohistochemical analysis of paraffin-embedded mouse spleen tissue with Rabbit anti-VASP antibody (NBP3-32815) at 1/1,000 dilution.


The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH2O and PBS, and then probed with the primary antibody (NBP3-32815) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.


Immunocytochemistry/ Immunofluorescence: VASP [p Ser156] Antibody (JE45-91) [NBP3-32815] - Immunocytochemistry analysis of A431 cells treated with or without 10µM forskolin for 1 hour labeling VASP with Rabbit anti-VASP antibody (NBP3-32815) at 1/100 dilution.

Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-VASP antibody (NBP3-32815) at 1/100 dilution in 1% BSA in PBST overnight at 4 □. Goat Anti-Rabbit IgG H&L (iFluor™ 488) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.

Beta tubulin (red) was stained at 1/100 dilution overnight at  $+4\square$ . Goat Anti-Mouse IgG H&L (iFluor<sup>TM</sup> 594) was used as the secondary antibody at 1/1,000 dilution.











### **Novus Biologicals USA**

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

#### **Products Related to NBP3-32815**

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

NBP1-51073-0.05mg Recombinant Human VASP His Protein

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-32815

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

