# **Product Datasheet**

## PSMC6 Antibody (PSH0-12) NBP3-32866

Unit Size: 100 ul

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-32866

Updated 8/7/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-32866



## NBP3-32866

PSMC6 Antibody (PSH0-12)

| 1 SMC0 Antibody (1 SH0-12)  |   |
|-----------------------------|---|
| Product Information         |   |
| Unit Size                   | 100 ul  |
| Concentration               | 1 mg/ml   |
| Storage                     | Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.                      |
| Clonality                   | Monoclonal  |
| Clone                       | PSH0-12   |
| Preservative                | 0.05% Sodium Azide  |
| Isotype                     | IgG   |
| Purity                      | Protein A purified  |
| Buffer                      | PBS (pH7.4), 0.1% BSA and 40% Glycerol  |
| Target Molecular Weight     | 44 kDa  |
| Product Description         |   |
| Host                        | Rabbit  |
| Gene ID                     | 5706  |
| Gene Symbol                 | PSMC6   |
| Species                     | Human, Mouse, Rat   |
| Immunogen                   | Recombinant protein within human PSMC6 aa 1-200 / 389. (Uniprot: P62333)                                    |
| Product Application Details |   |
| Applications                | Western Blot, Immunohistochemistry-Paraffin, Flow Cytometry, Immunohistochemistry                           |
| Recommended Dilutions       | Western Blot 1:1000, Flow Cytometry 1:500-1:1000, Immunohistochemistry, Immunohistochemistry-Paraffin 1:400 |
|                             |   |



#### **Images**

Western Blot: PSMC6 Antibody (PSH0-12) [NBP3-32866] - Western blot analysis of PSMC6 on different lysates with Rabbit anti-PSMC6 antibody (NBP3-32866) at 1/1,000 dilution.

Lane 1: Hela cell lysate (10 ug/Lane)

Lane 2: 293T cell lysate (10 ug/Lane)

Lane 3: SH-SY5Y cell lysate (10 ug/Lane)

Lane 4: HepG2 cell lysate (10 ug/Lane)

Lane 5: PC-12 cell lysate (10 ug/Lane)

Lane 6: NIH/3T3 cell lysate (10 ug/Lane)

Lane 7: Mouse kidney tissue lysate (20 ug/Lane)

Lane 8: Rat kidney tissue lysate (20 ug/Lane)

Lane 9: Rat brain tissue lysate (20 ug/Lane)

Predicted band size: 44 kDa Observed band size: 44 kDa

Exposure time: 2 minutes;

10% SDS-PAGE gel.


Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (NBP3-32866) at 1/1,000 dilution was used in 5% NFDM/TBST at room temperature for 2 hours. Goat Anti-Rabbit IgG - HRP Secondary Antibody at 1:300,000 dilution was used for 1 hour at room temperature.


Immunohistochemistry: PSMC6 Antibody (PSH0-12) [NBP3-32866] - Immunohistochemical analysis of paraffin-embedded human testis tissue with Rabbit anti-PSMC6 antibody (NBP3-32866) at 1/400 dilution.


The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) for 2 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH2O and PBS, and then probed with the primary antibody (NBP3-32866) at 1/400 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.

Flow Cytometry: PSMC6 Antibody (PSH0-12) [NBP3-32866] - Flow cytometric analysis of SH-SY5Y cells labeling PSMC6.

Cells were fixed and permeabilized. Then stained with the primary antibody (NBP3-32866, 1ug/ml) (red) compared with Rabbit IgG Isotype Control (green). After incubation of the primary antibody at +4□ for an hour, the cells were stained with a iFluor<sup>™</sup> 488 conjugate-Goat anti-Rabbit IgG Secondary antibody at 1/1,000 dilution for 30 minutes at +4□. Unlabelled sample was used as a control (cells without incubation with primary antibody; black).











### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

### **Products Related to NBP3-32866**

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

H00005706-P01-10ug Recombinant Human PSMC6 GST (N-Term) Protein

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-32866

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

