Product Datasheet

TIMM22 Antibody (JE66-67) NBP3-33021

Unit Size: 100 ul

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-33021

Updated 8/7/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP3-33021



NBP3-33021

TIMM22 Antibody (JE66-67)

TIIVIIVIZZ Antibody (JE66-67)	
Product Information	
Unit Size	100 ul
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	JE66-67
Preservative	0.05% Sodium Azide
Isotype	IgG
Purity	Protein A purified
Buffer	1*TBS (pH7.4), 0.05% BSA and 40% Glycerol
Target Molecular Weight	20 kDa
Product Description	
Host	Rabbit
Gene ID	29928
Gene Symbol	TIMM22
Species	Human, Mouse, Rat
Immunogen	Recombinant protein within Human TIMM22 aa 95-194 / 194. (Uniprot: Q9Y584)
Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry
Recommended Dilutions	Western Blot 1:1000, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1:100, Immunohistochemistry-Paraffin 1:200-1:1000



Images

Western Blot: TIMM22 Antibody (JE66-67) [NBP3-33021] - Western blot analysis of TIMM22 on different lysates with Rabbit anti-TIMM22 antibody (NBP3-33021) at 1/1,000 dilution.

Lane 1: K-562 cell lysate (15 ug/Lane) Lane 2: HepG2 cell lysate (15 ug/Lane)

Lane 3: A549 cell lysate (15 ug/Lane)
Lane 4: MCF7 cell lysate (15 ug/Lane)

Lane 5: HEK-293 cell lysate (15 ug/Lane)

Lane 6: HeLa cell lysate (15 ug/Lane) Lane 7: Jurkat cell lysate (15 ug/Lane)

Lane 8: Mouse testis tissue lysate (30 ug/Lane)

Predicted band size: 20 kDa Observed band size: 20 kDa

Exposure time: 3 minutes;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (NBP3-33021) at 1/1,000 dilution was used in 5% NFDM/TBST at room temperature for 2 hours. Goat Anti-Rabbit IgG - HRP Secondary Antibody at 1:100,000 dilution was used for 1 hour at room temperature.


Immunohistochemistry: TIMM22 Antibody (JE66-67) [NBP3-33021] - Immunohistochemical analysis of paraffin-embedded human testis tissue with Rabbit anti-TIMM22 antibody (NBP3-33021) at 1/1,000 dilution.


The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH2O and PBS, and then probed with the primary antibody (NBP3-33021) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.


Immunocytochemistry/ Immunofluorescence: TIMM22 Antibody (JE66-67) [NBP3-33021] - Immunocytochemistry analysis of K-562 cells labeling TIMM22 with Rabbit anti-TIMM22 antibody (NBP3-33021) at 1/100 dilution.

Cells were fixed in 4% paraformaldehyde for 10 minutes at 37 □, permeabilized with 0.05% Triton X-100 in PBS for 20 minutes, and then blocked with 2% negative goat serum for 30 minutes at room temperature. Cells were then incubated with Rabbit anti-TIMM22 antibody (NBP3-33021) at 1/100 dilution in 2% negative goat serum overnight at 4 □. Goat Anti-Rabbit IgG H&L (iFluor™ 488) was used as the secondary antibody at 1/1,000 dilution. Nuclear DNA was labelled in blue with DAPI.

Beta tubulin (red) was stained at 1/200 dilution overnight at +4□. Goat Anti-Mouse IgG H&L (iFluor™ 594) was used as the secondary antibody at 1/1,000 dilution."











Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP3-33021

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

H00029928-P01-10ug Recombinant Human TIMM22 GST (N-Term) Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-33021

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

