## **Product Datasheet**

# TRF-2 Antibody (JE37-57) NBP3-33049

Unit Size: 100 ul

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-33049

Updated 8/7/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-33049



## NBP3-33049

TRF-2 Antibody (JE37-57)

| TRF-2 Aniibody (JE37-37)    |   |
|-----------------------------|---|
| Product Information         |   |
| Unit Size                   | 100 ul  |
| Concentration               | 1 mg/ml   |
| Storage                     | Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.  |
| Clonality                   | Monoclonal  |
| Clone                       | JE37-57   |
| Preservative                | 0.05% Sodium Azide  |
| Isotype                     | IgG   |
| Purity                      | Protein A purified  |
| Buffer                      | 1*TBS (pH7.4), 0.05% BSA and 40% Glycerol   |
| Target Molecular Weight     | 60 kDa  |
| Product Description         |   |
| Host                        | Rabbit  |
| Gene ID                     | 7014  |
| Gene Symbol                 | TERF2   |
| Species                     | Human   |
| Immunogen                   | Recombinant protein within Human TRF-2 aa 1-100 / 542. (Uniprot: Q15554)  |
| Product Application Details |   |
| Applications                | Western Blot, Immunohistochemistry-Paraffin, Immunocytochemistry/<br>Immunofluorescence, Immunohistochemistry                             |
| Recommended Dilutions       | Western Blot 1:1000, Immunohistochemistry, Immunocytochemistry/<br>Immunofluorescence 20ug/mL, Immunohistochemistry-Paraffin 1:200-1:1000 |
| 4                           |   |



#### **Images**

Western Blot: TRF-2 Antibody (JE37-57) [NBP3-33049] - Western blot analysis of TRF-2 on different lysates with Rabbit anti-TRF-2 antibody (NBP3-33049) at 1/500 dilution.

Lane 1: SiHa cell lysate Lane 2: Jurkat cell lysate

Lysates/proteins at 10 ug/Lane.

Predicted band size: 60 kDa Observed band size: 60 kDa

Exposure time: 2 minutes;

10% SDS-PAGE gel.


Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (NBP3-33049) at 1/500 dilution was used in 5% NFDM/TBST at room temperature for 2 hours. Goat Anti-Rabbit IgG - HRP Secondary Antibody at 1:200,000 dilution was used for 1 hour at room temperature.


Immunohistochemistry: TRF-2 Antibody (JE37-57) [NBP3-33049] - Immunohistochemical analysis of paraffin-embedded human colon carcinoma tissue with Rabbit anti-TRF-2 antibody (NBP3-33049) at 1/1,000 dilution.


The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) for 2 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH2O and PBS, and then probed with the primary antibody (NBP3-33049) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.

Immunocytochemistry/ Immunofluorescence: TRF-2 Antibody (JE37-57) [NBP3-33049] - Immunocytochemistry analysis of SiHa cells labeling TRF-2 with Rabbit anti-TRF-2 antibody (NBP3-33049) at 20ug/mL dilution.

Cells were fixed in 4% paraformaldehyde for 10 minutes at 37 □, permeabilized with 0.05% Triton X-100 in PBS for 20 minutes, and then blocked with 2% negative goat serum for 30 minutes at room temperature. Cells were then incubated with Rabbit anti-TRF-2 antibody (NBP3-33049) at 20ug/mL dilution in 2% negative goat serum overnight at 4 □. Goat Anti-Rabbit IgG H&L (iFluor™ 488) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.











### **Novus Biologicals USA**

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

### **Products Related to NBP3-33049**

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

NBP1-86911PEP TRF-2 Recombinant Protein Antigen

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-33049

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

